prev_injuries ~ age+minutes_played+injured

0.996 + (−0.007)*age + (0)*minutes_played + (0.675)*injured

A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05, High R and Adj R,
Low AIC

F−statistic: 12.48142 on 3 and 616 Degrees of Freedom


F p−value: 0


R−Squared: 0.0573


Adjusted R−Squared: 0.0527118358989821


Root MSE: 1.05223721049096


AIC: 1832.62281312613



prev_injuries against age
prev_injuries
0
1
2
3
4
20
25
30
35
age



prev_injuries against minutes_played
prev_injuries
minutes_played
0
1
2
3
4
0
1000
2000
3000
4000



prev_injuries against injured
prev_injuries
injured
0
1
2
3
4
0.0
0.2
0.4
0.6
0.8
1.0



Residuals vs Fitted
Residuals
3
2
1
0
−1
−2
0.8
1.0
1.2
1.4
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
1294
608
613
−3
−2
−1
0
1
2
3


Scale−Location
√|Standardized residuals|
0.0
0.5
1.0
1.5
0.8
1.0
1.2
1.4
Fitted values
lm(formula)


Cook's distance
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
97
256
613
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
613
256
97
Cook's distance
3
2
1
0
−1
−2
0.000
0.005
0.010
0.015
0.020
0.025
0.030


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3.5
3
2.5
2
256
97
613
1.5
1
0.5
0
Cook's distance
0.025
0.020
0.015
0.010
0.005
0.000
0
0.005
0.01
0.015
0.02
0.025
0.03
Leverage h_{ii}
lm(formula)