


prev_injuries ~ age+minutes_played+training_load
0.683 + (0.002)*age + (0)*minutes_played + (0.07)*training_load


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 0.8592 on 3 and 616 Degrees of Freedom
F p−value: 0.46201
R−Squared: 0.00417
Adjusted R−Squared: −0.000682856678209776
Root MSE: 1.08148582144189
AIC: 1866.62024291008



prev_injuries against age
prev_injuries
0
1
2
3
4
20
25
30
35
age



prev_injuries against minutes_played
prev_injuries
minutes_played
0
1
2
3
4
0
1000
2000
3000
4000



prev_injuries against training_load
prev_injuries
training_load
0
1
2
3
4
1
2
3
4
5



Residuals vs Fitted
256
96
129
Residuals
3
2
1
0
−1
0.75
0.80
0.85
0.90
0.95
1.00
1.05
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
256
96
−3
−2
−1
0
1
2
3


Scale−Location
256
96
129
√|Standardized residuals|
0.0
0.5
1.0
1.5
0.75
0.80
0.85
0.90
0.95
1.00
1.05
Fitted values
lm(formula)


Cook's distance
256
115
613
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
3
2
1
0
−1
256
613
115
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
2
1.5
1
0.5
0
256
613
115
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0
0.005
0.01
0.015
0.02
0.025
Leverage h_{ii}
lm(formula)