


prev_injuries ~ age+minutes_played
0.824 + (0.003)*age + (0)*minutes_played


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 0.04396 on 2 and 617 Degrees of Freedom
F p−value: 0.957
R−Squared: 0.00014
Adjusted R−Squared: −0.00309855313660146
Root MSE: 1.08366894584711
AIC: 1867.12082644414



prev_injuries against age
prev_injuries
0
1
2
3
4
20
25
30
35
age



prev_injuries against minutes_played
prev_injuries
minutes_played
0
1
2
3
4
0
1000
2000
3000
4000



Residuals vs Fitted
256
439
515
Residuals
3
2
1
0
−1
0.88
0.89
0.90
0.91
0.92
0.93
0.94
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−3
−2
−1
0
1
2
3
−1
0
1
2
3
256


Scale−Location
256
439
515
√|Standardized residuals|
0.0
0.5
1.0
1.5
0.88
0.89
0.90
0.91
0.92
0.93
0.94
Fitted values
lm(formula)


Cook's distance
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
256
97
613
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
3
2
1
0
−1
97
613
256
Cook's distance
0.000
0.005
0.010
0.015
0.020
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
2
256
613
97
Cook's distance
0.030
0.025
0.020
0.015
0.010
0.005
0.000
1.5
1
0.5
0
0
0.005
0.01
0.015
0.02
Leverage h_{ii}
lm(formula)