


prev_injuries ~ age+training_load+bmi
0.497 + (0.002)*age + (0.067)*training_load + (0.006)*bmi


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 0.82799 on 3 and 616 Degrees of Freedom
F p−value: 0.47874
R−Squared: 0.00402
Adjusted R−Squared: −0.000834358286828651
Root MSE: 1.08156768586051
AIC: 1866.71410270486



prev_injuries against age
prev_injuries
0
1
2
3
4
20
25
30
35
age



prev_injuries against training_load
prev_injuries
training_load
0
1
2
3
4
1
2
3
4
5



prev_injuries against bmi
prev_injuries
bmi
0
1
2
3
4
20
22
24
26
28



Residuals vs Fitted
2596
129
Residuals
3
2
1
0
−1
0.75
0.80
0.85
0.90
0.95
1.00
1.05
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
129
96
256
−3
−2
−1
0
1
2
3


Scale−Location
√|Standardized residuals|
Fitted values
lm(formula)
0.0
0.5
1.0
1.5
0.75
0.80
0.85
0.90
0.95
1.00
1.05
256
96
129


Cook's distance
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
129
256
271
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
271
129
256
3
2
1
0
−1
0.000
0.005
0.010
0.015
0.020
0.025


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
2
256
129
271
Cook's distance
0.000
0.005
0.010
0.015
0.020
1.5
1
0.5
0
0
0.005
0.01
0.015
0.02
0.025
Leverage h_{ii}
lm(formula)