prev_injuries ~ age+training_load+injured

0.849 + (−0.007)*age + (0.034)*training_load + (0.662)*injured

A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05, High R and Adj R,
Low AIC

F−statistic: 12.67616 on 3 and 616 Degrees of Freedom


F p−value: 0


R−Squared: 0.05814


Adjusted R−Squared: 0.0535580311604819


Root MSE: 1.0517671333232


AIC: 1832.06873090721



prev_injuries against age
prev_injuries
0
1
2
3
4
20
25
30
35
age



prev_injuries against training_load
prev_injuries
training_load
0
1
2
3
4
1
2
3
4
5



prev_injuries against injured
prev_injuries
injured
0
1
2
3
4
0.0
0.2
0.4
0.6
0.8
1.0



Residuals vs Fitted
129
Residuals
3
2
1
0
−1
−2
0.8
1.0
1.2
1.4
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
129
96
408
−3
−2
−1
0
1
2
3


Scale−Location
129
√|Standardized residuals|
0.0
0.5
1.0
1.5
0.8
1.0
1.2
1.4
Fitted values
lm(formula)


Cook's distance
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
256
115
129
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
129
115
256
Standardized residuals
3
2
1
0
−1
−2
Cook's distance
0.000
0.005
0.010
0.015
0.020
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3.5
3
2.5
256
115
129
2
1.5
1
0.5
0
0.025
0.020
0.015
0.010
0.005
0.000
Cook's distance
0
0.005
0.01
0.015
0.02
Leverage h_{ii}
lm(formula)