prev_injuries ~ injured

0.78 + (0.663)*injured

A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05, High R and Adj R,
Low AIC

F−statistic: 37.0709 on 1 and 618 Degrees of Freedom


F p−value: 0


R−Squared: 0.05659


Adjusted R−Squared: 0.0550641158986732


Root MSE: 1.05263462773458


AIC: 1829.09105768049



prev_injuries against injured
prev_injuries
injured
0
1
2
3
4
0.0
0.2
0.4
0.6
0.8
1.0



Residuals vs Fitted
Residuals
Fitted values
lm(formula)
3
2
1
0
−1
−2
0.8
0.9
1.0
1.1
1.2
1.3
1.4


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−3
−2
−1
0
1
2
3


Scale−Location
√|Standardized residuals|
0.0
0.5
1.0
1.5
0.8
0.9
1.0
1.1
1.2
1.3
1.4
Fitted values
lm(formula)


Cook's distance
34
97
181
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
3
2
1
0
−1
Cook's distance
0.000
0.002
0.004
0.006
0.008
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3.5
3
2.5
2
1.5
1
0.5
0
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.001
0.002
0.003
0.004
0.005
0.006
0.007
0.008
0.009
Leverage h_{ii}
lm(formula)