


prev_injuries ~ minutes_played+bmi+injured
0.617 + (0)*minutes_played + (0.009)*bmi + (0.666)*injured


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 12.38232 on 3 and 616 Degrees of Freedom
F p−value: 0
R−Squared: 0.05687
Adjusted R−Squared: 0.0522806725700607
Root MSE: 1.052476648976
AIC: 1832.90494527572



prev_injuries against minutes_played
prev_injuries
minutes_played
0
1
2
3
4
0
1000
2000
3000
4000



prev_injuries against bmi
prev_injuries
bmi
0
1
2
3
4
20
22
24
26
28



prev_injuries against injured
prev_injuries
injured
0
1
2
3
4
0.0
0.2
0.4
0.6
0.8
1.0



Residuals vs Fitted
Residuals
3
2
1
0
−1
−2
0.8
1.0
1.2
1.4
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−3
−2
−1
0
1
2
3
613


Scale−Location
√|Standardized residuals|
0.0
0.5
1.0
1.5
0.8
1.0
1.2
1.4
Fitted values
lm(formula)


Cook's distance
97
271
579
0.025
0.020
0.015
0.010
0.005
0.000
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
3
2
1
0
−1
579
97
271
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3.5
3
2.5
2
271
579
97
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0
0.005
0.01
0.015
0.02
0.025
Leverage h_{ii}
lm(formula)
1.5
1
0.5
0