


prev_injuries ~ minutes_played+bmi
0.82 + (0)*minutes_played + (0.004)*bmi


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 0.00872 on 2 and 617 Degrees of Freedom
F p−value: 0.99132
R−Squared: 3e−05
Adjusted R−Squared: −0.00321314414604634
Root MSE: 1.08373084164577
AIC: 1867.19164936365



prev_injuries against minutes_played
prev_injuries
minutes_played
0
1
2
3
4
0
1000
2000
3000
4000



prev_injuries against bmi
prev_injuries
bmi
0
1
2
3
4
20
22
24
26
28



Residuals vs Fitted
579
534
408
Residuals
3
2
1
0
−1
0.885
0.890
0.895
0.900
0.905
0.910
0.915
0.920
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−1
0
1
2
3
−3
−2
−1
0
1
2
3
579


Scale−Location
579
534
613
√|Standardized residuals|
0.0
0.5
1.0
1.5
0.885
0.890
0.895
0.900
0.905
0.910
0.915
0.920
Fitted values
lm(formula)


Cook's distance
220
271
534
0.025
0.020
0.015
0.010
0.005
0.000
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
534
271
220
3
2
1
0
−1
0.000
0.005
0.010
0.015
0.020
0.025


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
2
534
271
220
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0
0.5
1
1.5
0
0.005
0.01
0.015
0.02
0.025
Leverage h_{ii}
lm(formula)