


prev_injuries ~ minutes_played
0.908 + (0)*minutes_played


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 0.00098 on 1 and 618 Degrees of Freedom
F p−value: 0.97501
R−Squared: 0
Adjusted R−Squared: −0.00161653078053292
Root MSE: 1.08374529119245
AIC: 1865.20818236035



prev_injuries against minutes_played
prev_injuries
minutes_played
0
1
2
3
4
0
1000
2000
3000
4000



Residuals vs Fitted
613
Residuals
3
2
1
0
−1
0.902
0.904
0.906
0.908
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−3
−2
−1
0
1
2
3
579
97
613


Scale−Location
613
579
97
1.5
1.0
0.5
0.0
√|Standardized residuals|
0.902
0.904
0.906
0.908
Fitted values
lm(formula)


Cook's distance
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0.035
97
129
613
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
3
2
1
0
−1
129
97
613
Cook's distance
0.000
0.005
0.010
0.015
0.020
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
2
613
97
129
Cook's distance
0.035
0.030
0.025
0.020
0.015
0.010
0.005
0.000
1.5
1
0.5
0
0
0.005
0.01
0.015
0.02
Leverage h_{ii}
lm(formula)