


prev_injuries ~ training_load+bmi
0.555 + (0.068)*training_load + (0.006)*bmi


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 1.22332 on 2 and 617 Degrees of Freedom
F p−value: 0.29496
R−Squared: 0.00395
Adjusted R−Squared: 0.000721032799220644
Root MSE: 1.08160378762412
AIC: 1864.75549210688



prev_injuries against training_load
prev_injuries
training_load
0
1
2
3
4
1
2
3
4
5



prev_injuries against bmi
prev_injuries
bmi
0
1
2
3
4
20
22
24
26
28



Residuals vs Fitted
96
256
129
Residuals
3
2
1
0
−1
0.75
0.80
0.85
0.90
0.95
1.00
1.05
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−3
−2
−1
0
1
2
3
96


Scale−Location
96
256
129
√|Standardized residuals|
0.0
0.5
1.0
1.5
0.75
0.80
0.85
0.90
0.95
1.00
1.05
Fitted values
lm(formula)


Cook's distance
271
129
534
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
129
534
271
3
2
1
0
−1
0.000
0.005
0.010
0.015
0.020
0.025


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
2
271
129
534
Cook's distance
0.000
0.005
0.010
0.015
0.020
1.5
1
0.5
0
0
0.005
0.01
0.015
0.02
0.025
Leverage h_{ii}
lm(formula)