


prev_injuries ~ training_load
0.688 + (0.067)*training_load


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 2.41267 on 1 and 618 Degrees of Freedom
F p−value: 0.12087
R−Squared: 0.00389
Adjusted R−Squared: 0.00227697640684843
Root MSE: 1.08163686019964
AIC: 1862.79340743885



prev_injuries against training_load
prev_injuries
training_load
0
1
2
3
4
1
2
3
4
5



Residuals vs Fitted
Residuals
3
2
1
0
−1
0.75
0.80
0.85
0.90
0.95
1.00
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−3
−2
−1
0
1
2
3
96
29
256


Scale−Location
√|Standardized residuals|
0.0
0.5
1.0
1.5
0.75
0.80
0.85
0.90
0.95
1.00
Fitted values
lm(formula)


Cook's distance
115
384
96
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
3
2
1
0
−1
96
Cook's distance
0.000
0.002
0.004
0.006
0.008
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
2
1.5
1
0.5
0
96
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0
0.002
0.004
0.006
0.008
Leverage h_{ii}
lm(formula)