


training_load ~ age+injured
2.859 + (0.011)*age + (0.334)*injured


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 6.35496 on 2 and 617 Degrees of Freedom
F p−value: 0.00185
R−Squared: 0.02018
Adjusted R−Squared: 0.0170077070032661
Root MSE: 0.994474585523846
AIC: 1760.61326847991



training_load against age
training_load
age
1
2
3
4
5
20
25
30
35



training_load against injured
training_load
injured
1
2
3
4
5
0.0
0.2
0.4
0.6
0.8
1.0



Residuals vs Fitted
Residuals
Fitted values
lm(formula)
2
1
0
−1
−2
−3
3.1
3.2
3.3
3.4
3.5
3.6
288
101


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−3
−2
−1
0
1
2
3
101
281


Scale−Location
√|Standardized residuals|
Fitted values
lm(formula)
0.0
0.5
1.0
1.5
3.1
3.2
3.3
3.4
3.5
3.6
101


Cook's distance
238
281
101
Cook's distance
0.000
0.005
0.010
0.015
0.020
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
101
288
0.000
0.005
0.010
0.015
2
1
0
−1
−2
−3


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
2.5
2
1.5
1
0.5
0
101
Cook's distance
0.000
0.005
0.010
0.015
0.020
0
0.005
0.01
0.015
Leverage h_{ii}
lm(formula)