

training_load ~ age+minutes_played+bmi+injured


2.595 + (0.012)*age + (0)*minutes_played + (−0.022)*bmi + (0.306)*injured


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05, High R and Adj R,
Low AIC


F−statistic: 10.59952 on 4 and 615 Degrees of Freedom
F p−value: 0
R−Squared: 0.06449
Adjusted R−Squared: 0.0584091522314205
Root MSE: 0.971728012710485
AIC: 1735.92140454175



training_load against age
training_load
age
1
2
3
4
5
20
25
30
35



training_load against minutes_played
training_load
minutes_played
1
2
3
4
5
0
1000
2000
3000
4000



training_load against bmi
training_load
bmi
1
2
3
4
5
20
22
24
26
28



training_load against injured
training_load
injured
1
2
3
4
5
0.0
0.2
0.4
0.6
0.8
1.0



Residuals vs Fitted
Residuals
Fitted values
lm(formula)
2
1
0
−1
−2
−3
2.5
3.0
3.5
107
101
281


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−3
−2
−1
0
1
2
3
281
101
107


Scale−Location
281
101
107
√|Standardized residuals|
0.0
0.5
1.0
1.5
2.5
3.0
3.5
Fitted values
lm(formula)


Cook's distance
101
238
474
0.020
0.015
0.010
0.005
0.000
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
2
1
0
−1
−2
−3
238
474
101


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
2
1.5
1
0.5
0
101
474
238
Cook's distance
0.000
0.005
0.010
0.015
0.020
0
0.005
0.01
0.015
0.02
0.025
0.03
Leverage h_{ii}
lm(formula)