

training_load ~ age+minutes_played+prev_injuries+injured


2.05 + (0.013)*age + (0)*minutes_played + (0.033)*prev_injuries + (0.286)*injured


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05, High R and Adj R,
Low AIC


F−statistic: 10.62787 on 4 and 615 Degrees of Freedom
F p−value: 0
R−Squared: 0.06466
Adjusted R−Squared: 0.0585715533058202
Root MSE: 0.971644209602791
AIC: 1735.81446069187



training_load against age
training_load
age
1
2
3
4
5
20
25
30
35



training_load against minutes_played
training_load
minutes_played
1
2
3
4
5
0
1000
2000
3000
4000



training_load against prev_injuries
training_load
prev_injuries
1
2
3
4
5
0
1
2
3
4



training_load against injured
training_load
injured
1
2
3
4
5
0.0
0.2
0.4
0.6
0.8
1.0



Residuals vs Fitted
Residuals
Fitted values
lm(formula)
2.6
2.8
3.0
3.2
3.4
3.6
3.8
−3
−2
−1
0
1
2
253
281


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−3
−2
−1
0
1
2
3
281


Scale−Location
281
252
253
√|Standardized residuals|
0.0
0.5
1.0
1.5
2.6
2.8
3.0
3.2
3.4
3.6
3.8
Fitted values
lm(formula)


Cook's distance
281
238
115
0.020
0.015
0.010
0.005
0.000
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
115
238
281
2
1
0
−1
−2
−3
0.000
0.005
0.010
0.015
0.020
0.025
0.030


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
2
281
238
115
Cook's distance
0.000
0.005
0.010
0.015
0.020
0
0.5
1
1.5
0
0.005
0.01
0.015
0.02
0.025
0.03
Leverage h_{ii}
lm(formula)