


training_load ~ age+minutes_played
2.004 + (0.017)*age + (0)*minutes_played


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 16.08199 on 2 and 617 Degrees of Freedom
F p−value: 0
R−Squared: 0.04955
Adjusted R−Squared: 0.0464658811784917
Root MSE: 0.979460104992943
AIC: 1741.74910339965



training_load against age
training_load
age
1
2
3
4
5
20
25
30
35



training_load against minutes_played
training_load
minutes_played
1
2
3
4
5
0
1000
2000
3000
4000



Residuals vs Fitted
Residuals
Fitted values
lm(formula)
2
1
0
−1
−2
−3
2.6
2.8
3.0
3.2
3.4
3.6
107
4232


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
252
413
107
−3
−2
−1
0
1
2
3


Scale−Location
√|Standardized residuals|
0.0
0.5
1.0
1.5
2.6
2.8
3.0
3.2
3.4
3.6
Fitted values
lm(formula)


Cook's distance
43
353
413
0.020
0.015
0.010
0.005
0.000
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
353
43
413
2
1
0
−1
−2
−3
0.000
0.005
0.010
0.015
0.020


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
2.5
2
1.5
1
0.5
0
43
353
413
Cook's distance
0.000
0.005
0.010
0.015
0
0.005
0.01
0.015
0.02
Leverage h_{ii}
lm(formula)