


training_load ~ age+prev_injuries+bmi
3.395 + (0.016)*age + (0.057)*prev_injuries + (−0.027)*bmi


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 1.93804 on 3 and 616 Degrees of Freedom
F p−value: 0.12221
R−Squared: 0.00935
Adjusted R−Squared: 0.00452564159994839
Root MSE: 0.999957277600994
AIC: 1769.43080426737



training_load against age
training_load
age
1
2
3
4
5
20
25
30
35



training_load against prev_injuries
training_load
prev_injuries
1
2
3
4
5
0
1
2
3
4



training_load against bmi
training_load
bmi
1
2
3
4
5
20
22
24
26
28



Residuals vs Fitted
Residuals
2
1
0
−1
−2
3.0
3.1
3.2
3.3
3.4
Fitted values
lm(formula)
474


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
466
413
474
−3
−2
−1
0
1
2
3
−2
−1
0
1
2


Scale−Location
√|Standardized residuals|
0.0
0.5
1.0
1.5
474
3.0
3.1
3.2
3.3
3.4
Fitted values
lm(formula)


Cook's distance
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
43
474
584
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
2
1
0
−1
−2
0.000
0.005
0.010
0.015
0.020
Leverage
lm(formula)
584
43
474
Cook's distance


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
2.5
2
474
584
43
1.5
1
0.5
0
0.020
0.015
0.010
0.005
0.000
Cook's distance
0
0.005
0.01
0.015
0.02
Leverage h_{ii}
lm(formula)