training_load ~ age+prev_injuries+injured

2.83 + (0.011)*age + (0.031)*prev_injuries + (0.313)*injured

A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05, High R and Adj R,
Low AIC

F−statistic: 4.44985 on 3 and 616 Degrees of Freedom


F p−value: 0.00419


R−Squared: 0.02121


Adjusted R−Squared: 0.0164448278280379


Root MSE: 0.993952819123151


AIC: 1761.96251266131



training_load against age
training_load
age
1
2
3
4
5
20
25
30
35



training_load against prev_injuries
training_load
prev_injuries
1
2
3
4
5
0
1
2
3
4



training_load against injured
training_load
injured
1
2
3
4
5
0.0
0.2
0.4
0.6
0.8
1.0



Residuals vs Fitted
Residuals
Fitted values
lm(formula)
2
1
0
−1
−2
3.1
3.2
3.3
3.4
3.5
3.6
288
101


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−3
−2
−1
0
1
2
3
101
281


Scale−Location
√|Standardized residuals|
0.0
0.5
1.0
1.5
3.1
3.2
3.3
3.4
3.5
3.6
Fitted values
lm(formula)


Cook's distance
238
281
101
0.020
0.015
0.010
0.005
0.000
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
101
288
0.000
0.005
0.010
0.015
0.020
2
1
0
−1
−2
−3


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
2.5
2
1.5
1
0.5
0
288
101
Cook's distance
0.000
0.005
0.010
0.015
0.020
0
0.005
0.01
0.015
0.02
Leverage h_{ii}
lm(formula)