


training_load ~ age+prev_injuries
2.749 + (0.016)*age + (0.057)*prev_injuries


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 2.38673 on 2 and 617 Degrees of Freedom
F p−value: 0.09278
R−Squared: 0.00768
Adjusted R−Squared: 0.00446056271324302
Root MSE: 1.00080131406128
AIC: 1768.47701271771



training_load against age
training_load
age
1
2
3
4
5
20
25
30
35



training_load against prev_injuries
training_load
prev_injuries
1
2
3
4
5
0
1
2
3
4



Residuals vs Fitted
Residuals
Fitted values
lm(formula)
2
1
0
−1
−2
3.1
3.2
3.3
3.4
43
46
613


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
413
466
43
−3
−2
−1
0
1
2
3
−2
−1
0
1
2


Scale−Location
√|Standardized residuals|
0.0
0.5
1.0
1.5
3.1
3.2
3.3
3.4
Fitted values
lm(formula)
43
46
413


Cook's distance
413
384
115
0.015
0.010
0.005
0.000
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
2
1
0
−1
−2
384
115
413
Cook's distance
0.000
0.005
0.010
0.015
0.020
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
2.5
2
1.5
413
384
115
0.015
0.010
0.005
0.000
Cook's distance
1
0.5
0
0
0.005
0.01
0.015
0.02
Leverage h_{ii}
lm(formula)