


training_load ~ age
2.796 + (0.016)*age


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 2.40695 on 1 and 618 Degrees of Freedom
F p−value: 0.12131
R−Squared: 0.00388
Adjusted R−Squared: 0.00226778786503568
Root MSE: 1.00271447625163
AIC: 1768.84517157981



training_load against age
training_load
age
1
2
3
4
5
20
25
30
35



Residuals vs Fitted
Residuals
Fitted values
lm(formula)
2
1
0
−1
−2
3.10
3.15
3.20
3.25
3.30
3.35
413


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
413
43
252
−3
−2
−1
0
1
2
3


Scale−Location
√|Standardized residuals|
Fitted values
lm(formula)
0.0
0.5
1.0
1.5
3.10
3.15
3.20
3.25
3.30
3.35
413


Cook's distance
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
156
381
413
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
2
1
0
−1
−2
413
Cook's distance
0.000
0.005
0.010
0.015
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
2.5
2
413
386
1.5
1
0.5
0
Cook's distance
0.000
0.005
0.010
0.015
0.020
0
0.002
0.004
0.006
0.008
0.01
0.012
0.014
Leverage h_{ii}
lm(formula)