


training_load ~ bmi+injured
3.728 + (−0.025)*bmi + (0.347)*injured


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 6.2276 on 2 and 617 Degrees of Freedom
F p−value: 0.0021
R−Squared: 0.01979
Adjusted R−Squared: 0.0166099056760923
Root MSE: 0.99467578918398
AIC: 1760.86412185187



training_load against bmi
training_load
bmi
1
2
3
4
5
20
22
24
26
28



training_load against injured
training_load
injured
1
2
3
4
5
0.0
0.2
0.4
0.6
0.8
1.0



Residuals vs Fitted
Residuals
Fitted values
lm(formula)
2
1
0
−1
−2
−3
3.1
3.2
3.3
3.4
3.5
3.6
238
281
101


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
101
281
238
−3
−2
−1
0
1
2
3
−2
−1
0
1
2


Scale−Location
√|Standardized residuals|
0.0
0.5
1.0
1.5
238
281
101
3.1
3.2
3.3
3.4
3.5
3.6
Fitted values
lm(formula)


Cook's distance
101
238
474
Cook's distance
0.00
0.01
0.02
0.03
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
474
238
101
2
1
0
−1
−2
−3
0.000
0.005
0.010
0.015
0.020


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
101
474
238
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0.035
0
0.5
1
1.5
2
0
0.005
0.01
0.015
0.02
Leverage h_{ii}
lm(formula)