training_load ~ injured

3.137 + (0.35)*injured

A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC

F−statistic: 11.56486 on 1 and 618 Degrees of Freedom


F p−value: 0.00072


R−Squared: 0.01837


Adjusted R−Squared: 0.016781215755145


Root MSE: 0.995394809019574


AIC: 1759.76015502349



training_load against injured
training_load
injured
1
2
3
4
5
0.0
0.2
0.4
0.6
0.8
1.0



Residuals vs Fitted
Residuals
Fitted values
lm(formula)
2
1
0
−1
−2
−3
3.15
3.20
3.25
3.30
3.35
3.40
3.45
3.50


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−3
−2
−1
0
1
2
3
101
281


Scale−Location
√|Standardized residuals|
0.0
0.5
1.0
1.5
3.15
3.20
3.25
3.30
3.35
3.40
3.45
3.50
Fitted values
lm(formula)


Cook's distance
101
238
281
0.030
0.025
0.020
0.015
0.010
0.005
0.000
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
2
1
0
−1
−2
−3
Cook's distance
0.000
0.002
0.004
0.006
0.008
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
Cook's distance
Leverage h_{ii}
lm(formula)
0.000
0.005
0.010
0.015
0.020
0.025
0.001
0.002
0.003
0.004
0.005
0.006
0.007
0.008
0.009
0
0.5
1
1.5
2
2.5
3