


training_load ~ minutes_played+bmi
3.016 + (0)*minutes_played + (−0.024)*bmi


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 15.06715 on 2 and 617 Degrees of Freedom
F p−value: 0
R−Squared: 0.04657
Adjusted R−Squared: 0.0434752218426148
Root MSE: 0.980994889243247
AIC: 1743.69062500027



training_load against minutes_played
training_load
minutes_played
1
2
3
4
5
0
1000
2000
3000
4000



training_load against bmi
training_load
bmi
1
2
3
4
5
20
22
24
26
28



Residuals vs Fitted
Residuals
Fitted values
lm(formula)
2
1
0
−1
−2
2.4
2.6
2.8
3.0
3.2
3.4
3.6
47
281
107


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
107
281
474
−3
−2
−1
0
1
2
3
−2
−1
0
1
2


Scale−Location
47
281
107
√|Standardized residuals|
0.0
0.5
1.0
1.5
2.4
2.6
2.8
3.0
3.2
3.4
3.6
Fitted values
lm(formula)


Cook's distance
474
43
101
0.030
0.025
0.020
0.015
0.010
0.005
0.000
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
43
101
474
2
1
0
−1
−2
−3
0.000
0.005
0.010
0.015
0.020
0.025


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
2.5
2
474
43
101
1.5
1
0.5
0
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0
0.005
0.01
0.015
0.02
0.025
Leverage h_{ii}
lm(formula)