


training_load ~ prev_injuries+bmi+injured
3.711 + (0.03)*prev_injuries + (−0.025)*bmi + (0.327)*injured


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 4.3572 on 3 and 616 Degrees of Freedom
F p−value: 0.00476
R−Squared: 0.02078
Adjusted R−Squared: 0.0160102382284714
Root MSE: 0.994172386801826
AIC: 1762.23640277835



training_load against prev_injuries
training_load
prev_injuries
1
2
3
4
5
0
1
2
3
4



training_load against bmi
training_load
bmi
1
2
3
4
5
20
22
24
26
28



training_load against injured
training_load
injured
1
2
3
4
5
0.0
0.2
0.4
0.6
0.8
1.0



Residuals vs Fitted
Residuals
Fitted values
lm(formula)
2
1
0
−1
−2
−3
3.0
3.1
3.2
3.3
3.4
3.5
3.6
238
281
101


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
101
281
238
−3
−2
−1
0
1
2
3


Scale−Location
√|Standardized residuals|
Fitted values
lm(formula)
238
281
101
0.0
0.5
1.0
1.5
3.0
3.1
3.2
3.3
3.4
3.5
3.6


Cook's distance
101
238
474
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
2
1
0
−1
−2
−3
0.000
0.005
0.010
0.015
0.020
0.025
Leverage
lm(formula)
474
238
101
Cook's distance


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
2
1.5
1
0.5
0
101
238
474
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0
0.005
0.01
0.015
0.02
0.025
Leverage h_{ii}
lm(formula)