


training_load ~ prev_injuries+bmi
3.809 + (0.058)*prev_injuries + (−0.028)*bmi


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 1.75656 on 2 and 617 Degrees of Freedom
F p−value: 0.1735
R−Squared: 0.00566
Adjusted R−Squared: 0.00243849063365009
Root MSE: 1.00181717830747
AIC: 1769.73503742698



training_load against prev_injuries
training_load
prev_injuries
1
2
3
4
5
0
1
2
3
4



training_load against bmi
training_load
bmi
1
2
3
4
5
20
22
24
26
28



Residuals vs Fitted
Residuals
Fitted values
lm(formula)
2
1
0
−1
−2
3.1
3.2
3.3
3.4
474
466
584


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
584
474
466
−3
−2
−1
0
1
2
3
−2
−1
0
1
2


Scale−Location
√|Standardized residuals|
0.0
0.5
1.0
1.5
3.1
3.2
3.3
3.4
Fitted values
lm(formula)


Cook's distance
474
507
101
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
2
1
0
−1
−2
507
101
474
Cook's distance
0.000
0.005
0.010
0.015
0.020
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
2.5
474
101
507
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0
0.5
1
1.5
2
0
0.005
0.01
0.015
0.02
Leverage h_{ii}
lm(formula)