


training_load ~ prev_injuries+injured
3.114 + (0.03)*prev_injuries + (0.331)*injured


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 6.08121 on 2 and 617 Degrees of Freedom
F p−value: 0.00242
R−Squared: 0.01933
Adjusted R−Squared: 0.0161523081462397
Root MSE: 0.994907186799957
AIC: 1761.15255721603



training_load against prev_injuries
training_load
prev_injuries
1
2
3
4
5
0
1
2
3
4



training_load against injured
training_load
injured
1
2
3
4
5
0.0
0.2
0.4
0.6
0.8
1.0



Residuals vs Fitted
Residuals
2
1
0
−1
−2
3.1
3.2
3.3
3.4
3.5
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−3
−2
−1
0
1
2
3
101
238
281


Scale−Location
√|Standardized residuals|
0.0
0.5
1.0
1.5
3.1
3.2
3.3
3.4
3.5
Fitted values
lm(formula)


Cook's distance
101
238
281
0.025
0.020
0.015
0.010
0.005
0.000
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
2
1
0
−1
−2
−3
0.000
0.005
0.010
0.015
Leverage
lm(formula)
Cook's distance


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
Cook's distance
Leverage h_{ii}
lm(formula)
0.000
0.005
0.010
0.015
0.020
0.025
0
0.005
0.01
0.015
2.5
2
1.5
1
0.5
0