


training_load ~ prev_injuries
3.149 + (0.058)*prev_injuries


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 2.41267 on 1 and 618 Degrees of Freedom
F p−value: 0.12087
R−Squared: 0.00389
Adjusted R−Squared: 0.00227697640684832
Root MSE: 1.0027098590282
AIC: 1768.83946170888



training_load against prev_injuries
training_load
prev_injuries
1
2
3
4
5
0
1
2
3
4



Residuals vs Fitted
Residuals
2
1
0
−1
−2
3.15
3.20
3.25
3.30
3.35
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
105
466
584
−3
−2
−1
0
1
2
3
−2
−1
0
1
2


Scale−Location
√|Standardized residuals|
0.0
0.5
1.0
1.5
3.15
3.20
3.25
3.30
3.35
Fitted values
lm(formula)


Cook's distance
115
384
96
0.020
0.015
0.010
0.005
0.000
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
2
1
0
−1
−2
96
Cook's distance
0.000
0.005
0.010
0.015
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
2.5
2
1.5
1
0.5
0
96
0.020
0.015
0.010
0.005
0.000
Cook's distance
0
0.002
0.004
0.006
0.008
0.01
0.012
0.014
Leverage h_{ii}
lm(formula)