

MATH 231-01: Homework Assignment 3

22 September 2025

Due: 29 September 2025 by 10:00pm Eastern time, submitted on Moodle as a single PDF.

Instructions: Write your solutions on the following pages. If you need more space, you may add pages, but make sure they are in order and label the problem number(s) clearly. You should attempt each problem on scrap paper first, before writing your solution here. Excessively messy or illegible work will not be graded. You must show your work/reasoning to receive credit. You do not need to include every minute detail; however the process by which you reached your answer should be evident. You may work with other students, but please write your solutions in your own words.

Name:

Score:

1. The angle between two curves at a point of intersection is defined to be the smallest angle between their tangent lines at that point. Let C_1 and C_2 be the curves parametrized by $\mathbf{r}_1(t) = \langle \cos(t), \sin(2t), t \rangle$ and $\mathbf{r}_2(t) = \langle \sin(2t), \cos(t), t \rangle$ for $t \in [0, \pi]$.
 - (a) Find all points of intersection between C_1 and C_2 . (The double-angle formula $\sin(2t) = 2 \sin(t) \cos(t)$ may be helpful.)
 - (b) Find the angle between C_1 and C_2 at each point of intersection. If you can't write down the exact value, round your answer to the nearest hundredth of a radian.

2. Prove the derivative rule for the dot product. In other words, given arbitrary vector functions $\mathbf{r}_1(t) = \langle x_1(t), y_1(t), z_1(t) \rangle$ and $\mathbf{r}_2(t) = \langle x_2(t), y_2(t), z_2(t) \rangle$, show that

$$(\mathbf{r}_1(t) \cdot \mathbf{r}_2(t))' = \mathbf{r}_1'(t) \cdot \mathbf{r}_2(t) + \mathbf{r}_1(t) \cdot \mathbf{r}_2'(t).$$

3. Let $\mathbf{r}_1(t) = \langle e^{2t}, t^2, \sin(3t) \rangle$ and $\mathbf{r}_2(t) = \langle \ln(1 + t^2), \cos(2t), t^3 \rangle$. Let C be the curve parametrized by $\mathbf{r}(t) = \mathbf{r}_1(t) \times \mathbf{r}_2(t)$. Find an equation for the line tangent to C at the point $\mathbf{r}(0)$.

4. The acceleration, initial velocity, and initial position of an object traveling through space are given by

$\mathbf{r}''(t) = \langle -6, 2, 4 \rangle$, $\mathbf{r}'(0) = \langle -5, 1, 3 \rangle$, and $\mathbf{r}(0) = \langle 6, -2, 1 \rangle$,
respectively. Find the object's position function $\mathbf{r}(t)$.

5. Find the arc length of the curve parametrized by $\mathbf{r}(t) = \langle t - \sin(t), \cos(t) \rangle$ for $t \in [0, 2\pi]$. (The half-angle formula $\sqrt{2}\sin(t/2) = \sqrt{1 - \cos(t)}$ for $t \in [0, 2\pi]$ may be helpful.)

6. Let C be the *moment curve*, parametrized by $\mathbf{r}(t) = \langle t, t^2, t^3 \rangle$.
- (a) Find the curvature of C .

- (b) Does the curvature increase or decrease as $t \rightarrow \infty$? (No need to be rigorous here; an educated guess is fine!)

7. Complete Problem 10 on p. 889 of the textbook (Section 11.4)¹

¹The names in this problem are purely coincidental and do not refer to individuals in our class!