Supplementary Online Content

Fu DJ, Keenan TD, Faes L, et al. Insights from survival analyses during 12 years of anti–vascular endothelial growth factor therapy for neovascular age-related macular degeneration. *JAMA Ophthalmol*. Published online November 19, 2020. doi:10.1001/jamaophthalmol.2020.5044

eFigure 1. Overview of MEH AMD Anti-VEGF Treatment Guidelines

eFigure 2. Overview of Treatment-Naïve Neovascular AMD Patients

eFigure 3. Cohort Selection

eFigure 4. Number of Cumulative Events

eTable. Demographics and Clinical Features of Cohort

This supplementary material has been provided by the authors to give readers additional information about their work.

© 2020 American Medical Association. All rights reserved.

eFigure 1. Overview of MEH AMD Anti-VEGF Treatment Guidelines



© 2020 American Medical Association. All rights reserved.

eFigure 2. Overview of Treatment-Naïve Neovascular AMD Patients



© 2020 American Medical Association. All rights reserved.

eFigure 3. Cohort Selection



	Potential observation time		Seen in clinic	
Year	**n**	**(%)**	**n**	**(%)**
1	7122	(91%)	5970	(84%)
2	6326	(81%)	4131	(65%)
3	5585	(72%)	2738	(49%)
4	4851	(62%)	1711	(35%)
5	4218	(54%)	973	(23%)
6	3568	(46%)	558	(16%)
7	2768	(35%)	305	(11%)
8	2084	(27%)	212	(10%)
9	1432	(18%)	134	(9%)
10	751	(10%)	63	(8%)

© 2020 American Medical Association. All rights reserved.

eFigure 4. Number of Cumulative Events



© 2020 American Medical Association. All rights reserved.

eTable. Demographics and Clinical Features of Cohort

	Overall	Aflibercept	Ranibizumab	Mean difference (95% CI)	P-value
Gender	n = 7802	n = 3951	n = 3851		
Female (%)	4776 (61.2)	2371 (60.0)	2405 (62.5)		.029
Male (%)	3026 (38.8)	1580 (40.0)	1446 (37.5)		
Age at baseline (Years)					
Mean (SD)	78.68 (8.75)	79.15 (8.70)	78.21 (8.77)	0.95 (0.55 - 1.33)	< .001
Ethnicity					
Afro Caribbean (%)	177 (2.3)	80 (2.0)	97 (2.5)		< .001
Caucasian (%)	4785 (61.3)	2081 (52.7)	2704 (70.2)		
Chinese (%)	41 (0.5)	17 (0.4)	24 (0.6)		
Mixed (%)	21 (0.3)	9 (0.2)	12 (0.3)		
South-East Asian (%)	652 (8.4)	350 (8.9)	302 (7.8)		
Unknown (%)	2126 (27.2)	1414 (35.8)	712 (18.5)		
VA at baseline in ETDRS letter score (approximate Snellen equivalent					
Mean	54.69 (20/80)	55.09 (20/80)	54.28 (20/80)	0.81 (0.10 - 1.52)	.025
SD	16.02 (3 lines)	15.76 (3 lines)	16.28 (3 lines)		
Treatment initiation					
Post-2013 (%)	4541 (58.2)	3951 (100.0)	590 (15.3)		< .001
Pre-2013 (%)	3261 (41.8)	0 (0.0)	3261 (84.7)		
Injection number					
Mean (SD)	10.25 (8.65)	12.11 (9.44)	8.34 (7.28)	3.77 (3.39 - 4.13)	< .001
Induction injection interval					
Mean (SD)	33.19 (34.14)	30.77 (6.16)	36.64 (52.94)	5.87 (3.74 - 8.0)	< .001
Induction phase					
Completed appropriately (%)	6212 (79.6)	3563 (90.2)	2649 (68.8)		< .001
Not completed (%)	1590 (20.4)	388 (9.8)	1202 (31.2)		

© 2020 American Medical Association. All rights reserved.