

Likelihood

Materials and Methods

Observed data (Y)

For each patient $i = 1, \dots, N$ admitted to one of the wards in the study period, we denote

- w_i the ward where the patient is admitted (1 for adult ICU, 2 for paediatric ICU)
- k_i the number of times the patient is admitted (1 if no readmission)
- A_i and D_i vectors containing the times of admission and discharge from the ward
- P_i and N_i vectors containing the times of positive and negative swabs (positive defined as any of the samples taken is positive ; negative defined as all samples taken are negative).
- p_i and n_i the size of those vectors, ie the number of positive and negative swabs.
- $S_i = \{s_i^1, \dots, s_i^{m_i}\}$ a set of m_i genetic sequences of MRSA isolated in patient i at times $T_i = \{t_i^1, \dots, t_i^{m_i}\}$; collection dates T_i are ordered so that $t_i^k \leq t_i^{k+1}$.
- $d_{s_i^k, s_j^q}$ the number of transitions between sequence k of patient i and sequence q of patient j .
- $g_{s_i^k, s_j^q}$ the number of transversions between sequence k of patient i and sequence q of patient j .
- $l_{s_i^k, s_j^q}$ the number of typed nucleotides common to sequences s_i^k and s_j^q .

Augmented (unobserved) data (Z)

For each patient i admitted to one of the wards in the study period, we denote

- C_i the colonisation time (we assume no supercolonisations)
- E_i the time of end of colonisation.

For each pair of sequences s_i^k and s_j^q in patients i and j , we denote

- $b_{s_i^k, s_j^q}$ an indicator function with value 1 if a direct ancestor to sequence s_i^k has been sampled in patients i or j , and 0 otherwise.

We denote $I_w(t) = \sum_{i=1}^N \mathbf{1}_{\{w_i=w\}} \mathbf{1}_{\{C_i \leq t < E_i\}} \sum_{l=1}^{k_i} \mathbf{1}_{\{A_i[l] \leq t < D_i[l]\}}$ the number of patients in ward w who are colonized at time t .

Parameters (θ)

Parameters of the model are:

- β a 2 by 2 matrix containing $\beta_{i \leftarrow j}$, the person to person transmission rate from ward j to ward i
- $\beta_{\text{ward} \leftarrow \text{out}}$ the force of infection from outside the 2 wards applied to patients in the wards
- $\beta_{\text{out} \leftarrow \text{out}}$ the force of infection from outside the 2 wards applied to patients when they are not in the wards (eg inbetween two admissions)

- Sp the specificity of the testing, ie the probability of getting a negative test given uncolonized (assumed 100%)
- Se the sensitivity of the testing, ie the probability of getting a positive test given colonized
- π the probability of being already colonized at first admission
- μ and σ the mean and standard deviation of the duration of colonization.
- ν_1 and ν_2 the rate of transitions ($A \leftrightarrow G$ and $C \leftrightarrow T$) and transversions (other changes) of the DNA sequences. In practice, we will use $\nu_2 = \kappa \nu_1$ with $\kappa \in \mathbb{R}_+$.

Statistical Model

In the following, $\mathbf{1}_{\{\cdot\}}$ denotes the indicator function, defined by $\mathbf{1}_{\{X\}} = 1$ if X is true, and 0 otherwise. The joint density of the observed data, the augmented data, and the model parameters is:

$$P(\mathbf{Y}, \mathbf{Z}, \boldsymbol{\theta}) = P(\mathbf{Y}|\mathbf{Z}) P(\mathbf{Z}|\boldsymbol{\theta}) P(\boldsymbol{\theta})$$

where $P(\mathbf{Y}|\mathbf{Z})$, $P(\mathbf{Z}|\boldsymbol{\theta})$ and $P(\boldsymbol{\theta})$ refer to the observation level, the transmission level and the prior level respectively.

Observation level

The observation level ensures that the observed data are consistent with the augmented data:

$$P(\mathbf{Y}|\mathbf{Z}) = \prod_{i=1}^N \mathbf{1}_{\{C_i < E_i\}} \prod_{j=1}^{p_i} ((\mathbf{1}_{\{P_i[j] < C_i\}} + \mathbf{1}_{\{P_i[j] \geq E_i\}}) \times (1 - Sp) + \mathbf{1}_{\{C_i \leq P_i[j] < E_i\}} \times Se) \prod_{k=1}^{n_i} ((\mathbf{1}_{\{N_i[k] < C_i\}} + \mathbf{1}_{\{N_i[k] \geq E_i\}}) \times Sp + \mathbf{1}_{\{C_i \leq N_i[k] < E_i\}} \times (1 - Se))$$

The first line describes the positive tests, which can be either false positives (first term) or true positives (second term). The second line describes the negative tests, which can be either true negatives (first term) or false negatives (second term).

Transmission level (discrete time version ; time step = half day or day ?)

In the discrete version $A_i[k]$ is the first time step where individual i is in hospital (for his/her k_{th} stay), and $D_i[k]$ is the first time step where he/she is out of hospital (after his/her k_{th} stay). Individual i can transmit staph aureus from time step C_i to time step $E_i - 1$.

$$P(\mathbf{Z}|\boldsymbol{\theta}) = \prod_{i=1}^N \left(\Omega_i^{(1)} + \Omega_i^{(2)} \right) \times (\Phi_{\mu, \sigma}(E_i - C_i + 0.5) - \Phi_{\mu, \sigma}(E_i - C_i - 0.5))$$

where $\Phi_{\mu, \sigma}$ is the cumulative density function of a Gamma distribution with mean μ and standard deviation σ (we assume that the duration of colonisation is Gamma distributed), and:

$$\begin{aligned}
\Omega_i^{(1)} &= \pi \times \mathbf{1}_{\{C_i < A_i[1]\}} \\
\Omega_i^{(2)} &= (1 - \pi) \times \mathbf{1}_{\{C_i \geq A_i[1]\}} \times e^{-\sum_{t=A_i[1]}^{C_i-1} \lambda_i(t)} \left(1 - e^{-\lambda_i(C_i)}\right) \eta_i(C_i)
\end{aligned}$$

$\Omega_i^{(1)}$ is the probability that individual i is colonized before his/her first admission in the wards ; $\Omega_i^{(2)}$ is the probability that individual i is colonized after his/her first admission in the wards. $\lambda_i(t)$ is the force of transmission applied to individual i at time t . It is equal to:

$$\begin{aligned}
\lambda_i^{(t)} &= \sum_{w=1}^2 \beta_{w_i \leftarrow w} I_w(t) + \beta_{\text{ward} \leftarrow \text{out}} \text{ if individual } i \text{ is in a ward at time } t \\
&= \beta_{\text{out} \leftarrow \text{out}} \text{ otherwise}
\end{aligned}$$

$\eta_i(C_i)$ is the probability of observing sequences S_i given the possible contamination sources at time C_i .

It is equal to:

$$\begin{aligned}
\eta_i(C_i) &= \frac{\sum_{j \text{ colonized and in a ward at time } C_i} \beta_{w_i \leftarrow w_j} f_{i \leftarrow j} + \beta_{\text{ward} \leftarrow \text{out}} f_{\text{i, ward} \leftarrow \text{out}}}{\sum_{j \text{ colonized and in a ward at time } C_i} \beta_{w_i \leftarrow w_j} + \beta_{\text{ward} \leftarrow \text{out}}} \text{ if individual } i \text{ is in a ward at time } C_i \\
&= f_{\text{i, out} \leftarrow \text{out}} \text{ otherwise}
\end{aligned}$$

$f_{i,j}$ is the probability of observing sequences s_i and s_j at respective time steps t_i and t_j given that individual j infected individual i (see next section on the genetic likelihood). Similarly, $f_{\text{i, ward} \leftarrow \text{out}}$ and $f_{\text{i, out} \leftarrow \text{out}}$ are the probability of observing sequence s_i at time t_i given that individual i was infected from outside the wards while he was in or outside hospital respectively.

Genetic likelihood

The genetic likelihood given the infection $i \leftarrow j$ relies on how likely it is that a sequence in S_i has its closest direct ancestor in S_i or S_j . The probability of observing the sequence s_i^k given that patient j infected patient i is:

$$p(s_i^k | s_i^1, \dots, s_i^{k-1}, S_j, i \leftarrow j) = b_{s_i^k, j} (\Xi_{s_i^k}^{(j)} + \Xi_{s_i^k}^{(i)}) + (1 - b_{s_i^k, j})$$

with:

- $\Xi_{s_i^k}^{(j)}$: direct ancestries to s_i^k from sequences in patient j (S_j)
- $\Xi_{s_i^k}^{(i)}$: direct ancestries to s_i^k from sequences in patient i (S_i)

These are given by:

$$\begin{aligned}\Xi_{s_i^k}^{(j)} &= \sum_{q=1}^{m_j} \left(\mathcal{P} \left(d_{s_i^k, s_j^q} | \nu_1 \mathbf{1}_{\{t_i^k \geq t_j^q\}} (t_i^k - t_j^q) l_{s_i^k, s_j^q} \right) + \mathcal{P} \left(g_{s_i^k, s_j^q} | \nu_2 \mathbf{1}_{\{t_i^k \geq t_j^q\}} (t_i^k - t_j^q) l_{s_i^k, s_j^q} \right) \right) \\ \Xi_{s_i^k}^{(i)} &= \sum_{r=1}^{k-1} \left(\mathcal{P} \left(d_{s_i^k, s_i^r} | \nu_1 (t_i^k - t_i^r) l_{s_i^k, s_i^r} \right) + \mathcal{P} \left(g_{s_i^k, s_i^r} | \nu_2 (t_i^k - t_i^r) l_{s_i^k, s_i^r} \right) \right)\end{aligned}$$

where $\mathcal{P}(\cdot|\lambda)$ is the probability mass function of a Poisson distribution with parameter λ . Left-hand terms correspond to transitions, while right-hand terms correspond to transversions.

The probability of observing a set of sequences S_i given that patient j (with sequences S_j) infected patient i is given by:

$$p(S_i | S_j, i \leftarrow j) = \prod_{k=1}^{m_i} p(s_i^k | s_i^1, \dots, s_i^{k-1}, S_j, i \leftarrow j)$$

Note that in the absence of genetic information ($m_i < 1$ and $m_j = 0$), $b_{s_i^k, j}$ is automatically set to 0.

Prior level

For all model parameters, independent prior distributions were chosen:

- uniform on $[0, 1]$ for Sp , Se , and π ,
- flat exponential (mean 1000) for all other parameters.

Parameter Estimation

A Markov chain Monte Carlo (MCMC) method was used to sample the joint posterior distribution $P(\mathbf{Y}, \mathbf{Z}, \boldsymbol{\theta})$. Sp , Se and π were updated using the Gibbs sampler, and all other parameters using a Metropolis algorithm.

Appendix: Full formulation of the transmission level

In the discrete version $A_i[k]$ is the first time step where individual i is in hospital (for his/her k_{th} stay), and $D_i[k]$ is the first time step where he/she is out of hospital (after his/her k_{th} stay). Individual i can transmit staph aureus from time step C_i to time step $E_i - 1$.

$$P(\mathbf{Z} | \boldsymbol{\theta}) = \prod_{i=1}^N \left(\Omega_i^{(1)} + \Omega_i^{(2)} + \Omega_i^{(3)} \right) \times (\Phi_{\mu, \sigma}(E_i - C_i + 0.5) - \Phi_{\mu, \sigma}(E_i - C_i - 0.5))$$

where $\Phi_{\mu, \sigma}$ is the cumulative density function of a Gamma distribution with mean μ and standard deviation σ (we assume that the duration of colonisation is Gamma distributed), and:

$$\begin{aligned}
\Omega_i^{(1)} &= \pi \times \mathbf{1}_{\{C_i < A_i[1]\}} \\
\Omega_i^{(2)} &= (1 - \pi) \sum_{l=1}^{k_i} \mathbf{1}_{\{A_i[l] \leq C_i < D_i[l]\}} \\
&\quad \times \exp \left(-\mathbf{1}_{\{l \geq 2\}} \sum_{s=1}^{l-1} \left[\sum_{t=A_i[s]}^{D_i[s]-1} \left(\sum_{w=1}^2 \beta_{w_i \leftarrow w} I_w(t) \right) + \beta_{\text{ward} \leftarrow \text{out}} (D_i[s] - A_i[s]) \right] \right) \\
&\quad \times \exp \left(-\mathbf{1}_{\{l \geq 2\}} \sum_{s=1}^{l-1} [\beta_{\text{out} \leftarrow \text{out}} (A_i[s+1] - D_i[s])] \right) \\
&\quad \times \exp \left(-\sum_{t=A_i[l]}^{C_i-1} \left(\sum_{w=1}^2 \beta_{w_i \leftarrow w} I_w(t) \right) - \beta_{\text{ward} \leftarrow \text{out}} (C_i - A_i[l]) \right) \\
&\quad \times \left(1 - \exp \left(-\sum_{w=1}^2 \beta_{w_i \leftarrow w} I_w(C_i) - \beta_{\text{ward} \leftarrow \text{out}} \right) \right) \\
&\quad \times \frac{\sum_{j=1}^N \mathbf{1}_{\{C_j \leq C_i < E_j\}} \sum_{r=1}^{k_j} \mathbf{1}_{\{A_j[r] \leq C_i < D_j[r]\}} \beta_{w_i \leftarrow w_j} f_{i \leftarrow j} + \beta_{\text{ward} \leftarrow \text{out}} f_{i, \text{ward} \leftarrow \text{out}}}{\sum_{j=1}^N \mathbf{1}_{\{C_j \leq C_i < E_j\}} \sum_{r=1}^{k_j} \mathbf{1}_{\{A_j[r] \leq C_i < D_j[r]\}} \beta_{w_i \leftarrow w_j} + \beta_{\text{ward} \leftarrow \text{out}}} \\
\Omega_i^{(3)} &= \mathbf{1}_{\{k_i > 1\}} (1 - \pi) \sum_{l=1}^{k_i-1} \mathbf{1}_{\{D_i[l] \leq C_i < A_i[l+1]\}} \\
&\quad \times \exp \left(-\sum_{s=1}^l \left[\sum_{t=A_i[s]}^{D_i[s]-1} \left(\sum_{w=1}^2 \beta_{w_i \leftarrow w} I_w(t) \right) + \beta_{\text{ward} \leftarrow \text{out}} (D_i[s] - A_i[s]) \right] \right) \\
&\quad \times \exp \left(-\mathbf{1}_{\{l \geq 2\}} \sum_{s=1}^{l-1} [\beta_{\text{out} \leftarrow \text{out}} (A_i[s+1] - D_i[s])] \right) \\
&\quad \times \exp (-\beta_{\text{out} \leftarrow \text{out}} (C_i - D_i[l])) \\
&\quad \times (1 - \exp (-\beta_{\text{out} \leftarrow \text{out}})) \\
&\quad \times f_{i, \text{out} \leftarrow \text{out}}
\end{aligned}$$

$\Omega_i^{(1)}$ is the probability that individual i is colonized before his/her first admission in the wards ; $\Omega_i^{(2)}$ is the probability that individual i is colonized during one of his/her stays in the wards ; $\Omega_i^{(3)}$ is the probability, that individual i , if admitted several times, is colonized between successive stays in the wards.

$f_{i,j}$ is the probability of observing sequences s_i and s_j at respective time steps t_i and t_j given that individual j infected individual i (see next section on the genetic likelihood). Similarly, $f_{i, \text{ward} \leftarrow \text{out}}$ and $f_{i, \text{out} \leftarrow \text{out}}$ are the probability of observing sequence s_i at time t_i given that individual i was infected from outside the wards while he was in or outside hospital respectively.