

Radiative transitions of strange heavy axial mesons in light-cone QCD sum rules

T. M. Aliev ^{1,*} S. Bilmis ^{1,2,†} and M. Savci ^{1,‡}

¹*Department of Physics, Middle East Technical University, Ankara, 06800, Turkey*

²*TUBITAK ULAKBIM, Ankara, 06510, Turkey*

(Dated: August 13, 2026)

Abstract

We calculate the radiative decay widths of $D_{s1}(2460) \rightarrow D_s\gamma$, $D_{s1}(2536) \rightarrow D_s\gamma$, $B_{s1}(5750) \rightarrow B_s\gamma$, and $B_{s1}(5830) \rightarrow B_s\gamma$ within light-cone QCD sum rules, taking into account the mixing between the 1P_1 and 3P_1 states.

Using the lattice-normalized leading-twist photon input and the preferred sector-specific mixing angles, we obtain $\Gamma[D_{s1}(2460) \rightarrow D_s\gamma] = 34.96^{+8.57}_{-6.90}$ keV, $\Gamma[D_{s1}(2536) \rightarrow D_s\gamma] = 0.99^{+2.57}_{-0.90}$ keV, $\Gamma[B_{s1}(5750) \rightarrow B_s\gamma] = 11.34^{+2.72}_{-2.20}$ keV, and $\Gamma[B_{s1}(5830) \rightarrow B_s\gamma] = 8.70^{+2.53}_{-2.23}$ keV.

The charm-strange results exhibit a pronounced hierarchy: the $D_{s1}(2536) \rightarrow D_s\gamma$ width is strongly suppressed relative to that of $D_{s1}(2460) \rightarrow D_s\gamma$. This pattern follows from constructive interference between the two basis-current contributions for the lower state and destructive interference for the higher state. In the bottom-strange sector, the two predicted widths are comparable within their uncertainties.

* taliev@metu.edu.tr

† sbilmis@metu.edu.tr

‡ savci@metu.edu.tr

I. INTRODUCTION

Radiative transitions of heavy-light mesons provide a sensitive probe of their internal structure. For positive-parity strange heavy mesons, such transitions are especially useful because they depend on the spin composition of the physical axial-vector states and on the coupling of the photon to both the heavy and strange quark degrees of freedom. They therefore complement mass spectroscopy and strong-decay measurements in testing the assignment of the observed states and the role of spin mixing.

The charm-strange axial-vector mesons $D_{s1}(2460)$ and $D_{s1}(2536)$ are particularly important examples. The $D_{s1}(2460)$ is a very narrow state below the D^*K threshold and its radiative decay to $D_s\gamma$ has been observed in B decays and in continuum production [1, 2]. The $D_{s1}(2536)$, on the other hand, is well established in hadronic channels such as D^*K and $D\pi K$ [3, 4], but its radiative decay to $D_s\gamma$ has not been observed with comparable significance. The measured width of the $D_{s1}(2536)$ is below the MeV scale, which makes a suppressed radiative branching fraction experimentally difficult to isolate [4]. In the bottom-strange sector the $B_{s1}(5830)$ state has been observed in B^*K spectroscopy [5, 6], whereas no direct information is available on its radiative transition to $B_s\gamma$.

Theoretical studies of these radiative modes have been carried out using constituent quark models, relativistic quark models, effective approaches, and QCD sum rules [7–19]. The light-cone QCD sum-rule calculation of Ref. [10] showed that $D_{s1}(2460) \rightarrow D_s\gamma$ is expected to be one of the leading radiative channels of the $D_{s1}(2460)$. More recently, Bondar and Milstein argued that the smallness of $D_{s1}(2536) \rightarrow D_s\gamma$ follows from a cancellation between the two spin components of the physical axial-vector state [18, 19]. Their analysis also considered the corresponding bottom-strange transitions, including both the established $B_{s1}(5830)$ state and a lower $B_{s1}(5750)$ configuration. This raises the question of whether the same cancellation mechanism emerges within light-cone QCD sum rules. Addressing this question is the principal objective of the present work.

In this work, we calculate the radiative transitions $D_{s1}(2460) \rightarrow D_s\gamma$, $D_{s1}(2536) \rightarrow D_s\gamma$, $B_{s1}(5750) \rightarrow B_s\gamma$, and $B_{s1}(5830) \rightarrow B_s\gamma$ within light-cone QCD sum rules. We retain the strange-quark-mass terms and use photon distribution amplitudes, including the recent lattice-normalized input for the leading-twist photon distribution amplitude. The two basis-current sum rules are combined into the physical axial-vector currents, which allows us to test whether the cancellation mechanism proposed for $D_{s1}(2536) \rightarrow D_s\gamma$ is reproduced in the sum-rule framework. For the bottom-strange sector, the lower $B_{s1}(5750)$ state is treated as a theoretical diagnostic, while the $B_{s1}(5830)$ result is a prediction for the experimentally established state.

The paper is organized as follows. In Sec. II, we define the axial-vector interpolating currents

and derive the sum rules for the transition form factors. Section III presents the numerical analysis, comparison with previous studies, and discussion of the experimental status. We summarize our results in Sec. IV.

II. THEORETICAL FRAMEWORK

We consider the radiative transition

$$A_i(p', \eta) \rightarrow P(p) \gamma(q, \varepsilon),$$

where A_i denotes a physical heavy axial-vector meson of mass M_i , P is the corresponding pseudoscalar meson, and γ is a real photon. The light quark field appearing below is understood to be the strange-quark field. This process is described by the vacuum correlation function

$$\Pi_{\mu\nu}^{(i)}(p', q) = i^2 \int d^4x \int d^4y e^{ip \cdot x + iq \cdot y} \langle 0 | T \left\{ J_\mu^{(i)}(x) J_\nu^{\text{em}}(y) J_5^\dagger(0) \right\} | 0 \rangle, \quad (1)$$

where

$$J_\nu^{\text{em}} = e_q \bar{q} \gamma_\nu q + e_Q \bar{Q} \gamma_\nu Q \quad (2)$$

is the electromagnetic current, with e_q and e_Q denoting the electric charges of the light and heavy quarks, respectively. The pseudoscalar meson interpolating current is

$$J_5 = \bar{q} i \gamma_5 Q. \quad (3)$$

For the axial-vector channel, we choose two independent interpolating currents,

$$\begin{aligned} J_\mu^A &= \bar{q} \gamma_\mu \gamma_5 Q, \\ J_\mu^B &= \frac{i}{m_Q + m_q} \bar{q} \sigma_{\mu\nu} p'^\nu \gamma_5 Q. \end{aligned} \quad (4)$$

The interpolating currents for the two physical states are related to the basis currents by the rotation

$$\begin{pmatrix} J_\mu^{(1)} \\ J_\mu^{(2)} \end{pmatrix} = \begin{pmatrix} \sin \theta & \cos \theta \\ \cos \theta & -\sin \theta \end{pmatrix} \begin{pmatrix} J_\mu^A \\ J_\mu^B \end{pmatrix}, \quad (5)$$

where θ denotes the 3P_1 - 1P_1 mixing angle.

Following the standard QCD sum-rule procedure, the correlation function is evaluated in two

complementary representations: in terms of hadronic states on the phenomenological side and in terms of quark and gluon degrees of freedom on the QCD side.

The hadronic representation is obtained by inserting complete sets of states with the same quantum numbers as the interpolating currents and isolating the ground-state contributions. This gives

$$\Pi_{\mu\nu}^{(i)} = \frac{\langle 0 | J_\mu^{(i)} | A_i(p', \eta) \rangle \langle A_i(p', \eta) | J_\nu^{\text{em}} | P(p) \rangle \langle P(p) | J_5^\dagger | 0 \rangle}{(M_i^2 - p'^2)(m_P^2 - p^2)} + \dots, \quad (6)$$

where the ellipsis denotes contributions from higher resonances and the continuum.

The relevant matrix elements are defined as

$$\langle 0 | J_\mu^{(i)} | A_i(p', \eta) \rangle = M_i f_i \eta_\mu, \quad (7)$$

and

$$\langle P(p) | J_5^\dagger | 0 \rangle = \frac{m_P^2 f_P}{m_Q + m_q}. \quad (8)$$

Gauge invariance constrains the electromagnetic-current matrix element to the form

$$\langle A_i(p', \eta) | J_\nu^{\text{em}} | P(p) \rangle = e g_i [(p \cdot q) \eta_\nu - (\eta \cdot q) p_\nu], \quad (9)$$

where

$$q = p' - p,$$

and $g_i(q^2)$ denotes the electric-dipole transition form factor. We determine this form factor at the real-photon point, $q^2 = 0$.

Substituting the above matrix elements into the hadronic representation, summing over the polarization of the axial-vector meson according to

$$\sum_\lambda \eta_\mu^{(\lambda)} \eta_\nu^{(\lambda)*} = -g_{\mu\nu} + \frac{p'_\mu p'_\nu}{M_i^2},$$

one obtains

$$\Pi_{\mu\nu}^{(i)} = \frac{M_i f_i g_i m_P^2 f_P}{(m_Q + m_q)(M_i^2 - p'^2)(m_P^2 - p^2)} \left[(p \cdot q) \left(-g_{\mu\nu} + \frac{p'_\mu p'_\nu}{M_i^2} \right) - p_\nu \left(-q_\mu + \frac{p'_\mu (p' \cdot q)}{M_i^2} \right) \right]. \quad (10)$$

Multiplying both sides by the photon polarization vector ε^ν , the correlation function can be written as

$$\Pi_{\mu\nu}^{(i)} \varepsilon^\nu = \frac{M_i f_i g_i m_P^2 f_P}{(m_Q + m_q)(M_i^2 - p'^2)(m_P^2 - p^2)} \{ -\varepsilon_\mu (p \cdot q) + (p \cdot \varepsilon) q_\mu + \dots \}, \quad (11)$$

where the omitted terms correspond to other independent Lorentz structures.

We use the coefficient of the Lorentz structure

$$(p \cdot \varepsilon) q_\mu - \varepsilon_\mu (p \cdot q)$$

to derive the corresponding light-cone QCD sum rules for the transition form factors $g_i(q^2 = 0)$.

On the QCD side, the correlation function is evaluated in the deep Euclidean region using the background-field light-cone expansion. The invariant amplitude multiplying the selected Lorentz structure is decomposed as

$$\Pi^{\text{OPE}} = \Pi^{\text{pert}} + \Pi^{2\text{P}} + \Pi^{3\text{P}}, \quad (12)$$

where the last two terms contain the two- and three-particle photon DAs up to twist four. The explicit spectral densities and auxiliary convolution operators are collected in Appendix A.

We denote the physical axial-vector mass by M_i and reserve m_Q and m_s for the heavy- and strange-quark masses. The double Borel parameters in the axial-vector and pseudoscalar channels are denoted by $\mathcal{M}_A^2$ and $\mathcal{M}_P^2$, respectively. We define

$$d \equiv m_Q + m_s, \quad \mathcal{M}^2 = \frac{\mathcal{M}_A^2 \mathcal{M}_P^2}{\mathcal{M}_A^2 + \mathcal{M}_P^2}, \quad u_0 = \frac{\mathcal{M}_P^2}{\mathcal{M}_A^2 + \mathcal{M}_P^2}, \quad a_0 = 1 - u_0, \quad (13)$$

A. Background-field reduction

For a real photon with momentum q and polarization vector ε_μ , the external electromagnetic field-strength tensor is

$$F_{\mu\nu}(x) = i(\varepsilon_\nu q_\mu - \varepsilon_\mu q_\nu) e^{iq \cdot x}. \quad (14)$$

Within this formalism, the correlation function in Eq. (1) can be rewritten as

$$\Pi_\mu^{(i)}(p, q) = i \int d^4x e^{ip \cdot x} \left\langle 0 \left| T \left\{ J_\mu^{(i)}(x) J_5^\dagger(0) \right\} \right| 0 \right\rangle_F, \quad (15)$$

where the subscript F indicates that the vacuum expectation value is evaluated in the presence of the external electromagnetic field.

The correlation function receives two distinct contributions. The first is generated by perturbative photon emission from a quark propagator. The second describes the long-distance interaction of the photon with the quark fields and is parameterized in terms of photon light-cone distribution amplitudes (DAs).

Applying Wick's theorem, the correlation function is expressed in terms of heavy- and light-quark propagators,

$$\Pi_\mu^{(i)} = iN_c \int d^4x e^{ip \cdot x} \text{Tr} [\Gamma_\mu^{(i)} S_Q(x) \gamma_5 S_q(-x)]_F, \quad (16)$$

where $\Gamma_\mu^{(i)}$ denotes the Dirac structure associated with the chosen interpolating current, and N_c is the number of colors. For the two basis currents,

$$\Gamma_\mu^A = \gamma_\mu \gamma_5, \quad \Gamma_\mu^B = \frac{i}{d} \sigma_{\mu\lambda} p'^\lambda \gamma_5. \quad (17)$$

B. Background-field propagators

Equation (16) requires the light- and heavy-quark propagators in external gluonic and electromagnetic background fields. The corresponding propagator is

$$\begin{aligned} S_q(x) = & \int \frac{d^4k}{(2\pi)^4} e^{-ik \cdot x} \frac{i(\not{k} + m_q)}{k^2 - m_q^2} \\ & - g_s \int \frac{d^4k}{(2\pi)^4} e^{-ik \cdot x} \int_0^1 du \left[\frac{\not{k} + m_q}{2(k^2 - m_q^2)^2} G^{\mu\nu}(ux) \sigma_{\mu\nu} - \frac{ux_\mu G^{\mu\nu}(ux) \gamma_\nu}{k^2 - m_q^2} \right] \\ & - ie_q \int \frac{d^4k}{(2\pi)^4} e^{-ik \cdot x} \int_0^1 du \left[\frac{\not{k} + m_q}{2(k^2 - m_q^2)^2} F^{\mu\nu}(ux) \sigma_{\mu\nu} + \frac{ux_\mu F^{\mu\nu}(ux) \gamma_\nu}{k^2 - m_q^2} \right] + \dots, \end{aligned} \quad (18)$$

where the subscript q denotes either a light or a heavy quark, $G_{\mu\nu}$ and $F_{\mu\nu}$ are the external gluonic and electromagnetic field strength tensors, respectively, and e_q is the electric charge of the corresponding quark.

The nonperturbative contribution is obtained through the replacement

$$S_{\alpha\beta}(x) = -\frac{1}{4} (\Gamma_i)_{\alpha\beta} \bar{q}(x) \Gamma_i q(0), \quad (19)$$

where

$$\Gamma_i = \left\{ I, \gamma_5, \gamma_\mu, i\gamma_\mu \gamma_5, \frac{\sigma_{\mu\nu}}{\sqrt{2}} \right\} \quad (20)$$

forms a complete basis of Dirac matrices.

C. Nonlocal photon matrix elements

The nonperturbative replacement generates the nonlocal matrix elements

$$\begin{aligned} & \langle \gamma(q) | \bar{q}(x) \Gamma_i q(0) | 0 \rangle, \\ & \langle \gamma(q) | \bar{q}(x) \Gamma_i G_{\mu\nu} q(0) | 0 \rangle, \\ & \langle \gamma(q) | \bar{q}(x) \Gamma_i F_{\mu\nu} q(0) | 0 \rangle, \end{aligned} \quad (21)$$

which are parameterized in terms of photon light-cone distribution amplitudes of definite twist. In the present analysis, we employ the complete set of two-particle and three-particle photon distribution amplitudes up to twist four. Their explicit definitions and normalization conditions are collected in Appendix B and follow the convention of Ref. [20].

We implement continuum subtraction through the threshold s_0 . Matching the hadronic and OPE representations after the double Borel transformation gives

$$g_i M_i f_i \frac{m_P^2 f_P}{m_Q + m_s} \exp\left(-\frac{M_i^2}{\mathcal{M}_A^2} - \frac{m_P^2}{\mathcal{M}_P^2}\right) = \hat{\Pi}_i(\mathcal{M}^2, s_0). \quad (22)$$

The physical invariant amplitudes follow directly from the current rotation,

$$\begin{aligned} \hat{\Pi}_1 &= \sin \theta \hat{\Pi}_A + \cos \theta \hat{\Pi}_B, \\ \hat{\Pi}_2 &= \cos \theta \hat{\Pi}_A - \sin \theta \hat{\Pi}_B. \end{aligned} \quad (23)$$

The sum rules for the electric dipole form factor $g_i(q^2 = 0)$ involve the residues of the physical axial-vector mesons. These quantities are determined from the two-point correlation function,

$$\Pi_{\mu\nu}^{(i)}(p) = i \int d^4x e^{ip \cdot x} \langle 0 | T \{ J_\mu^{(i)}(x) J_\nu^{(i)\dagger}(0) \} | 0 \rangle, \quad (24)$$

where the interpolating currents $J_\mu^{(i)}$ are defined in Eq. (5).

Following the standard QCD sum-rule procedure, the hadronic and OPE representations are matched after a Borel transformation with parameter $\mathcal{M}_f^2$. The resulting physical-state projections are

$$f_1^2 M_1^2 e^{-M_1^2/\mathcal{M}_f^2} = \sin^2 \theta \tilde{\Pi}_{AA} + 2 \sin \theta \cos \theta \tilde{\Pi}_{AB} + \cos^2 \theta \tilde{\Pi}_{BB}, \quad (25)$$

and

$$f_2^2 M_2^2 e^{-M_2^2/\mathcal{M}_f^2} = \cos^2 \theta \tilde{\Pi}_{AA} - 2 \sin \theta \cos \theta \tilde{\Pi}_{AB} + \sin^2 \theta \tilde{\Pi}_{BB}, \quad (26)$$

where $\tilde{\Pi}_{AA}$, $\tilde{\Pi}_{AB}$, and $\tilde{\Pi}_{BB}$ represent the Borel-transformed invariant amplitudes obtained from the corresponding two-point correlation functions.

The construction of the mixed-current two-point sum rules follows Ref. [21]; their lengthy invariant amplitudes are not repeated here.

Using the residues obtained from Eqs. (25) and (26), the transition form factors are extracted from the light-cone QCD sum rules.

For the physical states one finds

$$g_1 = \frac{m_Q + m_s}{m_P^2 f_P M_1 f_1} \exp \left(\frac{M_1^2}{\mathcal{M}_A^2} + \frac{m_P^2}{\mathcal{M}_P^2} \right) \left(\sin \theta \hat{\Pi}_A + \cos \theta \hat{\Pi}_B \right), \quad (27)$$

and

$$g_2 = \frac{m_Q + m_s}{m_P^2 f_P M_2 f_2} \exp \left(\frac{M_2^2}{\mathcal{M}_A^2} + \frac{m_P^2}{\mathcal{M}_P^2} \right) \left(\cos \theta \hat{\Pi}_A - \sin \theta \hat{\Pi}_B \right), \quad (28)$$

where $\hat{\Pi}_A$ and $\hat{\Pi}_B$ are decomposed explicitly in Appendix A 1.

These expressions constitute the principal theoretical result of this work and enable the determination of the radiative transition form factors once the input parameters and photon distribution amplitudes are specified.

The electric dipole transition form factors obtained from the light-cone QCD sum rules can be used to determine the radiative decay widths of the physical axial-vector mesons.

The partial decay width for the process

$$A_i \rightarrow P\gamma$$

is given by

$$\Gamma(A_i \rightarrow P\gamma) = \frac{\alpha_{\text{em}}}{3} g_i^2 \left(\frac{M_i^2 - m_P^2}{2M_i} \right)^3, \quad (29)$$

where α_{em} denotes the electromagnetic fine-structure constant.

Once the transition form factors are determined, Eq. (29) gives the radiative decay widths of the physical heavy axial-vector mesons. Their numerical analysis is presented in the next section.

TABLE I: External inputs and physical mixed-current residues used in the numerical analysis.
Photon-DA normalizations are quoted at $\mu = 1$ GeV.

Parameter	Value
m_c, m_b, m_s	1.27(2), 4.18(3), 0.093(11) GeV [22]
$M_{D_{s1}(2460)}, M_{D_{s1}(2536)}, m_{D_s}$	2.4595, 2.53511, 1.96835 GeV [22]
$M_{B_{s1}(5830)}, m_{B_s}$	5.82870, 5.36692 GeV [6, 22]
$M_{B_{s1}(5750)}$	5.750(26) GeV [26]
f_{D_s}, f_{B_s}	0.2499, 0.2303 GeV [23]
$f_1^{(D_s)}, f_2^{(D_s)}$	0.405, 0.168 GeV
$f_1^{(B_s)}, f_2^{(B_s)}$	0.536, 0.089 GeV
$\chi_s(1 \text{ GeV})$	+3.15(30) GeV ⁻² [24]
	<i>Legacy susceptibility normalization; comparison only.</i>
$f_{\gamma,s}^\perp(1 \text{ GeV})$	-55.1(1.9) MeV [25]
	<i>Preferred lattice-normalized input.</i>

III. NUMERICAL ANALYSIS

The numerical analysis requires the quark and hadron masses, pseudoscalar decay constants, photon distribution-amplitude parameters, and the auxiliary parameters entering the sum rules. The external inputs used in the calculation are collected in Table I. The quark and hadron masses are taken from the Particle Data Group [22], while the pseudoscalar decay constants are taken from the lattice averages summarized by FLAG [23]. For the photon distribution amplitudes, we use the BBK/Rohrwild convention [20, 24].

For the normalization of the leading-twist photon distribution amplitude, we consider both the conventional susceptibility normalization proportional to $\chi_s \langle \bar{s}s \rangle$ and the recent lattice determination of $f_{\gamma,s}^\perp$ [25]. The physical mixed-current residues entering the final normalization are obtained from Eqs. (25) and (26) at the preferred sector-specific mixing angles.

The preferred predictions use the directly normalized lattice input $f_{\gamma,s}^\perp$; the susceptibility parameter χ_s is retained only for comparison with the traditional Rohrwild/BBK normalization and for quantifying the reduction of the leading-twist normalization uncertainty. The lattice-normalized input substantially reduces this uncertainty, although the cancellation-sensitive channels remain controlled by the relative interference of the two current contributions and by their residual mixing-angle dependence.

The Borel windows and continuum thresholds used in the numerical analysis are listed in Table II. The working regions are chosen according to the standard QCD sum-rule criteria. The extracted form factors should exhibit a sufficiently weak dependence on the Borel parameter, the continuum contribution should remain under control, and higher-twist terms should not dominate the OPE.

TABLE II: Final Borel windows and continuum-threshold intervals used in the Monte Carlo analysis.

Channel	$\mathcal{M}^2$ window	s_0 window
$D_{s1}(2460), D_{s1}(2536) \rightarrow D_s \gamma$	3.0–4.5 GeV ²	7.5–8.5 GeV ²
$B_{s1}(5750) \rightarrow B_s \gamma$	10.0–14.0 GeV ²	36.5–40.5 GeV ²
$B_{s1}(5830) \rightarrow B_s \gamma$	10.0–14.0 GeV ²	38.0–41.0 GeV ²

Since the initial and final mesons in each channel are relatively close in mass, we adopt the symmetric choice $\mathcal{M}_A^2 = \mathcal{M}_P^2$.

For the preferred numerical analysis, we use the sector-specific mixing angles obtained in the dedicated QCD sum-rule analysis of Ref. [27],

$$\theta_{D_s} = 26.6^\circ \pm 0.6^\circ, \quad \theta_{B_s} = 38.5^\circ \pm 0.1^\circ. \quad (30)$$

These Gaussian priors are used in determining the final central values and uncertainty intervals. To illustrate the sensitivity of the cancellation mechanism independently of these priors, we also perform the broader scan

$$25^\circ \leq \theta \leq 45^\circ.$$

This broad scan is used only as a benchmark and is not included in the preferred final uncertainty estimate.

Figure 1 shows the dependence of the form factors on the Borel parameter for representative values of the continuum threshold. The residual dependence on $\mathcal{M}^2$ and s_0 is included in the quoted uncertainties.

A second important source of sensitivity is the mixing angle. Since the physical amplitudes are linear combinations of the two basis-current contributions, variations in θ may either enhance or suppress the resulting transition amplitude. This effect is expected to be mild when the two components add with the same sign, but can become substantial when they have opposite signs.

Figure 2 displays the dependence of the radiative widths on the mixing angle. The minima in the higher-state curves make the cancellation-sensitive regions explicit. This figure represents the broad sensitivity test rather than the angle priors used for the final results.

The final uncertainties are obtained by varying the input parameters entering the calculation. Quantities with quoted errors are sampled within their stated uncertainties, while $\mathcal{M}^2$ and s_0 are varied over the intervals in Table II. The mixing angles are sampled from the sector-specific Gaussian priors in Eq. (30). For each set of input parameters, the transition form factor is calculated from




FIG. 1: Borel-parameter dependence of the transition form factors for $D_{s1}(2460) \rightarrow D_s\gamma$, $D_{s1}(2536) \rightarrow D_s\gamma$, $B_{s1}(5750) \rightarrow B_s\gamma$, and $B_{s1}(5830) \rightarrow B_s\gamma$. The curves correspond to representative values of the continuum threshold s_0 .

the corresponding light-cone QCD sum rule and the radiative width is then obtained from Eq. (29). We quote the median together with the 16th and 84th percentiles.

This procedure naturally yields asymmetric uncertainties. Such an uncertainty estimate is particularly appropriate for the $D_{s1}(2536) \rightarrow D_s\gamma$ and $B_{s1}(5830) \rightarrow B_s\gamma$ channels, where cancellations at the amplitude level generate strongly asymmetric distributions.

The resulting form factors and radiative widths obtained with the preferred lattice-normalized photon input are summarized in Table III.

The form-factor entries in Table III are the medians and percentile intervals of the sampled magnitudes $|g_i|$, rather than the absolute values of the medians of the signed g_i distributions. The decay width is evaluated sample by sample from the signed form factor through Eq. (29); it is not reconstructed from the displayed median of $|g_i|$.

Table III shows a clear suppression of the higher charm-strange state. The $D_{s1}(2536) \rightarrow D_s\gamma$




FIG. 2: Dependence of the radiative decay widths on the mixing angle θ . The charm-strange and bottom-strange channels are shown in the left and right panels, respectively. The vertical dashed line marks $\theta = 35.3^\circ$, the heavy-quark-limit reference value. The 25° – 45° range is shown only as a sensitivity test.

TABLE III: Transition form-factor magnitudes and radiative widths obtained with the lattice-normalized leading-twist photon input and the sector-specific mixing-angle priors of Eq. (30). The quoted values are reported as the median, with the lower and upper deviations set by the 16th and 84th percentiles, respectively. The form-factor column reports the distribution of $|g_i|$.

Channel	$ g_i $ (GeV $^{-1}$)	Γ (keV)
$D_{s1}(2460) \rightarrow D_s \gamma$	$0.408^{+0.047}_{-0.042}$	$34.96^{+8.57}_{-6.90}$
$D_{s1}(2536) \rightarrow D_s \gamma$	$0.0564^{+0.0506}_{-0.0392}$	$0.99^{+2.57}_{-0.90}$
$B_{s1}(5750) \rightarrow B_s \gamma$	$0.303^{+0.034}_{-0.031}$	$11.34^{+2.72}_{-2.20}$
$B_{s1}(5830) \rightarrow B_s \gamma$	$0.203^{+0.028}_{-0.028}$	$8.70^{+2.53}_{-2.23}$

amplitude remains cancellation-sensitive even with the narrow preferred angle prior, producing a strongly asymmetric relative uncertainty. In the bottom-strange sector, the two widths are comparable within their uncertainties.

For the preferred $f_{\gamma,s}^\perp$ normalization, the two mixed basis-current contributions to the normalized $D_{s1}(2460) \rightarrow D_s \gamma$ amplitude are approximately -0.0715 and -0.205 ; they therefore add constructively. For $D_{s1}(2536) \rightarrow D_s \gamma$, the corresponding contributions are -0.0917 and $+0.132$, so that the physical amplitude is determined by their difference. The small value of the resulting amplitude is therefore a direct consequence of destructive interference between the two mixed components.

These amplitude components provide the LCSR realization of the cancellation mechanism discussed in Refs. [18, 19].

The higher bottom-strange amplitude also remains sensitive to the relative contributions of the two basis currents. With the preferred sector-specific mixing-angle prior, however, the $B_{s1}(5750)$ and $B_{s1}(5830)$ widths are comparable within their uncertainties. We emphasize that $B_{s1}(5750)$ is

TABLE IV: Comparison of our predictions for radiative decay widths with results available in the literature. All widths are in keV.

Publication	$D_{s1}(2460) \rightarrow D_s \gamma$	$D_{s1}(2536) \rightarrow D_s \gamma$	$B_{s1}(5750) \rightarrow B_s \gamma$	$B_{s1}(5830) \rightarrow B_s \gamma$
[10]	19–29	–	–	–
[28, 29]	3.3	–	–	–
[8]	6.2	15	–	–
[30]	5.08	–	–	–
[9]	6.2	15	–	–
[7]	10.3–17.2	25.2–31.1	–	–
[14]	13.2	61.2	–	–
[11]	12.8	54.5	–	–
[15]	3.53–3.61	18.18–18.85	–	–
[31]	–	1.6 ± 2.3	–	–
[16]	–	–	37	27
[12]	–	–	70.6	47.8
[13]	–	–	97.7	56.6
[19]	297	12	47	28
This work	$34.96^{+8.57}_{-6.90}$	$0.99^{+2.57}_{-0.90}$	$11.34^{+2.72}_{-2.20}$	$8.70^{+2.53}_{-2.23}$

The LCSR and VMD estimates are quoted from Ref. [10]; the latter is based on Refs. [28, 29]. Bondar and Milstein [19] denote the bottom-strange states as $B_{s1}(5748)$ and $B_{s1}(5829)$; here they are identified with the corresponding $B_{s1}(5750)$ and $B_{s1}(5830)$ channels.

not an experimentally established resonance. In the present analysis it is included as a theoretical configuration motivated by predictions for the lower $J^P = 1^+$ bottom-strange state, with the mass input $M_{B_{s1}(5750)} = 5.750(26)$ GeV.

For comparison with earlier photon-LCSR calculations, we also repeated the analysis using the conventional susceptibility-normalized leading-twist input $\chi_s \langle \bar{s}s \rangle$. These results are not used for the final predictions in Table III, but they provide a useful estimate of the sensitivity to the normalization of the leading-twist photon matrix element. The lattice quantity $f_{\gamma,s}^\perp$ replaces the product $\chi_s \langle \bar{s}s \rangle$ by a directly normalized photon matrix element and is therefore used in our preferred numerical analysis.

It is also useful to compare our predictions with selected results obtained using other theoretical approaches. This comparison is presented in Table IV.

The comparison shows that the absolute widths remain model dependent. In particular, the potential-model predictions of Ref. [19] are larger than our preferred LCSR results in all four channels. Nevertheless, the qualitative pattern in the charm-strange system is similar: the higher axial-vector state is sensitive to cancellation between the two mixed spin components.

The difference in the absolute widths is not unexpected. In the potential-model treatment, the




FIG. 3: Monte Carlo distributions of the radiative decay widths from the broad $25^\circ \leq \theta \leq 45^\circ$ sensitivity scan. The solid and dashed vertical lines indicate the median and the 16th–84th percentile interval, respectively. These benchmark distributions are not used for the preferred final uncertainties.

radiative amplitudes depend on the meson wave functions and on sizable spin-dependent relativistic corrections. In the present LCSR calculation, the normalization is instead controlled by the photon distribution amplitudes, continuum subtraction, and the residues of the physical interpolating currents. In channels close to a cancellation, even a moderate change in the relative magnitude of the two components can lead to a large change in the decay width after the amplitude is squared.

For completeness, Fig. 3 shows the Monte Carlo distributions obtained in the broad $25^\circ \leq \theta \leq 45^\circ$ sensitivity scan. The long asymmetric tails in the cancellation-sensitive channels illustrate the strong dependence on the mixing angle. These benchmark distributions are not used to determine the preferred results quoted in Table III, which instead employ the Gaussian angle priors in Eq. (30).

Finally, the predicted width of the $D_{s1}(2536) \rightarrow D_s \gamma$ channel can be used to estimate its branching

fraction. Using $\Gamma_{\text{tot}}[D_{s1}(2536)] \simeq 0.92 \text{ MeV}$, the central value of our prediction gives

$$\mathcal{B}[D_{s1}(2536) \rightarrow D_s \gamma] \sim 1.1 \times 10^{-3}. \quad (31)$$

The 16th–84th percentile interval of the radiative width corresponds to an approximate branching-fraction range of 1.0×10^{-4} – 3.9×10^{-3} . The small predicted branching fraction makes this channel experimentally challenging, while also providing a useful target for dedicated searches.

These results also help explain the wide range of predictions found in the literature. For cancellation-sensitive transitions, relatively small changes in the mixing angle, current residues, or relative size of the two basis-current amplitudes can produce large changes in the decay width. Future measurements of $D_{s1}(2536) \rightarrow D_s \gamma$ and $B_{s1}(5830) \rightarrow B_s \gamma$, or improved upper limits on these channels, would therefore provide useful tests of the mixing mechanism in strange heavy axial-vector mesons.

IV. CONCLUSION

We have studied the radiative decays $D_{s1}(2460) \rightarrow D_s \gamma$, $D_{s1}(2536) \rightarrow D_s \gamma$, $B_{s1}(5750) \rightarrow B_s \gamma$, and $B_{s1}(5830) \rightarrow B_s \gamma$ within the light-cone QCD sum-rule framework. The calculation was carried out for two independent axial-vector interpolating currents, whose contributions were combined according to the 1P_1 – 3P_1 mixing of the physical 1^+ states. We retained the strange-quark-mass terms and used photon distribution amplitudes up to twist four. For the leading-twist photon distribution amplitude, we compared the conventional susceptibility normalization with the recent lattice-normalized input and used the latter for our preferred predictions.

The resulting radiative widths are

$$\Gamma[D_{s1}(2460) \rightarrow D_s \gamma] = 34.96^{+8.57}_{-6.90} \text{ keV}, \quad \Gamma[D_{s1}(2536) \rightarrow D_s \gamma] = 0.99^{+2.57}_{-0.90} \text{ keV}, \quad (32)$$

$$\Gamma[B_{s1}(5750) \rightarrow B_s \gamma] = 11.34^{+2.72}_{-2.20} \text{ keV}, \quad \Gamma[B_{s1}(5830) \rightarrow B_s \gamma] = 8.70^{+2.53}_{-2.23} \text{ keV}. \quad (33)$$

A pronounced hierarchy is found between the two charm-strange states. The $D_{s1}(2460) \rightarrow D_s \gamma$ prediction is stable at approximately 35 keV, whereas the $D_{s1}(2536) \rightarrow D_s \gamma$ width is strongly suppressed to approximately 1 keV. This hierarchy follows directly from the mixing structure: the two basis-current contributions add constructively for the lower state and destructively for the higher state. The result thus supports the cancellation mechanism discussed in Refs. [18, 19].

The preferred result for $D_{s1}(2536) \rightarrow D_s \gamma$ corresponds to a branching fraction of order 1.1×10^{-3}

when the measured total width is used. The large relative uncertainty of this estimate reflects the sensitivity of the amplitude to the cancellation between the two mixed components.

In the bottom-strange sector, the lower $B_{s1}(5750)$ state is included as a theoretically expected $J^P = 1^+$ configuration rather than an experimentally established resonance. Its predicted radiative width provides a useful benchmark for future studies of the lower axial-vector bottom-strange state. For the established $B_{s1}(5830)$, the preferred sector-specific-angle scenario gives a width of approximately 9 keV. Thus, this channel is not strongly suppressed in the preferred analysis. The broad 25° – 45° scan is presented separately only as a sensitivity test and is not used to define the final uncertainty.

The comparison with previous calculations shows that the absolute radiative widths remain model dependent, particularly in channels close to a cancellation. Nevertheless, the present analysis indicates that the relative size of the two mixed-current contributions is a central element in determining the decay pattern. Future measurements of $D_{s1}(2536) \rightarrow D_s \gamma$ and $B_{s1}(5830) \rightarrow B_s \gamma$, as well as improved limits on these channels, would therefore provide valuable tests of the 1P_1 – 3P_1 mixing structure and of the radiative dynamics of strange heavy axial-vector mesons.

Appendix A: OPE expressions and auxiliary definitions

This appendix collects the explicit Borel-transformed invariant amplitudes used in the main text. The charges are quoted in units of the positive elementary charge: $e_s = -1/3$, $e_c = 2/3$, and $e_b = -1/3$. Here m_Q and m_s denote quark masses, whereas M_i denotes a physical axial-vector mass. We use

$$d \equiv m_Q + m_s, \quad u_0 = \frac{\mathcal{M}_P^2}{\mathcal{M}_A^2 + \mathcal{M}_P^2}, \quad a_0 = 1 - u_0, \quad \mathcal{M}^2 = \frac{\mathcal{M}_A^2 \mathcal{M}_P^2}{\mathcal{M}_A^2 + \mathcal{M}_P^2}. \quad (\text{A1})$$

1. Support projections and invariant amplitudes

For an operator-weighted three-particle DA $X(\alpha_i, v)$, the momentum fraction entering the heavy-quark denominator is

$$a(\alpha_i, v) = \alpha_{\bar{q}} + v\alpha_g. \quad (\text{A2})$$

The support projection used below is

$$\mathcal{I}_z[X] = \int_0^z dv \int_0^{(1-z)/(1-v)} d\alpha_g X(1 - z - (1 - v)\alpha_g, z - v\alpha_g, \alpha_g, v)$$

$$+ \int_z^1 dv \int_0^{z/v} d\alpha_g X(1-z-(1-v)\alpha_g, z-v\alpha_g, \alpha_g, v). \quad (\text{A3})$$

Terms containing $1/(q \cdot x)$ require

$$\begin{aligned} \mathcal{L}_z[X] = & \int_0^{1-z} d\alpha_q \int_0^{\alpha_q} d\alpha'_q \int_0^z d\alpha_{\bar{q}} \frac{z-\alpha_{\bar{q}}}{(1-\alpha_q-\alpha_{\bar{q}})^2} X(\alpha_q, \alpha_{\bar{q}}, 1-\alpha_q-\alpha_{\bar{q}}) \\ & - \int_0^{1-z} d\alpha_q \int_z^{1-\alpha_q} d\beta \frac{z}{\beta^2} \int_0^\beta d\alpha'_{\bar{q}} X(\alpha_q, \alpha'_{\bar{q}}, 1-\alpha_q-\alpha'_{\bar{q}}) \\ & + \int_0^{1-z} d\alpha_q \frac{z}{(1-\alpha_q)^2} \int_0^{\alpha_q} d\alpha'_q \int_0^{1-\alpha'_q} d\alpha_{\bar{q}} X(\alpha'_q, \alpha_{\bar{q}}, 1-\alpha'_q-\alpha_{\bar{q}}). \end{aligned} \quad (\text{A4})$$

The electromagnetic three-particle DAs $\mathcal{S}_\gamma$ and $\mathcal{T}_4^\gamma$ are the scalar and tensor functions multiplying the corresponding $F_{\mu\nu}$ operator structures. Their basis-current weights are

$$\mathcal{F}_A^\gamma = (1-2v)\mathcal{S}_\gamma - \mathcal{T}_4^\gamma, \quad \mathcal{F}_B^\gamma = \frac{m_Q}{d}(\mathcal{S}_\gamma - \mathcal{T}_4^\gamma). \quad (\text{A5})$$

$$\begin{aligned} \hat{\Pi}_{A(B)} = & \hat{\Pi}_{A(B)}^{\text{pert}} + \hat{\Pi}_{A(B)}^{\text{twist2}} + \hat{\Pi}_{A(B)}^{\text{twist3}} \\ & + \hat{\Pi}_{A(B)}^{\text{twist4}} + \hat{\Pi}_{A(B)}^{3\text{p,g}} + \hat{\Pi}_{A(B)}^{3\text{p},\gamma}. \end{aligned} \quad (\text{A6})$$

The individual components are

$$\begin{aligned} \hat{\Pi}_A^{\text{pert}} &= \int_{(m_Q+m_s)^2}^{s_0} ds e^{-s/\mathcal{M}^2} \rho_A^P(s), \\ \hat{\Pi}_A^{\text{twist2}} &= e_s \langle \bar{s}s \rangle \left(e^{-m_Q^2/\mathcal{M}^2} - e^{-s_0/\mathcal{M}^2} \right) \mathcal{M}^2 \chi_s \phi_\gamma(u_0), \\ \hat{\Pi}_A^{\text{twist3}} &= e_s f_{3\gamma} m_Q e^{-m_Q^2/\mathcal{M}^2} \Psi^{(v)}(u_0), \\ \hat{\Pi}_A^{\text{twist4}} &= \frac{e_s \langle \bar{s}s \rangle e^{-m_Q^2/\mathcal{M}^2}}{4} [\mathcal{A}(u_0) + 2\mathcal{B}(u_0)] \left(1 + \frac{m_Q^2}{\mathcal{M}^2} \right), \\ \hat{\Pi}_A^{3\text{p,g}} &= e_s \langle \bar{s}s \rangle e^{-m_Q^2/\mathcal{M}^2} \mathcal{I}_{F_1}(u_0). \end{aligned} \quad (\text{A7})$$

The momentum fractions u_0 and $a_0 = 1 - u_0$ are defined in Eq. (13).

The superscripts label the perturbative, two-particle twist-2-4, gluonic three-particle, and electromagnetic three-particle contributions, respectively.

The perturbative axial-current density and the gluonic three-particle convolution are

$$\rho_a^A(s; m_a, m_b, e_a) = - \frac{3e_a}{8\pi^2} \left[2m_a \ln \left(\frac{s - m_b^2 + m_a^2 - \lambda^{1/2}(s, m_b^2, m_a^2)}{s - m_b^2 + m_a^2 + \lambda^{1/2}(s, m_b^2, m_a^2)} \right) \right]$$

$$+ (m_b - m_a) \frac{m_b^2 - m_a^2 - s}{s^2} \lambda^{1/2}(s, m_b^2, m_a^2) \Bigg],$$

$$\rho_A^P(s) = \rho_a^A(s; m_s, m_Q, e_s) - \rho_a^A(s; m_Q, m_s, e_Q). \quad (\text{A8})$$

$$\begin{aligned} \mathcal{I}_{F_1}(u_0) &= \int_0^{1-u_0} dv \int_0^{u_0/(1-v)} d\alpha_g F_1(u_0 - (1-v)\alpha_g, 1 - u_0 - v\alpha_g, \alpha_g, v) \\ &\quad + \int_{1-u_0}^1 dv \int_0^{(1-u_0)/v} d\alpha_g F_1(u_0 - (1-v)\alpha_g, 1 - u_0 - v\alpha_g, \alpha_g, v), \\ F_1(\alpha_i, v) &= \mathcal{S} + \tilde{\mathcal{S}} - \mathcal{T}_1 - \mathcal{T}_2 + \mathcal{T}_3 + \mathcal{T}_4 + 2v(-\mathcal{S} - \mathcal{T}_3 + \mathcal{T}_2). \end{aligned} \quad (\text{A9})$$

Here $\lambda(a, b, c) = (a - b - c)^2 - 4bc$ is the Källén function.

For the tensor current, the perturbative density entering the diagonal double-dispersion representation is

$$\begin{aligned} \rho_B^P(s) &= -\frac{3}{8\pi^2} \left\{ e_s \left[2 \frac{m_Q m_s}{m_Q + m_s} \ln \left(\frac{s - m_Q^2 + m_s^2 - \sqrt{\lambda(s, m_Q^2, m_s^2)}}{s - m_Q^2 + m_s^2 + \sqrt{\lambda(s, m_Q^2, m_s^2)}} \right) \right. \right. \\ &\quad \left. \left. + \frac{s}{m_Q + m_s} \frac{m_Q^2 - m_s^2 - s}{s^2} \sqrt{\lambda(s, m_Q^2, m_s^2)} \right] \right. \\ &\quad \left. - e_Q \left[2 \frac{2m_Q^2 - m_Q m_s}{m_Q + m_s} \ln \left(\frac{s - m_s^2 + m_Q^2 - \sqrt{\lambda(s, m_s^2, m_Q^2)}}{s - m_s^2 + m_Q^2 + \sqrt{\lambda(s, m_s^2, m_Q^2)}} \right) \right. \right. \\ &\quad \left. \left. - \frac{s}{m_Q + m_s} \frac{m_s^2 - m_Q^2 - s}{s^2} \sqrt{\lambda(s, m_s^2, m_Q^2)} \right] \right\}. \end{aligned} \quad (\text{A10})$$

The tensor-current twist-3 contribution is

$$\hat{\Pi}_B^{\text{twist3}} = \frac{e_s f_{3\gamma} e^{-m_Q^2/\mathcal{M}^2}}{d} \left[m_Q^2 \psi^{(v)}(u_0) - \frac{\mathcal{M}^2}{2} \frac{d}{da} \{ (1-a) \psi^{(v)}(a) \} \Big|_{a=u_0} \right]. \quad (\text{A11})$$

The twist-2 and twist-4 tensor-current contributions obey

$$\hat{\Pi}_B^{\text{twist2}} = \frac{m_Q}{d} \hat{\Pi}_A^{\text{twist2}}, \quad \hat{\Pi}_B^{\text{twist4}} = \frac{m_Q}{d} \hat{\Pi}_A^{\text{twist4}}. \quad (\text{A12})$$

The electromagnetic three-particle contributions are

$$\begin{aligned}\widehat{\Pi}_A^{3p,\gamma} &= e_Q \langle \bar{s}s \rangle \left(e^{-m_Q^2/\mathcal{M}^2} - e^{-s_0/\mathcal{M}^2} \right) \left\{ \mathcal{I}_{a_0}[\mathcal{F}_A^\gamma] + 2 \frac{d}{dz} \mathcal{L}_z[\mathcal{T}_4^\gamma] \Big|_{z=a_0} \right\}, \\ \widehat{\Pi}_B^{3p,\gamma} &= e_Q \langle \bar{s}s \rangle \left(e^{-m_Q^2/\mathcal{M}^2} - e^{-s_0/\mathcal{M}^2} \right) \frac{m_Q}{d} \left\{ \mathcal{I}_{a_0}[\mathcal{S}_\gamma - \mathcal{T}_4^\gamma] + 2 \frac{d}{dz} \mathcal{L}_z[\mathcal{T}_4^\gamma] \Big|_{z=a_0} \right\}.\end{aligned}\quad (\text{A13})$$

Here $a_0 = 1 - u_0$ and the convolution operators are defined in Eqs. (A3) and (A4).

Appendix B: Photon distribution amplitudes and auxiliary functions

For the nonperturbative photon matrix elements we use the photon distribution amplitudes (DAs) in the notation of Rohrwild [24], which is based on the BBK classification [20]. We collect here the functions and auxiliary combinations entering the sum rules of Sec. II. The variable u denotes the quark momentum fraction, $\bar{u} = 1 - u$, and $\alpha_q + \alpha_{\bar{q}} + \alpha_g = 1$ for the three-particle DAs.

The leading-twist two-particle DA is taken in the asymptotic form

$$\phi_\gamma(u) = 6u\bar{u}. \quad (\text{B1})$$

The twist-3 vector DA is

$$\psi^{(v)}(u) = -20u\bar{u}(2u-1) + \frac{15}{16} (\omega_\gamma^A - 3\omega_\gamma^V) u\bar{u}(2u-1) [7(2u-1)^2 - 3]. \quad (\text{B2})$$

For the twist-4 two-particle DAs we use

$$\begin{aligned}\mathcal{A}(u) &= 40u\bar{u} (3\kappa - \kappa^+ + 1) + 8 (\zeta_2^+ - 3\zeta_2) \left[u\bar{u}(2 + 13u\bar{u}) \right. \\ &\quad \left. + u^3(10 - 15u + 6u^2) \ln u + \bar{u}^3(10 - 15\bar{u} + 6\bar{u}^2) \ln \bar{u} \right],\end{aligned}\quad (\text{B3})$$

$$\mathcal{B}(u) = 40(1 + 3\kappa^+) \int_0^u d\alpha (u - \alpha) \left[-\frac{1}{2} + \frac{3}{2}(2\alpha - 1)^2 \right]. \quad (\text{B4})$$

The twist-4 three-particle DAs used in the numerical analysis are

$$\begin{aligned}\mathcal{S}(\alpha_i) &= 30\alpha_g^2 \left[(\kappa + \kappa^+)(1 - \alpha_g) + (\zeta_1 + \zeta_1^+)(1 - \alpha_g)(1 - 2\alpha_g) \right. \\ &\quad \left. + \zeta_2 \{ 3(\alpha_{\bar{q}} - \alpha_q)^2 - \alpha_g(1 - \alpha_g) \} \right],\end{aligned}\quad (\text{B5})$$

$$\tilde{\mathcal{S}}(\alpha_i) = -30\alpha_g^2 \left[(\kappa - \kappa^+)(1 - \alpha_g) + (\zeta_1 - \zeta_1^+)(1 - \alpha_g)(1 - 2\alpha_g) \right]$$

$$+ \zeta_2 \{ 3(\alpha_{\bar{q}} - \alpha_q)^2 - \alpha_g(1 - \alpha_g) \} \Big], \quad (\text{B6})$$

$$\mathcal{T}_1(\alpha_i) = -120(3\zeta_2 + \zeta_2^+)(\alpha_q - \alpha_{\bar{q}})\alpha_{\bar{q}}\alpha_q\alpha_g, \quad (\text{B7})$$

$$\mathcal{T}_2(\alpha_i) = 30\alpha_g^2(\alpha_q - \alpha_{\bar{q}}) \left[\kappa - \kappa^+ + (\zeta_1 - \zeta_1^+)(1 - 2\alpha_g) + \zeta_2(3 - 4\alpha_g) \right], \quad (\text{B8})$$

$$\mathcal{T}_3(\alpha_i) = -120(3\zeta_2 - \zeta_2^+)(\alpha_q - \alpha_{\bar{q}})\alpha_{\bar{q}}\alpha_q\alpha_g, \quad (\text{B9})$$

$$\mathcal{T}_4(\alpha_i) = 30\alpha_g^2(\alpha_q - \alpha_{\bar{q}}) \left[\kappa + \kappa^+ + (\zeta_1 + \zeta_1^+)(1 - 2\alpha_g) + \zeta_2(3 - 4\alpha_g) \right]. \quad (\text{B10})$$

The two-particle functions entering the Borel-transformed amplitudes are used through

$$\Psi^{(v)}(u_0) = \int_0^{u_0} du \psi^{(v)}(u), \quad (\text{B11})$$

$$\mathcal{B}(u_0) = -8 \int_0^{u_0} du (u_0 - u) h_\gamma(u), \quad (\text{B12})$$

where Eq. (B12) is the notation dictionary connecting the Rohrwild/BBK convention to the form used in the single-current benchmark of Ref. [10].

For the three-particle contribution we define

$$\begin{aligned} \mathcal{I}_F(u_0) = & \int_0^{1-u_0} dv \int_0^{u_0/(1-v)} d\alpha_g F(u_0 - (1-v)\alpha_g, 1 - u_0 - v\alpha_g, \alpha_g, v) \\ & + \int_{1-u_0}^1 dv \int_0^{(1-u_0)/v} d\alpha_g F(u_0 - (1-v)\alpha_g, 1 - u_0 - v\alpha_g, \alpha_g, v), \end{aligned} \quad (\text{B13})$$

with

$$F = \mathcal{S} + \tilde{\mathcal{S}} - \mathcal{T}_1 - \mathcal{T}_2 + \mathcal{T}_3 + \mathcal{T}_4 + 2v(-\mathcal{S} - \mathcal{T}_3 + \mathcal{T}_2). \quad (\text{B14})$$

All three-particle DAs in Eq. (B14) are evaluated at $(\alpha_q, \alpha_{\bar{q}}, \alpha_g)$, with the arguments specified in Eq. (B13). In the numerical analysis we use the symmetric point $u_0 = 1/2$.

In the preferred modern scenario we replace the legacy normalization $\chi_s \langle \bar{s}s \rangle$ of the leading-twist matrix element by the lattice-normalized input $f_{\gamma,s}^\perp$. The DA shapes in this appendix are otherwise kept fixed in the same Rohrwild/BBK convention.

-
- [1] BaBar Collaboration, B. Aubert *et al.*, “A study of the $D_{sJ}^*(2317)^+$ and $D_{sJ}(2460)^+$ mesons in inclusive $c\bar{c}$ production near $\sqrt{s} = 10.6$ GeV,” Phys. Rev. D **74** (2006) 032007, [hep-ex/0604030].
- [2] BaBar Collaboration, B. Aubert *et al.*, “Study of $B \rightarrow D^{(*)} D_{s(J)}^{(*)}$ decays and measurement of D_s^- and

TABLE V: Photon-DA parameters used at $\mu = 1$ GeV in Rohrwild notation.

Parameter	Value	Comment
χ_s	$3.15 \pm 0.30 \text{ GeV}^{-2}$	legacy susceptibility input
$f_{3\gamma}$	$-(4.0 \pm 2.0) \times 10^{-3} \text{ GeV}^2$	twist-3 coupling
ω_γ^A	-2.1 ± 1.0	twist-3 shape
ω_γ^V	3.8 ± 1.8	twist-3 shape
κ	0.15	twist-4 shape
κ^+	-0.05	twist-4 shape
ζ_1	0.40	twist-4 shape
ζ_1^+	0	twist-4 shape
ζ_2	0.30	twist-4 shape
ζ_2^+	0	twist-4 shape

$D_{sJ}(2460)^-$ branching fractions,” Phys. Rev. D **74** (2006) 031103, [hep-ex/0605036].

- [3] Belle Collaboration, V. Balagura *et al.*, “Observation of $D_{s1}(2536)^+ \rightarrow D^+ \pi^- K^+$ and angular decomposition of $D_{s1}(2536)^+ \rightarrow D^{*+} K_S^0$,” Phys. Rev. D **77** (2008) 032001, [0709.4184].
- [4] BaBar Collaboration, J. P. Lees *et al.*, “Measurement of the $D_{s1}(2536)$ mass and width,” Phys. Rev. D **83** (2011) 072003.
- [5] LHCb Collaboration, R. Aaij *et al.*, “First observation of the decay $B_{s2}^*(5840)^0 \rightarrow B^{*+} K^-$ and studies of excited B_s^0 mesons,” Phys. Rev. Lett. **110** (2013) 151803, [1211.5994].
- [6] CMS Collaboration, A. M. Sirunyan *et al.*, “Studies of $B_{s2}^*(5840)^0$ and $B_{s1}(5830)^0$ mesons including the observation of the $B_{s2}^*(5840)^0 \rightarrow B^0 K_S^0$ decay in proton-proton collisions at $\sqrt{s} = 8 \text{ TeV}$,” Eur. Phys. J. C **78** no. 11, (2018) 939, [1809.03578].
- [7] J. L. Goity and W. Roberts, “Radiative transitions in heavy mesons in a relativistic quark model,” Phys. Rev. D **64** (2001) 094007, [hep-ph/0012314].
- [8] S. Godfrey, “Testing the nature of the $D_{sJ}^*(2317)^+$ and $D_{sJ}(2463)^+$ states using radiative transitions,” Phys. Lett. B **568** (2003) 254–260, [hep-ph/0305122].
- [9] S. Godfrey, “Properties of the charmed P -wave mesons,” Phys. Rev. D **72** (2005) 054029, [hep-ph/0508078].
- [10] P. Colangelo, F. De Fazio, and A. Ozpineci, “Radiative transitions of $D_{sJ}^*(2317)$ and $D_{sJ}(2460)$,” Phys. Rev. D **72** (2005) 074004, [hep-ph/0505195].
- [11] S. F. Radford, W. W. Repko, and M. J. Saelim, “Potential model calculations and predictions for $c\bar{s}$ quarkonia,” Phys. Rev. D **80** (2009) 034012, [0903.0551].
- [12] S. Godfrey, K. Moats, and E. S. Swanson, “ B and B_s meson spectroscopy,” Phys. Rev. D **94** (2016) 054025, [1607.02169].
- [13] Q.-F. Lü, T.-T. Pan, Y.-Y. Wang, E. Wang, and D.-M. Li, “Excited bottom and bottom-strange

- mesons in the quark model*,” Phys. Rev. D **94** (2016) 074012, [1607.02812].
- [14] N. Green, W. W. Repko, and S. F. Radford, “*Note on predictions for $c\bar{s}$ quarkonia using a three-loop static potential*,” Nucl. Phys. A **958** (2017) 71–77, [1605.06393].
- [15] S.-F. Chen, J. Liu, H.-Q. Zhou, and D.-Y. Chen, “*Electric transitions of the charmed-strange mesons in a relativistic quark model*,” Eur. Phys. J. C **80** (2020) 290, [2003.07988].
- [16] Q. Li, R.-H. Ni, and X.-H. Zhong, “*Towards establishing an abundant B and B_s spectrum up to the second orbital excitations*,” Phys. Rev. D **103** (2021) 116010, [2102.03694].
- [17] B. Pullin and R. Zwicky, “*Radiative decays of heavy-light mesons and the decay constants $f_{H,H^*,H_1}^{(T)}$* ,” JHEP **09** (2021) 023, [2106.13617].
- [18] A. E. Bondar, “*Why We Do Not See the Radiative Decays of the $D_{s1}^+(2536)$ Meson?*,” JETP Lett. **121** no. 4, (2025) 231–235, [2311.15078].
- [19] A. E. Bondar and A. I. Milstein, “*Phenomenology of D_{s1} mesons radiative transitions*,” Phys. Rev. D **111** no. 11, (2025) 114019, [2505.01856].
- [20] P. Ball, V. M. Braun, and N. Kivel, “*Photon distribution amplitudes in QCD*,” Nucl. Phys. B **649** (2003) 263–296, [hep-ph/0207307].
- [21] T. M. Aliev, A. Ozpineci, and V. Zamiralov, “*Mixing Angle of Hadrons in QCD: A New View*,” Phys. Rev. D **83** (2011) 016008, [1007.0814].
- [22] Particle Data Group Collaboration, F. Takahashi *et al.*, “*Review of Particle Physics*,” Int. J. Mod. Phys. A **41** no. 22, (2026) 2630011.
- [23] Flavour Lattice Averaging Group (FLAG) Collaboration, Y. Aoki *et al.*, “*FLAG Review 2024*,” Phys. Rev. D **113** no. 1, (2026) 014508, [2411.04268].
- [24] J. Rohrwild, “*Determination of the magnetic susceptibility of the quark condensate using radiative heavy meson decays*,” JHEP **09** (2007) 073, [0708.1405].
- [25] S. Bacchio, D. Becirevic, G. Gagliardi, and F. Sanfilippo, “*Lattice QCD determination of the normalization of the leading-twist photon distribution amplitude and susceptibility of the quark condensate*,” Phys. Rev. D **110** (2024) 034511, [2404.10644].
- [26] C. B. Lang, D. Mohler, S. Prelovsek, and R. M. Woloshyn, “*Predicting positive parity B_s mesons from lattice QCD*,” Phys. Lett. B **750** (2015) 17–21, [1501.01646].
- [27] T. M. Aliev, S. Bilmis, and M. Savci, “*Mixing angle between 3P_1 and 1P_1 states in heavy axial vector mesons within the QCD sum rules framework*,” Phys. Rev. D **111** no. 1, (2025) 014005, [2408.08014].
- [28] P. Colangelo, F. De Fazio, and R. Ferrandes, “*Excited charmed mesons: Observations, analyses and puzzles*,” Mod. Phys. Lett. A **19** (2004) 2083–2102, [hep-ph/0407137].
- [29] P. Colangelo and F. De Fazio, “*Understanding $D_{sJ}(2317)$* ,” Phys. Lett. B **570** (2003) 180–184,

[hep-ph/0305140].

- [30] W. A. Bardeen, E. J. Eichten, and C. T. Hill, “*Chiral multiplets of heavy-light mesons*,” Phys. Rev. D **68** (2003) 054024, [hep-ph/0305049].
- [31] J. G. Korner, D. Pirjol, and K. Schilcher, “*Radiative decays of the P-wave charmed mesons*,” Phys. Rev. D **47** (1993) 3955–3961, [hep-ph/9212220].