
Magnetic moment μ/μ_N vs M^2

Nominal mixing angles | $+\chi$: $f_\perp = -55.1$ MeV




Thick curves: $\sqrt{s_0} - m_H = 0.50$ GeV (or nearest accepted threshold). Faint curves: other accepted s_0 values.

Only accepted windows are shown; points at different thresholds are never joined. Each point uses its own two-point residue.

Frozen CSV: commit 54e6958. Both sign pages use identical axis limits for each state.

Magnetic moment μ/μ_N vs M^2


Nominal mixing angles | $-\chi$: $f_\perp = +55.1$ MeV



Thick curves: $\sqrt{s_0} - m_H = 0.50$ GeV (or nearest accepted threshold). Faint curves: other accepted s_0 values.

Only accepted windows are shown; points at different thresholds are never joined. Each point uses its own two-point residue.


Frozen CSV: commit 54e6958. Both sign pages use identical axis limits for each state.



Thick curves: $\sqrt{s_0} - m_H = 0.50$ GeV (or nearest accepted threshold). Faint curves: other accepted s_0 values.

Only accepted windows are shown; points at different thresholds are never joined. Each point uses its own two-point residue.


Frozen CSV: commit 54e6958. Quadrupole total excludes charge: add Q_H=1 for Ds, 0 for Bs.



Thick curves: $\sqrt{s_0} - m_H = 0.50$ GeV (or nearest accepted threshold). Faint curves: other accepted s_0 values.

Only accepted windows are shown; points at different thresholds are never joined. Each point uses its own two-point residue.

Frozen CSV: commit 54e6958. Quadrupole total excludes charge: add Q_H=1 for Ds, 0 for Bs.




Thick curves: $\sqrt{s_0} - m_H = 0.50$ GeV (or nearest accepted threshold). Faint curves: other accepted s_0 values.

Only accepted windows are shown; points at different thresholds are never joined. Each point uses its own two-point residue.

Frozen CSV: commit 54e6958. Both sign pages use identical axis limits for each state.

Magnetic moment μ/μ_N vs M^2


$\theta = 90^\circ$: refitted pure-current windows | $-\chi$: $f_\perp = +55.1$ MeV



Thick curves: $\sqrt{s_0} - m_H = 0.50$ GeV (or nearest accepted threshold). Faint curves: other accepted s_0 values.

Only accepted windows are shown; points at different thresholds are never joined. Each point uses its own two-point residue.


Frozen CSV: commit 54e6958. Both sign pages use identical axis limits for each state.



Thick curves: $\sqrt{s_0} - m_H = 0.50$ GeV (or nearest accepted threshold). Faint curves: other accepted s_0 values.

Only accepted windows are shown; points at different thresholds are never joined. Each point uses its own two-point residue.

Frozen CSV: commit 54e6958. Quadrupole total excludes charge: add Q_H=1 for Ds, 0 for Bs.



Thick curves: $\sqrt{s_0} - m_H = 0.50$ GeV (or nearest accepted threshold). Faint curves: other accepted s_0 values.

Only accepted windows are shown; points at different thresholds are never joined. Each point uses its own two-point residue.

Frozen CSV: commit 54e6958. Quadrupole total excludes charge: add Q_H=1 for Ds, 0 for Bs.