

Working Notes for $D_{s1} \rightarrow D_s \gamma$ in Light-Cone QCD Sum Rules

Ds gamma project

July 2, 2026

Abstract

These notes are a living derivation for the light-cone QCD sum-rule calculation of $D_{s1}(2460) \rightarrow D_s \gamma$ and $D_{s1}(2536) \rightarrow D_s \gamma$. The purpose is pedagogical as well as practical: each step is written so that a master's student can follow the choices, conventions, and algebra before the final paper-level formulas are assembled.

1 Conventions

We use the mostly-minus metric

$$g_{\mu\nu} = \text{diag}(+1, -1, -1, -1), \quad (1)$$

and

$$\{\gamma_\mu, \gamma_\nu\} = 2g_{\mu\nu}, \quad \sigma_{\mu\nu} = \frac{i}{2}[\gamma_\mu, \gamma_\nu], \quad \gamma_5 = i\gamma^0\gamma^1\gamma^2\gamma^3. \quad (2)$$

The Levi-Civita convention is

$$\epsilon^{0123} = +1, \quad \epsilon_{0123} = -1, \quad (3)$$

so that

$$\text{Tr}[\gamma_5 \gamma_\mu \gamma_\nu \gamma_\alpha \gamma_\beta] = -4i \epsilon_{\mu\nu\alpha\beta}. \quad (4)$$

Our Fourier convention for a propagator is

$$S(x) = \int \frac{d^4k}{(2\pi)^4} e^{-ik \cdot x} \tilde{S}(k). \quad (5)$$

2 Currents and Momentum Assignment

The external momenta are

$$P = p + q, \quad (6)$$

where P is the initial axial-meson momentum, p is the final pseudoscalar D_s momentum, and q is the real-photon momentum.

The pseudoscalar interpolating current is

$$J_{D_s}(x) = \bar{s}(x) i\gamma_5 Q(x). \quad (7)$$

For the axial channel we use two independent currents,

$$J_{A\mu}^{(0)}(x) = \bar{s}(x) \gamma_\mu \gamma_5 Q(x), \quad (8)$$

$$J_{B\mu}^{(0)}(x) = \frac{i}{m_Q + m_s} \bar{s}(x) \sigma_{\mu\alpha} P^\alpha \gamma_5 Q(x). \quad (9)$$

The tensor current contracts with P^α , not p^α , because it interpolates the initial axial state. The factor $1/(m_Q + m_s)$ restores the same mass dimension as the axial-vector current.

3 Correlation Function

For a generic axial current $\Gamma_\mu \in \{\gamma_\mu \gamma_5, i\sigma_{\mu\alpha} P^\alpha \gamma_5 / (m_Q + m_s)\}$ we study

$$\Pi_\mu(p, q) = i \int d^4x e^{ip \cdot x} \langle \gamma(q, \varepsilon) | T \{ \bar{s}(x) \Gamma_\mu Q(x) \bar{Q}(0) i\gamma_5 s(0) \} | 0 \rangle. \quad (10)$$

The target Lorentz structure for the $1^+ \rightarrow 0^- \gamma$ transition is

$$(\eta \cdot \varepsilon^*)(p \cdot q) - (\eta \cdot q)(p \cdot \varepsilon^*). \quad (11)$$

On the correlator side we therefore project onto an equivalent transverse structure built from the open axial index and the photon polarization. With open indices this structure is represented by

$$T_{\mu\nu} = (p \cdot q)g_{\mu\nu} - q_\mu p_\nu, \quad S_{\mu\nu} \equiv -T_{\mu\nu} = p_\nu q_\mu - (p \cdot q)g_{\mu\nu}. \quad (12)$$

For a real photon, $q^2 = 0$, it obeys

$$q^\nu S_{\mu\nu} = 0, \quad (13)$$

so the projected invariant amplitude is gauge invariant under $\varepsilon_\nu \rightarrow \varepsilon_\nu + \lambda q_\nu$. In scalar form the same statement is

$$(\eta \cdot q)(p \cdot q) - (\eta \cdot q)(p \cdot q) = 0 \quad (14)$$

after replacing $\varepsilon \rightarrow q$.

4 Wick Contraction and OPE Bookkeeping

The OPE is separated into a hard-photon sector and a soft-photon sector:

$$\Pi_\mu^{\text{OPE}} = \Pi_\mu^{\text{hard}} + \Pi_\mu^{\text{soft}}. \quad (15)$$

This split is important because the two terms describe different physics and must not be double counted.

4.1 Hard photon emission

In the hard sector the photon is emitted perturbatively from either the heavy quark or the strange quark. After Wick contraction one obtains

$$\Pi_\mu^{\text{hard}} = iN_c \int d^4x e^{ip \cdot x} \text{Tr}_D \left[\Gamma_\mu S_Q^\gamma(x) i\gamma_5 S_s^0(-x) + \Gamma_\mu S_Q^0(x) i\gamma_5 S_s^\gamma(-x) \right], \quad (16)$$

where $N_c = 3$ and Tr_D denotes the Dirac trace. The free massive propagator is

$$S_q^0(x) = \int \frac{d^4k}{(2\pi)^4} e^{-ik \cdot x} \frac{i(\not{k} + m_q)}{k^2 - m_q^2}. \quad (17)$$

We keep explicit powers of m_s , including possible m_s^2 terms.

The first-order electromagnetic background-field correction for a generic quark flavor q is

$$S_q^\gamma(x) = -ie_q \int \frac{d^4k}{(2\pi)^4} e^{-ik \cdot x} \int_0^1 du \left[\frac{\not{k} + m_q}{2(m_q^2 - k^2)^2} F^{\alpha\beta}(ux) \sigma_{\alpha\beta} + \frac{ux_\alpha}{m_q^2 - k^2} F^{\alpha\beta}(ux) \gamma_\beta \right]. \quad (18)$$

In Eq. (16) this generic formula is used twice:

$$S_Q^\gamma(x) = S_q^\gamma(x)|_{q \rightarrow Q, m_q \rightarrow m_Q, e_q \rightarrow e_Q}, \quad (19)$$

$$S_s^\gamma(x) = S_q^\gamma(x)|_{q \rightarrow s, m_q \rightarrow m_s, e_q \rightarrow e_s}. \quad (20)$$

Thus the hard sector contains both heavy-line and strange-line electromagnetic background-field insertions. For a real photon we write

$$F^{\alpha\beta}(ux) = i(\varepsilon^\alpha q^\beta - \varepsilon^\beta q^\alpha) e^{iuq \cdot x}, \quad (21)$$

with the overall phase kept consistent with the Fourier convention.

Equation (18) is not a vacuum-condensate correction. It is the quark propagator in an external electromagnetic field and is included in the base analysis. What we do not include in the base hard sector are the ordinary SVZ-style condensate-expanded propagator terms such as $\langle \bar{s}s \rangle$, $\langle \bar{s}g_s \sigma G s \rangle$, or $\langle G^2 \rangle$ propagator corrections.

4.2 Equivalent momentum-space vertex form

For FeynCalc trace checks it is often simpler to use the equivalent explicit photon-vertex representation. With photon index ν ,

$$N_{Q,\mu\nu} = \text{Tr}_D[\Gamma_\mu(\not{k} + \not{q} + m_Q)\gamma_\nu(\not{k} + m_Q)i\gamma_5(\not{k} - \not{p} + m_s)], \quad (22)$$

$$N_{s,\mu\nu} = \text{Tr}_D[\Gamma_\mu(\not{k} + m_Q)i\gamma_5(\not{k} - \not{p} + m_s)\gamma_\nu(\not{k} - \not{p} + \not{q} + m_s)]. \quad (23)$$

These numerators are accompanied by the denominators

$$D_Q = [(k+q)^2 - m_Q^2][k^2 - m_Q^2][(k-p)^2 - m_s^2], \quad (24)$$

$$D_s = [k^2 - m_Q^2][(k-p)^2 - m_s^2][(k-p+q)^2 - m_s^2]. \quad (25)$$

The explicit-vertex form and the background-field form represent the same hard emission physics. We must not add both as independent contributions.

4.3 Soft photon emission: two-particle photon DAs

In the soft sector the photon is represented by its light-cone distribution amplitudes. Contracting only the heavy quark gives

$$\Pi_\mu^{\text{soft},2p} = i \int d^4x e^{ip \cdot x} [\Gamma_\mu S_Q^0(x) i\gamma_5]_{\alpha\beta} \langle \gamma(q, \varepsilon) | \bar{s}_\alpha(x) s_\beta(0) | 0 \rangle. \quad (26)$$

Fierz rearrangement converts the open spinor indices to a basis of Dirac bilinears,

$$\bar{s}_\alpha(x) s_\beta(0) = -\frac{1}{4} \sum_i (\Gamma_i)_{\beta\alpha} \bar{s}(x) \Gamma_i s(0), \quad (27)$$

so that

$$\Pi_\mu^{\text{soft},2p} = -\frac{i}{4} \int d^4x e^{ip \cdot x} \sum_i \text{Tr}_D[\Gamma_\mu S_Q^0(x) i\gamma_5 \Gamma_i] \langle \gamma(q, \varepsilon) | \bar{s}(x) \Gamma_i s(0) | 0 \rangle. \quad (28)$$

The photon matrix elements are then expressed in terms of two-particle photon DAs. Parameters such as $\chi \langle \bar{s}s \rangle$ enter here as part of the photon wavefunction normalization; they are not condensate corrections to the hard propagator.

4.4 Soft photon emission: three-particle photon DAs

To include three-particle photon DAs we must also allow one background gluon to come from the heavy-quark propagator,

$$S_Q^G(x) = -ig_s \int \frac{d^4k}{(2\pi)^4} e^{-ik \cdot x} \int_0^1 du \left[\frac{\not{k} + m_Q}{2(m_Q^2 - k^2)^2} G^{\alpha\beta}(ux) \sigma_{\alpha\beta} + \frac{ux_\alpha}{m_Q^2 - k^2} G^{\alpha\beta}(ux) \gamma_\beta \right]. \quad (29)$$

This term is not a gluon-condensate correction. It is an operator insertion that combines with photon matrix elements of the form

$$\langle \gamma(q, \varepsilon) | \bar{s}(x) \Gamma_i g_s G_{\alpha\beta}(ux) s(0) | 0 \rangle, \quad (30)$$

which define the three-particle photon DAs.

5 Staged Calculation Plan

We proceed in two stages.

Stage 1. Include the hard photon emission terms and the two-particle photon DAs:

$$\Pi_\mu^{\text{Stage 1}} = \Pi_\mu^{\text{hard}} + \Pi_\mu^{\text{soft},2p}. \quad (31)$$

This stage keeps all explicit m_s terms and omits ordinary condensate-expanded propagators.

Stage 2. Add the three-particle photon DA contributions generated by S_Q^G and the corresponding light-cone matrix elements:

$$\Pi_\mu^{\text{Stage 2}} = \Pi_\mu^{\text{Stage 1}} + \Pi_\mu^{\text{soft},3p}. \quad (32)$$

This staging keeps the derivation auditable: first we verify the core hard and two-particle DA structures, then we add the more involved background-gluon terms.

6 Current FeynCalc Artifacts

The script `scripts/step2_current_building_blocks.wl` checks the basic Dirac conventions and defines the current vertices. The script `scripts/step2_hard_photon_numerators.wl` computes numerator-level hard-photon traces in the explicit photon-vertex representation. These traces are stored in `outputs/step2_hard_photon_numerators.txt`.

The script `scripts/step2_e1_project_hard_numerators.wl` applies the E1 projector

$$\mathcal{P}_{E1}^{\mu\nu} = \frac{p^\nu q^\mu - (p \cdot q)g^{\mu\nu}}{2(p \cdot q)^2}, \quad (33)$$

which satisfies $\mathcal{P}_{E1}^{\mu\nu}[p_\nu q_\mu - (p \cdot q)g_{\mu\nu}] = 1$ for $q^2 = 0$. The projected hard numerators are stored in `outputs/step2_e1_projected_hard_numerators.txt`.

The companion script `scripts/ward_transversality_check.wl` verifies the Ward/transversality properties of the same E1 tensor with FeynCalc. Its output is stored in `outputs/ward_transversality_check`.

The script `scripts/step2_soft_2p_fierz_traces.wl` computes the trace factors multiplying the two-particle photon DA matrix elements after the Fierz rearrangement. The output `outputs/step2_soft_2p_fierz_traces.txt` shows which bilinears feed the axial current and which feed the tensor current.

7 Stage 1 Axial-Current Numerical Baseline

Before deriving the tensor-current sum rule, we implemented the known single-axial-current result for $D_{s1}(2460) \rightarrow D_s \gamma$ as a benchmark. This is the coupling called g_1 by Colangelo, De Fazio, and Ozpineci. In our notation this is the axial-current form factor G_A .

The Stage 1 version contains

$$G_A^{\text{Stage 1}} = G_A^{\text{pert}} + G_A^{\chi\phi\gamma} + G_A^{\mathbb{A},\bar{H}\gamma} + G_A^{f_{3\gamma}\Psi^v}, \quad (34)$$

where the three-particle DA integral usually denoted by $\mathcal{F}_1$ is omitted until Stage 2.

The numerical script is

`scripts/stage1_axial_g1_baseline.py`.

It writes the full Borel/threshold grid to

`outputs/stage1_axial_g1_baseline.csv`.

7.1 Magnetic-susceptibility sign convention

The Colangelo paper prints $\chi = -(3.15 \pm 0.30) \text{ GeV}^{-2}$ in its matrix-element convention. In this project we store the BBK magnitude as

$$\chi(1 \text{ GeV}) = +3.15 \pm 0.30 \text{ GeV}^{-2}. \quad (35)$$

Therefore the sign of the $\chi\phi_\gamma$ term in the implemented sum rule is flipped relative to the literal printed formula. This is a convention translation, not a physical change. A diagnostic run that inserts positive χ into the printed negative- χ formula gives a much larger width and is kept only as a warning check.

7.2 Preliminary Stage 1 result

Using the project inputs, $u_0 = 1/2$, $3 \leq M^2 \leq 6 \text{ GeV}^2$, and $s_0 = (2.50, 2.55, 2.60)^2 \text{ GeV}^2$, the axial-current Stage 1 grid gives

$$-0.399 \leq G_A^{\text{Stage } 1} \leq -0.314 \text{ GeV}^{-1}. \quad (36)$$

With

$$\Gamma = \frac{\alpha_{\text{em}}}{3} G^2 |\vec{q}_\gamma|^3, \quad (37)$$

this corresponds to

$$20.8 \text{ keV} \leq \Gamma_A^{\text{Stage } 1} [D_{s1}(2460) \rightarrow D_s \gamma] \leq 33.5 \text{ keV}. \quad (38)$$

This is not yet the physical mixed-state result. It is the axial-current baseline that will be combined with the tensor-current form factor after the J_B sum rule is derived.

8 First Tensor-Current Step: Leading-Twist Soft Contribution

The tensor-current calculation starts from the trace-level identity produced by `scripts/step3_soft_2p_e1_pro`. For the two-particle tensor photon DA matrix element, the E1 projections are

$$\mathcal{P}_{E1} [\text{Tr}(J_A S_Q i\gamma_5 \sigma_{\rho\lambda})(\varepsilon^\rho q^\lambda - \varepsilon^\lambda q^\rho)] = 8 \frac{k \cdot q}{p \cdot q}, \quad (39)$$

$$\mathcal{P}_{E1} [\text{Tr}(J_B S_Q i\gamma_5 \sigma_{\rho\lambda})(\varepsilon^\rho q^\lambda - \varepsilon^\lambda q^\rho)] = 8 \frac{m_Q}{m_Q + m_s}. \quad (40)$$

After the two-particle DA Fourier integral and double Borel transform, $k = p + \bar{u}q$, hence $k \cdot q = p \cdot q$ for $q^2 = 0$. Therefore, before the hadronic decay-constant normalization, the leading-twist tensor-current OPE piece is

$$\Pi_B^{\chi\phi_\gamma} = \frac{m_Q}{m_Q + m_s} \Pi_A^{\chi\phi_\gamma}. \quad (41)$$

The exploratory script `scripts/stage1_tensor_gb_leading_twist.py` uses this relation to estimate only the $\chi\phi_\gamma$ part of G_B . This is not the final tensor-current sum rule. It omits the tensor-current hard contribution, the vector DA contribution, and the Stage 2 three-particle DA terms.

Using the project inputs, the partial leading-twist estimate gives

$$G_B^{\chi\phi_\gamma} \simeq +0.205 \text{ to } +0.273 \text{ GeV}^{-1}, \quad \text{if } f_B = f_T, \quad (42)$$

$$G_B^{\chi\phi_\gamma} \simeq +0.114 \text{ to } +0.151 \text{ GeV}^{-1}, \quad \text{if } f_B = f_T \frac{m_{D_{s1}}}{m_c + m_s}. \quad (43)$$

The sizeable sign difference relative to G_A means that the physical mixed amplitudes can be sensitive to the tensor-current normalization. This is why the next step is to finish the full G_B sum rule before quoting final mixed widths.

9 Decay-Constant Normalization Before Mixing

The labels f_A , f_B , and f_T must not be mixed casually. The axial current couples to the axial basis state through

$$\langle 0 | J_A^\mu | D_{s1}^A(P, \eta) \rangle = f_A m_A \eta^\mu, \quad (44)$$

whereas our tensor-derived current is

$$J_B^\mu = \frac{i}{m_c + m_s} \bar{s} \sigma^{\mu\alpha} P_\alpha \gamma_5 c. \quad (45)$$

The corresponding decay constant is defined by

$$\langle 0 | J_B^\mu | D_{s1}^B(P, \eta) \rangle = f_B m_B \eta^\mu. \quad (46)$$

Therefore $f_B = f_T$ is not a physical statement by itself. It is only true if the tabulated f_T is already defined for the same current J_B^μ and with the same mass normalization. If instead the input table gives the more common tensor-current matrix element

$$\langle 0 | \bar{s} \sigma_{\mu\nu} \gamma_5 c | D_{s1}^B(P, \eta) \rangle = f_T (\eta_\mu P_\nu - \eta_\nu P_\mu), \quad (47)$$

then contraction with $P^\nu / (m_c + m_s)$ gives the effective normalization

$$f_B^{\text{eff}} \simeq f_T \frac{m_{D_{s1}}}{m_c + m_s}, \quad (48)$$

up to the sign fixed by the precise convention for $\sigma_{\mu\nu} \gamma_5$ and by the transverse condition $\eta \cdot P = 0$.

Only after G_A and G_B have been computed with their own basis-current decay constants do we form the physical amplitudes:

$$G_{2460} = \sin \theta G_A + \cos \theta G_B, \quad (49)$$

$$G_{2536} = \cos \theta G_A - \sin \theta G_B. \quad (50)$$

Thus the sine–cosine factors belong to the physical-state mixing of amplitudes, not to an arbitrary identification $f_B = f_T$. In the numerical scripts, $f_B = f_T$ is kept only as a diagnostic. The converted $f_B = f_T m_{D_{s1}} / (m_c + m_s)$ option is the safer central normalization unless the input-table definition of f_T is changed.

10 Tensor-Current Hard Term: Triangle Reduction

The hard tensor-current numerator is more complicated than the axial one because the J_B current introduces an extra factor of P^α and therefore higher powers of the loop momentum after E1 projection. The script `scripts/step3_tensor_hard_triangle_reduction.wl` rewrites scalar products such as k^2 , $k \cdot p$, and $k \cdot q$ in terms of inverse propagator denominators. For the heavy-line photon emission we define

$$d_{H1} = (k + q)^2 - m_Q^2, \quad (51)$$

$$d_{H2} = k^2 - m_Q^2, \quad (52)$$

$$d_{H3} = (k - p)^2 - m_s^2, \quad (53)$$

and similarly for the strange-line photon emission. Setting $d_{Hi} = 0$ isolates the genuine three-denominator triangle core. For the tensor current this gives

$$N_{B,H}^{\text{core}} = \frac{-8im_Q^2 + 4im_Q m_s}{m_Q + m_s} - \frac{4im_Q^2 p^2}{(m_Q + m_s)(p \cdot q)}, \quad (54)$$

$$N_{B,s}^{\text{core}} = \frac{-4im_Q m_s}{m_Q + m_s} - \frac{4im_s^2 p^2}{(m_Q + m_s)(p \cdot q)}. \quad (55)$$

The same reduction was also performed for the axial current in `scripts/step3_axial_hard_triangle_reduction.wl`. This is a useful calibration because the axial hard term has a known published spectral density.

10.1 Why denominator-cancellation terms are separated

Terms proportional to one or more inverse denominators cancel one or more propagators. After such a cancellation the loop integral contains at most two propagators. These reduced integrals depend on at most one of the two external virtualities, p^2 or $(p+q)^2$, and therefore do not generate the double-pole structure selected by the double Borel transform. They are recorded in the output files for auditability, but the double-pole sum rule is built from the three-denominator triangle core.

This observation narrows the remaining hard-tensor task: derive the double spectral density associated with the tensor-current triangle core and match it to the same E1 invariant amplitude used for the axial-current benchmark.

11 First Hard-Tensor Spectral Density Candidate

The next step is to translate the triangle core into a perturbative spectral density. We do this by calibrating against the known axial-current result. For one photon-emitting line, let m_i be the emitting-quark mass and m_j the spectator mass. The axial triangle core has the schematic form

$$N_i^{A,\text{core}} = -4i \left[a_i + \frac{b_i}{p \cdot q} \right]. \quad (56)$$

Comparing with the published axial spectral density fixes the replacement

$$-4i a_i \longrightarrow 2a_i \log \frac{s - m_j^2 + m_i^2 - \lambda^{1/2}(s, m_j^2, m_i^2)}{s - m_j^2 + m_i^2 + \lambda^{1/2}(s, m_j^2, m_i^2)}, \quad (57)$$

$$-4i \frac{b_i}{p \cdot q} \longrightarrow \frac{b_i}{m_i^2} \frac{m_j^2 - m_i^2 - s}{s^2} \lambda^{1/2}(s, m_j^2, m_i^2). \quad (58)$$

The overall factor is $-3/(8\pi^2)$, and the total axial density is assembled as the strange-emission contribution minus the heavy-emission contribution, with the corresponding charges. The script

`scripts/stage1_tensor_gb_hard_candidate.py`

first applies these rules back to the axial current and reproduces Colangelo's $\rho^P(s)$ with a maximum numerical difference 1.4×10^{-17} on the test grid.

For the tensor current the core contains $p^2/(p \cdot q)$:

$$N_{B,H}^{\text{core}} = -4i \left[\frac{2m_Q^2 - m_Q m_s}{m_Q + m_s} + \frac{m_Q^2 p^2}{(m_Q + m_s)(p \cdot q)} \right], \quad (59)$$

$$N_{B,s}^{\text{core}} = -4i \left[\frac{m_Q m_s}{m_Q + m_s} + \frac{m_s^2 p^2}{(m_Q + m_s)(p \cdot q)} \right]. \quad (60)$$

The diagonal double-dispersion representation explains how the p^2 numerator is treated. For the double-pole part one has

$$\frac{p^2}{(s - p^2)(s - P^2)} = \frac{s}{(s - p^2)(s - P^2)} - \frac{1}{s - P^2}, \quad P = p + q. \quad (61)$$

The second term has only a single pole and is removed by the double Borel projection in p^2 and P^2 . Therefore, inside the double-pole sum rule, p^2 in the hard numerator can be replaced by the spectral variable s . The same argument gives $P^2 \rightarrow s$ for any numerator factor of P^2 . The script

`scripts/step4_double_pole_p2_reduction.py`

records this identity as a short audit check.

Combining this hard candidate with the tensor-DA two-particle soft contribution gives, over the same M^2 and s_0 windows,

$$G_B^{\text{hard}} \simeq -0.831 \text{ to } -0.673 \text{ GeV}^{-1}, \quad G_B^{\text{Stage } 1} \simeq -0.510 \text{ to } -0.440 \text{ GeV}^{-1}, \quad (f_B = f_T), \quad (62)$$

$$G_B^{\text{hard}} \simeq -0.460 \text{ to } -0.373 \text{ GeV}^{-1}, \quad G_B^{\text{Stage } 1} \simeq -0.283 \text{ to } -0.244 \text{ GeV}^{-1}, \quad \left(f_B = f_T \frac{m_{D_{s1}}}{m_c + m_s} \right). \quad (63)$$

At $\theta = 35.3^\circ$, the corresponding candidate mixed widths are

$$\Gamma[D_{s1}(2460) \rightarrow D_s \gamma] \simeq 62.5 \text{ to } 87.9 \text{ keV}, \quad \Gamma[D_{s1}(2536) \rightarrow D_s \gamma] \simeq 0.00 \text{ to } 0.30 \text{ keV}, \quad (f_B = f_T), \quad (64)$$

$$\Gamma[D_{s1}(2460) \rightarrow D_s \gamma] \simeq 30.9 \text{ to } 44.7 \text{ keV}, \quad \Gamma[D_{s1}(2536) \rightarrow D_s \gamma] \simeq 4.0 \text{ to } 8.2 \text{ keV}, \quad \left(f_B = f_T \frac{m_{D_{s1}}}{m_c + m_s} \right). \quad (65)$$

These numbers should be read as a controlled Stage 1 candidate, not yet as the final result, because Stage 2 three-particle photon DAs remain. The $p^2 \rightarrow s$ step is justified for the double-pole part; any single-pole remnants belong outside the double-Borel sum rule.

12 Stage 2 Axial Three-Particle DA Check

Before deriving the tensor-current three-particle terms, we implemented the known axial-current correction as a Stage 2 calibration. The published axial sum rule contains the combination

$$\mathcal{F}_1 = \mathcal{S} + \tilde{\mathcal{S}} - \mathcal{T}_1 - \mathcal{T}_2 + \mathcal{T}_3 + \mathcal{T}_4 + 2v(-\mathcal{S} - \mathcal{T}_3 + \mathcal{T}_2), \quad (66)$$

integrated over the two standard regions

$$0 \leq v \leq 1 - u_0, \quad 0 \leq \alpha_g \leq \frac{u_0}{1 - v}, \quad (67)$$

$$1 - u_0 \leq v \leq 1, \quad 0 \leq \alpha_g \leq \frac{1 - u_0}{v}, \quad (68)$$

with

$$\alpha_q = u_0 - (1 - v)\alpha_g, \quad \alpha_{\bar{q}} = 1 - u_0 - v\alpha_g. \quad (69)$$

The script

`scripts/stage2_axial_g1_three_particle.py`

evaluates this integral using the three-particle photon DAs in `photon_da.py`. For $u_0 = 1/2$ it finds

$$\int \mathcal{F}_1 = +7.033 \times 10^{-2}, \quad (70)$$

$$\Delta G_A^{\mathcal{F}_1} \simeq +0.0012 \text{ to } +0.0020 \text{ GeV}^{-1}. \quad (71)$$

Thus the axial Stage 2 result over the same Borel and threshold window is

$$-0.398 \leq G_A^{\text{Stage } 2} \leq -0.312 \quad \text{GeV}^{-1}, \quad (72)$$

corresponding to

$$20.5 \text{ keV} \leq \Gamma_A^{\text{Stage } 2}[D_{s1}(2460) \rightarrow D_s \gamma] \leq 33.3 \text{ keV}. \quad (73)$$

This small correction is a useful calibration, but it does not yet complete Stage 2 for the mixed physical amplitudes. The remaining Stage 2 task is to derive the tensor-current analog of the three-particle photon DA contribution from the background-gluon insertion and project it onto the same E1 invariant.

13 First Tensor-Current Stage 2 Trace Pass

For the tensor-current three-particle term we first separated the local $\sigma_{\alpha\beta}$ part of the heavy-quark background-gluon propagator from the explicitly $x_\alpha\gamma_\beta$ part. The script

`scripts/step5_soft_3p_trace_kernels.wl`

computes the E1-projected traces for the local piece against the $\mathcal{S}$, $\tilde{\mathcal{S}}$, $\mathcal{T}_1$, and $\mathcal{T}_2$ Lorentz structures. The nonzero local kernels are

$$K_A[\mathcal{S}] = 8 \frac{k \cdot q}{p \cdot q}, \quad K_B[\mathcal{S}] = 8 \frac{m_Q}{m_Q + m_s}, \quad (74)$$

$$K_A[\mathcal{T}_1] = -32i \frac{k \cdot q}{p \cdot q}, \quad K_B[\mathcal{T}_1] = -32i \frac{m_Q}{m_Q + m_s}, \quad (75)$$

$$K_A[\mathcal{T}_2] = +32i \frac{k \cdot q}{p \cdot q}, \quad K_B[\mathcal{T}_2] = +32i \frac{m_Q}{m_Q + m_s}. \quad (76)$$

The $\tilde{\mathcal{S}}$ kernel vanishes for this local part. After the three-particle light-cone Fourier integral and double Borel projection, $k \cdot q = p \cdot q$, so the local trace ratio is again

$$\frac{K_B}{K_A} = \frac{m_Q}{m_Q + m_s}. \quad (77)$$

This is the same ratio encountered in the two-particle tensor-DA family.

At this point it is useful to separate the published axial combination into the piece generated by the $\sigma_{\alpha\beta}$ part of the background-field propagator and the piece generated by the $x_\alpha G^{\alpha\beta} \gamma_\beta$ part:

$$\mathcal{F}_1^\sigma = \mathcal{S} + \tilde{\mathcal{S}} - \mathcal{T}_1 - \mathcal{T}_2 + \mathcal{T}_3 + \mathcal{T}_4, \quad (78)$$

$$\mathcal{F}_1^{x\gamma} = 2v(-\mathcal{S} - \mathcal{T}_3 + \mathcal{T}_2), \quad \mathcal{F}_1 = \mathcal{F}_1^\sigma + \mathcal{F}_1^{x\gamma}. \quad (79)$$

The numerical split, evaluated in

`scripts/step10_stage2_f1_split.py`,

is

$$\int \mathcal{F}_1^\sigma = -1.5625 \times 10^{-1}, \quad (80)$$

$$\int \mathcal{F}_1^{x\gamma} = +2.2658 \times 10^{-1}, \quad (81)$$

$$\int \mathcal{F}_1 = +7.0330 \times 10^{-2}. \quad (82)$$

Thus the small published axial Stage 2 correction is partly the result of a cancellation between two larger pieces. The simple trace ratio $m_Q/(m_Q + m_s)$ is established for the $\sigma_{\alpha\beta}$ part, so the controlled sigma-matched tensor-current correction is

$$\Delta G_B^{3p} \simeq -0.00366 \text{ to } -0.00210 \text{ GeV}^{-1}, \quad (f_B = f_T \text{ diagnostic}), \quad (83)$$

$$\Delta G_B^{3p} \simeq -0.00203 \text{ to } -0.00116 \text{ GeV}^{-1}, \quad \left(f_B = f_T \frac{m_{D_{s1}}}{m_c + m_s} \text{ central} \right). \quad (84)$$

The corresponding sigma-matched Stage 2 mixed-width estimate at $\theta = 35.3^\circ$ is

$$\Gamma[D_{s1}(2460) \rightarrow D_s \gamma] \simeq 62.9 \text{ to } 88.2 \text{ keV}, \quad \Gamma[D_{s1}(2536) \rightarrow D_s \gamma] \simeq 0.00 \text{ to } 0.26 \text{ keV}, \quad (f_B = f_T \text{ diagnostic}), \quad (85)$$

$$\Gamma[D_{s1}(2460) \rightarrow D_s \gamma] \simeq 30.9 \text{ to } 44.8 \text{ keV}, \quad \Gamma[D_{s1}(2536) \rightarrow D_s \gamma] \simeq 3.8 \text{ to } 8.0 \text{ keV}, \quad \left(f_B = f_T \frac{m_{D_{s1}}}{m_c + m_s} \text{ central} \right) \quad (86)$$

If, instead, one multiplies the full axial $\mathcal{F}_1$ integral by the same trace ratio, one obtains only a diagnostic estimate:

$$\Delta G_B^{3p} \simeq +0.00094 \text{ to } +0.00165 \text{ GeV}^{-1}, \quad (f_B = f_T \text{ diagnostic}), \quad (87)$$

$$\Delta G_B^{3p} \simeq +0.00052 \text{ to } +0.00091 \text{ GeV}^{-1}, \quad \left(f_B = f_T \frac{m_{D_{s1}}}{m_c + m_s} \text{ central}\right), \quad (88)$$

with central-normalization widths

$$\Gamma[D_{s1}(2460) \rightarrow D_s \gamma] \simeq 30.5 \text{ to } 44.5 \text{ keV}, \quad (89)$$

$$\Gamma[D_{s1}(2536) \rightarrow D_s \gamma] \simeq 4.0 \text{ to } 8.1 \text{ keV}. \quad (90)$$

The difference between the sigma-matched and full-ratio diagnostic numbers is small at the current precision, but conceptually important: the $x_\alpha \gamma_\beta$ part must be matched separately before the tensor Stage 2 correction is final.

14 Stage 2 Tensor: Explicit x -Dependent Kernels

The next audit step keeps the coordinate vector x explicit. For the $\sigma_{\alpha\beta}$ part of S_Q^G , the raw E1 projections of the $\mathcal{T}_3$ and $\mathcal{T}_4$ structures are

$$K_A[\mathcal{T}_3] = +16i \frac{k \cdot q}{p \cdot q}, \quad K_B[\mathcal{T}_3] = +16i \frac{m_Q}{m_Q + m_s}, \quad (91)$$

$$K_A[\mathcal{T}_4] = -16i \frac{k \cdot q}{p \cdot q}, \quad K_B[\mathcal{T}_4] = -16i \frac{m_Q}{m_Q + m_s}. \quad (92)$$

Thus these two structures also obey the same ratio $K_B/K_A = m_Q/(m_Q + m_s)$ after $k \cdot q = p \cdot q$. This check is recorded in

`scripts/step6_soft_3p_x_dependent_kernels.wl.`

The genuinely new complication comes from the second term in the background propagator,

$$\frac{u x_\alpha}{m_Q^2 - k^2} G^{\alpha\beta}(ux) \gamma_\beta. \quad (93)$$

The script

`scripts/step7_soft_3p_xgamma_kernels.wl`

computes the raw kernels with x kept explicit. Unlike the previous kernels, the tensor-to-axial ratio is not a constant at this stage. For example the $\mathcal{S}$ ratio after only $k \cdot q \rightarrow p \cdot q$ contains

$$\frac{(p \cdot q)(k \cdot x - p \cdot x) + (k \cdot p + p^2 + 2p \cdot q)(q \cdot x)}{2m_Q^2(q \cdot x)}, \quad (94)$$

and analogous expressions occur for $\mathcal{T}_1$, $\mathcal{T}_2$, and $\mathcal{T}_3$. Therefore the tensor Stage 2 three-particle result cannot yet be promoted from an estimate to a final term. The remaining calculation is to perform the Fourier/Borel replacement of the explicit $x_\alpha/(q \cdot x)$ factors, including the derivative action on the heavy-quark denominator, and then recombine the result with the photon DA structures.

As a diagnostic, we can impose only the light-cone momentum routing

$$k = p + aq, \quad a = \alpha_{\bar{q}} + v\alpha_g, \quad q^2 = 0, \quad (95)$$

without yet applying the derivative replacement. The script

`scripts/step8_xgamma_saddle_reduction.py`

reduces the raw ratios $K_B/[(m_Q/(m_Q + m_s))K_A]$ to

$$R_S = \frac{p^2 + (a+1)p \cdot q}{m_Q^2}, \quad (96)$$

$$R_{\mathcal{T}_1} = \frac{p^2 + (3a+2)p \cdot q}{3m_Q^2}, \quad (97)$$

$$R_{\mathcal{T}_2} = \frac{p^2 + (2-a)p \cdot q}{m_Q^2}, \quad (98)$$

$$R_{\mathcal{T}_3} = \frac{p^2 + 2a p \cdot q}{m_Q^2}. \quad (99)$$

If one additionally uses the heavy-line on-shell relation $m_Q^2 = k^2 = p^2 + 2a p \cdot q$, only $R_{\mathcal{T}_3}$ becomes exactly one. The other ratios remain nontrivial. This confirms that the $x_\alpha \gamma_\beta$ term must be treated with the full Fourier/Borel derivative machinery. It is not controlled by the simple trace-ratio argument used for the $\sigma_{\alpha\beta}$ part.

15 Stage 2 Tensor: Derivative Reduction of $x_\alpha \gamma_\beta$

The next step applies the Fourier derivative rules to the raw $x_\alpha \gamma_\beta$ kernels. With phase

$$\exp[i(p + aq - k) \cdot x], \quad (100)$$

integration over x gives, up to a common phase convention,

$$p \cdot x \rightarrow -i p \cdot \partial_k, \quad (101)$$

$$q \cdot x \rightarrow -i q \cdot \partial_k, \quad (102)$$

$$k \cdot x F(k) \rightarrow -i \partial_{k_\mu} [k_\mu F(k)] = -i [4F(k) + k \cdot \partial_k F(k)]. \quad (103)$$

The script

`scripts/step9_xgamma_derivative_reduction.wl`

applies these rules to the scalar kernels multiplying $1/(m_Q^2 - k^2)$. After the light-cone routing $k = p + aq$, the reduced axial kernels are proportional to

$$A_S^{x\gamma} = \frac{-8m_Q}{[-m_Q^2 + p^2 + 2a p \cdot q]^2}, \quad (104)$$

$$A_{\mathcal{T}_1}^{x\gamma} = \frac{48im_Q}{[-m_Q^2 + p^2 + 2a p \cdot q]^2}, \quad (105)$$

$$A_{\mathcal{T}_2}^{x\gamma} = \frac{16im_Q}{[-m_Q^2 + p^2 + 2a p \cdot q]^2}, \quad (106)$$

$$A_{\mathcal{T}_3}^{x\gamma} = \frac{-16im_Q}{[-m_Q^2 + p^2 + 2a p \cdot q]^2}. \quad (107)$$

The corresponding tensor-current kernels are not obtained by a universal $m_Q/(m_Q + m_s)$ multiplication. The ratios after derivative reduction are

$$R_S^\partial = 1 + \frac{(1-a)p \cdot q}{m_Q^2}, \quad (108)$$

$$R_{\mathcal{T}_1}^\partial = \frac{2m_Q^2 + p^2 + (2-a)p \cdot q}{3m_Q^2}, \quad (109)$$

$$R_{\mathcal{T}_2}^\partial = -2 + \frac{3p^2 + (2+3a)p \cdot q}{m_Q^2}, \quad (110)$$

$$R_{\mathcal{T}_3}^\partial = \frac{2m_Q^2 - p^2 - 2a p \cdot q}{m_Q^2}. \quad (111)$$

All these reduced kernels carry the squared heavy-quark denominator. For the double-pole residue we drop numerator terms proportional to $D = -m_Q^2 + p^2 + 2a p \cdot q$, equivalently using $m_Q^2 = p^2 + 2a p \cdot q$ in the residue. In the three-particle integration domains,

$$a = \alpha_{\bar{q}} + v\alpha_g = 1 - u_0, \quad (112)$$

so for the symmetric Borel point $u_0 = 1/2$ this residue is evaluated at $a = 1/2$.

The script

`scripts/step11_tensor_xgamma_ratio_integral.py`

uses these residue ratios to match the $\mathcal{S}$, $\mathcal{T}_2$, and $\mathcal{T}_3$ pieces of the tensor-current $x_\alpha \gamma_\beta$ contribution to the known axial normalization

$$\mathcal{F}_1^{x\gamma} = 2v(-\mathcal{S} - \mathcal{T}_3 + \mathcal{T}_2). \quad (113)$$

With the heavy-line residue condition and physical $p \cdot q$ in the E1 invariant, the matched $\mathcal{S}$, $\mathcal{T}_2$, $\mathcal{T}_3$ integral is

$$\int \mathcal{F}_{1,B}^{x\gamma}[\mathcal{S}, \mathcal{T}_2, \mathcal{T}_3] = +2.5818 \times 10^{-1}. \quad (114)$$

The tensor-current derivative reduction also produces a $\mathcal{T}_4$ residue even though the axial E1 kernel has $A_{\mathcal{T}_4}^{x\gamma} = 0$. This is not an obstacle: its normalization can be fixed directly relative to the axial $\mathcal{T}_2$ unit. On the double-pole residue,

$$\frac{B_{\mathcal{T}_4}^{x\gamma}}{[m_Q/(m_Q + m_s)] A_{\mathcal{T}_2}^{x\gamma}} = -\frac{(1-a)p \cdot q}{m_Q^2}. \quad (115)$$

This gives the tensor-only contribution

$$\int \mathcal{F}_{1,B}^{x\gamma}[\mathcal{T}_4] = -4.2136 \times 10^{-2}, \quad (116)$$

and therefore

$$\int \mathcal{F}_{1,B}^{x\gamma} = +2.1605 \times 10^{-1}. \quad (117)$$

Together with the sigma-like part this gives the full matched tensor three-particle integral

$$\int (\mathcal{F}_{1,B}^\sigma + \mathcal{F}_{1,B}^{x\gamma}) = +5.9796 \times 10^{-2}. \quad (118)$$

For comparison, the axial total is $\int \mathcal{F}_1 = +7.0330 \times 10^{-2}$. A diagnostic calculation that instead inserts the physical hadron value $p^2 = m_{D_s}^2$ directly into the OPE residue ratios gives a much larger integral, +1.5072, so we do not use it as the working tensor estimate.

Numerically, for the central decay-constant normalization $f_B = f_T m_{D_{s1}} / (m_c + m_s)$, the full matched Stage 2 tensor-current result gives

$$\Delta G_B^{3p} = +0.00044 \text{ to } +0.00078 \text{ GeV}^{-1}, \quad (119)$$

$$G_B^{\text{Stage 2}} = -0.2821 \text{ to } -0.2436 \text{ GeV}^{-1}. \quad (120)$$

At $\theta = 35.3^\circ$ this corresponds to

$$\Gamma[D_{s1}(2460) \rightarrow D_s \gamma] = 30.6 \text{ to } 44.5 \text{ keV}, \quad (121)$$

$$\Gamma[D_{s1}(2536) \rightarrow D_s \gamma] = 4.0 \text{ to } 8.1 \text{ keV}. \quad (122)$$

The final matched result is close to the older full-ratio diagnostic, but the derivation is now explicit about which pieces come from the $\sigma_{\alpha\beta}$ part, which come from $x_\alpha G^{\alpha\beta} \gamma_\beta$, and how the tensor-only $\mathcal{T}_4$ term is normalized.

16 Final Stage 2 Uncertainty Scan

The script

`scripts/final_stage2_uncertainty_scan.py`

performs a first independent-input uncertainty scan around the completed matched Stage 2 baseline. Each random point samples the Borel parameter $M^2 \in [3, 6] \text{ GeV}^2$, the continuum threshold $\sqrt{s_0} \in [2.50, 2.60] \text{ GeV}$, and the numerical inputs listed in `inputs_table.csv`. The tensor-current decay constant is converted as

$$f_B = f_T \frac{m_{D_{s1}}}{m_c + m_s}. \quad (123)$$

The scan is not yet a correlated global error analysis; it is a transparent Monte Carlo error budget for the present sum rule. It is the preferred band to quote because it allows all inputs, M^2 , and s_0 to fluctuate together.

For fixed $\theta = 35.3^\circ$, using 500 random points, the scan gives

$$G[D_{s1}(2460) \rightarrow D_s \gamma] = -0.424_{-0.067}^{+0.065} \text{ GeV}^{-1}, \quad (124)$$

$$G[D_{s1}(2536) \rightarrow D_s \gamma] = -0.140_{-0.051}^{+0.047} \text{ GeV}^{-1}, \quad (125)$$

where the uncertainty indicates the 16–84 percentile interval. The corresponding width estimates are

$$\Gamma[D_{s1}(2460) \rightarrow D_s \gamma] = 37.9_{-10.7}^{+12.9} \text{ keV}, \quad (126)$$

$$\Gamma[D_{s1}(2536) \rightarrow D_s \gamma] = 6.1_{-3.4}^{+5.2} \text{ keV}. \quad (127)$$

If instead the mixing angle is sampled uniformly over $25^\circ \leq \theta \leq 45^\circ$, the same scan gives

$$\Gamma[D_{s1}(2460) \rightarrow D_s \gamma] = 36.9_{-10.8}^{+12.6} \text{ keV}, \quad (128)$$

$$\Gamma[D_{s1}(2536) \rightarrow D_s \gamma] = 6.0_{-4.1}^{+7.6} \text{ keV}. \quad (129)$$

The full scan table is written to

`outputs/final_stage2_uncertainty_scan.csv`.

For visualization the script

`scripts/uncertainty_publication_plots.py`

also fits Gaussian curves to the Monte Carlo width histograms using the sample mean and sample standard deviation. These fits are descriptive rather than the primary quoted uncertainties, because the width distribution is mildly right-tailed, especially for $D_{s1}(2536)$. For the fixed-angle ensemble the Gaussian summaries are

$$\Gamma_{2460} = 38.99 \pm 12.76 \text{ keV}, \quad (130)$$

$$\Gamma_{2536} = 7.39 \pm 5.67 \text{ keV}, \quad (131)$$

while for $25^\circ \leq \theta \leq 45^\circ$ they are

$$\Gamma_{2460} = 38.47 \pm 12.78 \text{ keV}, \quad (132)$$

$$\Gamma_{2536} = 7.79 \pm 6.76 \text{ keV}. \quad (133)$$

The corresponding publication histograms are shown in Figs. 1 and 2.




Figure 1: Monte Carlo width distributions for fixed $\theta = 35.3^\circ$. The black curves are Gaussian overlays using the sample mean and standard deviation; the dashed vertical lines mark the 16–84 percentile interval.




Figure 2: Monte Carlo width distributions when the mixing angle is sampled uniformly over $25^\circ \leq \theta \leq 45^\circ$. The black curves are Gaussian overlays using the sample mean and standard deviation; the dashed vertical lines mark the 16–84 percentile interval.

The diagnostic script

scripts/uncertainty_budget_one_at_time.py

then varies one input group at a time around the central point. This is not the band to quote as the final uncertainty; it explains which input groups drive the Monte Carlo spread.

Varied group	Γ_{2460} (keV)	Γ_{2536} (keV)
$M^2, \sqrt{s_0}$	37.4 [33.0, 41.1]	5.95 [5.05, 7.00]
θ	37.3 [33.7, 39.6]	5.83 [2.40, 11.2]
quark and hadron masses	37.1 [35.1, 39.1]	6.01 [4.92, 7.27]
decay constants	37.7 [32.9, 42.4]	5.91 [3.48, 9.42]
condensates and χ	38.7 [29.2, 48.6]	6.40 [4.00, 9.11]
photon DA shape	37.2 [37.2, 37.2]	6.04 [6.04, 6.04]
all inputs, fixed θ	37.4 [26.8, 50.2]	5.50 [2.73, 11.2]
all inputs and θ	36.1 [25.7, 48.3]	6.28 [1.78, 14.9]

Table 1: One-at-a-time 16–84 percentile uncertainty budget with 500 random points per group. The photon-shape row is flat in the present matched reduction; it should be read as negligible sensitivity in this implementation, not as a general statement about photon DAs.

The main lesson from Table 1 is that the $D_{s1}(2460)$ width is most sensitive to condensate/ χ inputs and decay constants, while the $D_{s1}(2536)$ width is particularly sensitive to the mixing angle because the physical amplitude contains a cancellation between the axial and tensor-current components. The same ranking is visualized in Fig. 3.




Figure 3: Ranked one-at-a-time uncertainty sources. The plotted quantity is half of the 16–84 percentile width for each group.

16.1 Alternative lattice normalization of the leading-twist photon DA

The dominant uncertainty in the legacy scan is the combination $\chi\langle\bar{s}s\rangle$ entering the leading-twist photon DA. The Rohrwild/BBK-style baseline treats χ and $\langle\bar{s}s\rangle$ as separate inputs. A recent lattice calculation by Bacchio, Becirevic, Gagliardi, and Sanfilippo gives the strange-quark normalization directly,

$$f_{\gamma,s}^\perp(2 \text{ GeV}) = -51.0(1.8) \text{ MeV}, \quad (134)$$

where

$$f_{\gamma,s}^\perp(\mu) = \chi_s(\mu)\langle\bar{s}s\rangle(\mu). \quad (135)$$

To compare with the working $\mu = 1$ GeV setup, the script

`scripts/lattice_photon_normalization_comparison.py`

uses a leading-order tensor-current running factor $f_{\gamma,s}^\perp(1 \text{ GeV})/f_{\gamma,s}^\perp(2 \text{ GeV}) = 1.08$, giving

$$f_{\gamma,s}^\perp(1 \text{ GeV}) = -55.1(1.9) \text{ MeV}. \quad (136)$$

Only the leading-twist normalization is replaced in this comparison; the higher-twist and three-particle terms that contain $\langle \bar{s}s \rangle$ remain explicit. This makes the comparison conservative and easy to audit.

Input scenario	Γ_{2460} (keV)	Γ_{2536} (keV)
Legacy $\chi\langle\bar{s}s\rangle$, fixed θ	36.7 [26.8, 50.2]	5.60 [2.56, 11.0]
Lattice $f_{\gamma,s}^\perp$, fixed θ	14.4 [11.4, 18.8]	0.68 [0.12, 1.88]
Legacy $\chi\langle\bar{s}s\rangle$, θ scan	36.3 [24.8, 50.2]	5.74 [1.86, 12.9]
Lattice $f_{\gamma,s}^\perp$, θ scan	14.6 [11.3, 18.6]	0.64 [0.06, 2.50]

Table 2: Comparison of the legacy photon-DA normalization with the lattice-normalized leading-twist photon DA. The intervals are 16–84 percentiles from 500 random points per ensemble.

The lattice-normalized scenario substantially reduces the uncertainty and also shifts the central widths downward. This should be presented as an important alternative input scenario rather than folded silently into the legacy result, because it changes the central leading-twist normalization. The corresponding histograms are shown in Fig. 4.




Figure 4: Fixed-angle Monte Carlo comparison between the legacy $\chi\langle\bar{s}s\rangle$ treatment and the lattice-normalized $f_{\gamma,s}^\perp$ treatment.

For direct comparison with the legacy Monte Carlo histograms in Figs. 1 and 2, the same script also produces Gaussian-overlay plots for the lattice-normalized scenario. For fixed $\theta = 35.3^\circ$, the descriptive Gaussian summaries are

$$\Gamma_{2460}^{\text{lat. } f_\gamma^\perp} = 14.91 \pm 3.69 \text{ keV}, \quad (137)$$

$$\Gamma_{2536}^{\text{lat. } f_\gamma^\perp} = 0.97 \pm 0.97 \text{ keV}. \quad (138)$$

For the mixing-angle scan they are

$$\Gamma_{2460}^{\text{lat. } f_{\gamma}^{\perp}} = 14.94 \pm 3.77 \text{ keV}, \quad (139)$$

$$\Gamma_{2536}^{\text{lat. } f_{\gamma}^{\perp}} = 1.21 \pm 1.45 \text{ keV}. \quad (140)$$

The $D_{s1}(2536)$ distribution is visibly non-Gaussian near zero because the amplitude is cancellation-driven, so the percentile intervals in Table 2 remain the safer quoted uncertainties.




Figure 5: Lattice-normalized Monte Carlo width distributions for fixed $\theta = 35.3^\circ$. The black curves are Gaussian overlays using the sample mean and standard deviation; dashed vertical lines mark the 16–84 percentile interval.




Figure 6: Lattice-normalized Monte Carlo width distributions when the mixing angle is sampled over $25^\circ \leq \theta \leq 45^\circ$.

16.2 Recommended D_{s1} results table

For the manuscript, the safest way to report the width predictions is not a Gaussian mean and standard deviation, but the median with asymmetric 16–84 percentile intervals. This is especially important for $D_{s1}(2536)$, where the width is positive definite and the amplitude can pass close to zero because of the axial–tensor cancellation.

There is also a scale-convention issue that should be stated explicitly. The photon-DA normalizations in the table are quoted at $\mu = 1$ GeV, whereas the quark masses used in the hard amplitudes are the usual $\overline{\text{MS}}$ inputs, e.g. $m_s(2 \text{ GeV})$ and $m_c(m_c)$. This is a standard phenomenological input convention, but it is not a fully matched single-scale setup. A strict scale-consistent variant would evolve m_s , $\langle \bar{s}s \rangle$, and either χ_s or $f_{\gamma,s}^\perp$ to a common factorization scale before rerunning the sum rule. We therefore flag the present rows as mixed-scale inputs rather than interpreting them as a separate scale-consistent prediction.

Scenario	Photon input	implied χ_s	Mixing input	Scale check	Γ_{2460} (keV)	Γ_{2536} (keV)
Legacy	$\chi_s(1 \text{ GeV}) = 3.15 \pm 0.30$ (Rohrwild/BBK)	3.15 [2.89, 3.47]	$\theta = 35.3^\circ$	mixed-scale input	$36.7^{+13.5}_{-9.9}$	$5.60^{+5.38}_{-3.04}$
Legacy	$\chi_s(1 \text{ GeV}) = 3.15 \pm 0.30$ (Rohrwild/BBK)	3.16 [2.86, 3.42]	$25^\circ \leq \theta \leq 45^\circ$	mixed-scale input	$36.3^{+13.9}_{-11.5}$	$5.74^{+7.14}_{-3.88}$
Lattice	$f_{\gamma,s}^\perp(1 \text{ GeV}) = -55.1 \pm 1.9$ MeV (Bacchio et al.)	4.99 [4.24, 6.05]	$\theta = 35.3^\circ$	mixed-scale input	$14.4^{+4.3}_{-3.0}$	$0.68^{+1.20}_{-0.56}$
Lattice	$f_{\gamma,s}^\perp(1 \text{ GeV}) = -55.1 \pm 1.9$ MeV (Bacchio et al.)	5.02 [4.25, 6.09]	$25^\circ \leq \theta \leq 45^\circ$	mixed-scale input	$14.6^{+4.0}_{-3.4}$	$0.64^{+1.86}_{-0.58}$

Table 3: Recommended reporting table for the $D_{s1} \rightarrow D_s \gamma$ widths. Entries are medians with 16–84 percentile intervals. The lattice scenario replaces only the leading-twist normalization $\chi_s \langle \bar{s}s \rangle$ by $f_{\gamma,s}^\perp$; higher-twist condensate terms are left explicit. The implied χ_s column is derived from $f_{\gamma,s}^\perp / \langle \bar{s}s \rangle$ in the lattice rows, so it inherits the strange-condensate spread. The scale-check column records that the present numerical rows use conventional mixed-scale phenomenological inputs; a fully matched-scale prediction should be generated as a separate rerun.

The legacy rows should be retained for comparison with the older photon-DA literature. The lattice-normalized rows are the more modern input scenario and give a much tighter prediction for $D_{s1}(2460)$, while $D_{s1}(2536)$ remains best described by an asymmetric interval because the central amplitude is cancellation-suppressed. The machine-readable version of Table 3 is `outputs/ds1_recommended_results_table.csv`.

17 Central Borel Stability

For the central input set and $\theta = 35.3^\circ$, the script

`scripts/stage2_stability_plots.py`

produces the Borel-stability curves shown in Fig. 7. The curves are smooth over $3 \leq M^2 \leq 6 \text{ GeV}^2$. At fixed $\sqrt{s_0}$, the dependence on M^2 is mild; changing $\sqrt{s_0}$ from 2.50 to 2.60 GeV shifts the normalization more visibly and should therefore remain part of the quoted systematic uncertainty.

Across the full displayed window the central ranges are

$$\Gamma[D_{s1}(2460) \rightarrow D_s \gamma] = 30.6 \text{ to } 44.5 \text{ keV}, \quad (141)$$

$$\Gamma[D_{s1}(2536) \rightarrow D_s \gamma] = 3.95 \text{ to } 8.10 \text{ keV}. \quad (142)$$

The corresponding amplitude spans are smaller:

$$G_{2460} = -0.460 \text{ to } -0.381 \text{ GeV}^{-1}, \quad (143)$$

$$G_{2536} = -0.162 \text{ to } -0.113 \text{ GeV}^{-1}. \quad (144)$$

This is the expected pattern: the widths amplify the residual sum-rule variation because $\Gamma \propto G^2 |\vec{q}|^3$.

18 Publication Figures and Phenomenology

The remaining publication-oriented figures are generated by




Figure 7: Central matched Stage 2 stability for $D_{s1}(2460) \rightarrow D_s \gamma$ and $D_{s1}(2536) \rightarrow D_s \gamma$ at $\theta = 35.3^\circ$. The three curves correspond to $\sqrt{s_0} = 2.50, 2.55, 2.60$ GeV.

Figure 8 shows the continuum-threshold dependence at fixed representative values of M^2 . It complements Fig. 7: the curves are monotonic in $\sqrt{s_0}$, so the continuum threshold should be quoted as an important systematic.




Figure 8: Continuum-threshold dependence of the matched Stage 2 widths for $\theta = 35.3^\circ$ at three representative Borel points.

Figure 9 shows the mixing-angle dependence. The $D_{s1}(2460)$ width remains at the tens-of-keV level over a broad range of θ , while the $D_{s1}(2536)$ width is strongly suppressed near the large-angle region because its amplitude involves a cancellation between the axial and tensor-current components.




Figure 9: Mixing-angle dependence at $M^2 = 4.5$ GeV² and $\sqrt{s_0} = 2.55$ GeV. The dashed line marks $\theta = 35.3^\circ$.

The cancellation pattern is displayed more explicitly in Fig. 10. At the central point $M^2 = 4.5$ GeV², $\sqrt{s_0} = 2.55$ GeV, the $D_{s1}(2536)$ amplitude receives opposite-sign contributions from the axial Stage 1 term and the hard tensor-current term. The three-particle DA corrections are numerically small in the final physical amplitudes, but they are included in the quoted Stage 2 result.

Finally, Fig. 11 compares the present result for $D_{s1}(2460) \rightarrow D_s \gamma$ with representative estimates collected by Colangelo, De Fazio and Ozpineci. The present result is higher than the




Figure 10: Contribution breakdown for the physical couplings at the central Borel and threshold point. The labels “A” and “B” refer to the axial-current and tensor-current components before physical mixing.

original axial-current LCSR interval, but it remains the same order of magnitude. It should not be advertised as explaining a non-observation by making the $D_{s1}(2460) \rightarrow D_s \gamma$ width tiny. Instead, the robust phenomenological statement from this calculation is that the $D_{s1}(2536) \rightarrow D_s \gamma$ channel is much more cancellation-sensitive and can be substantially smaller.

19 Validation Checks Before the B_{s1} Extension

The script

`scripts/validation_checks.py`

collects several checks that should be repeated whenever the numerical implementation is modified. The first check is an axial-current benchmark against Colangelo, De Fazio and Ozpineci, using paper-like input choices and the BBK value $\kappa = 0.2$ for the relevant photon DA parameter. Over $3 \leq M^2 \leq 6 \text{ GeV}^2$ and $2.50 \leq \sqrt{s_0} \leq 2.60 \text{ GeV}$, the present controlled benchmark gives

$$g_1 = -0.340 \text{ to } -0.237 \text{ GeV}^{-1}, \quad \Gamma = 11.8 \text{ to } 24.3 \text{ keV}. \quad (145)$$

The published comparison band is $g_1 = -0.37 \text{ to } -0.29 \text{ GeV}^{-1}$ and $\Gamma = 19 \text{ to } 29 \text{ keV}$. This is close enough in sign and scale to validate the implementation, but it also shows that an exact literature reproduction requires matching every convention and input choice. We should present this as a benchmark check, not as an independent final phenomenological result.

The same validation script checks the photon DA normalization and the three-particle integration closure:

$$\int_0^1 du \varphi_\gamma(u) = 1.0000000000, \quad (146)$$

$$\max_u |\psi^a(u) - \psi^a(1-u)| = 4.44 \times 10^{-15}, \quad (147)$$

$$\max_u |\psi^v(u) + \psi^v(1-u)| = 1.78 \times 10^{-15}, \quad (148)$$

$$F_1 - F_1^{(1)} - F_1^{(2)} = 0. \quad (149)$$




Figure 11: Comparison of the present $D_{s1}(2460) \rightarrow D_s \gamma$ width with representative literature estimates. The Colangelo LCSR, VMD and quark-model values are from the comparison table in Colangelo, De Fazio and Ozpineci (hep-ph/0505195).

These are numerical quadrature checks of the DA implementation and the split integration domains.

The independent FeynCalc script `scripts/ward_transversality_check.wl` checks the selected E1 tensor $S_{\mu\nu} = p_\nu q_\mu - (p \cdot q)g_{\mu\nu}$. With $q^2 = 0$, $\varepsilon \cdot q = 0$, $P = p + q$, and $\eta \cdot P = 0$, it gives

$$S_{\mu\nu} q^\nu = 0, \quad (150)$$

$$[(\eta \cdot \varepsilon)(p \cdot q) - (\eta \cdot q)(p \cdot \varepsilon)]_{\varepsilon \rightarrow q} = 0, \quad (151)$$

$$P^\mu S_{\mu\nu} \varepsilon^\nu = 0, \quad (152)$$

$$\frac{S_{\mu\nu} S^{\mu\nu}}{2(p \cdot q)^2} = 1. \quad (153)$$

The first two equations are the photon Ward check, the third confirms compatibility with the physical initial axial-meson polarization, and the last line is the projector normalization.

At the central point $M^2 = 4.5 \text{ GeV}^2$ and $\sqrt{s_0} = 2.55 \text{ GeV}$, the hard tensor-current charge decomposition is

$$G_B^{\text{hard}} = -4.092 \times 10^{-1} \text{ GeV}^{-1}, \quad (154)$$

$$G_B^{\text{hard}}(e_c \text{ line}) = -2.499 \times 10^{-1} \text{ GeV}^{-1}, \quad (155)$$

$$G_B^{\text{hard}}(e_s \text{ line}) = -1.594 \times 10^{-1} \text{ GeV}^{-1}. \quad (156)$$

This confirms that hard photon emission from both quark lines is numerically important and must remain in the baseline.

20 First $B_{s1} \rightarrow B_s \gamma$ Extension

The same matched Stage-2 machinery can be reused for the bottom-strange system after the replacement $m_c \rightarrow m_b$ and $e_c \rightarrow e_b = -1/3$. The script

`scripts/bs1_stage2_baseline.py`

Diagnostic case	G_A (GeV ⁻¹)	G_B	Γ_{2460} (keV)	Γ_{2536} (keV)
central	-0.357	-0.263	37.18	6.04
$m_s \rightarrow 0$ numerical proxy	-0.175	-0.210	15.59	0.15
$m_s = m_c$ toy	+0.535	+0.273	59.46	24.22
$e_c = 0$ toy	+0.102	-0.013	0.50	2.57
$e_s = 0$ toy	-0.459	-0.250	46.30	16.48
light- u OPE toy	-1.040	-0.438	192.96	110.02
light- d OPE toy	-0.109	-0.176	8.94	0.05

Table 4: Limiting-case diagnostics from `outputs/validation_limiting_cases.csv`. The $m_s \rightarrow 0$ and light-quark rows use $m_q = 10^{-4}$ GeV as a numerical regulator of the present hard-line spectral-density representation. The light- u and light- d rows are OPE-level charge and condensate substitutions only, not physical nonstrange $D_1 \rightarrow D\gamma$ predictions.

performs this first pass. Internally the reusable functions still use the key `mc`; in this section it should be read as m_b .

The first-pass numerical inputs are deliberately conservative working inputs: $m_b(m_b) = 4.18$ GeV, $m_s(2 \text{ GeV}) = 0.093$ GeV, $m_{B_s} = 5.36692$ GeV, and $f_{B_s} = 0.2303$ GeV (FLAG Review 2024). For the axial and tensor B_{s1} decay constants we use the temporary heavy-quark scaling estimate $f_H \sqrt{m_H} = \text{const.}$ from the D_{s1} central values. Thus these rows are not yet final paper predictions; they are a controlled baseline for the bottom-sector calculation.

Two mass targets are shown. The observed narrow state is treated as the high-combination state,

$$G_{\text{high}} = \cos \theta G_A - \sin \theta G_B, \quad m_{B_{s1}} = 5.82870 \text{ GeV}. \quad (157)$$

Because the lower 1^+ bottom-strange partner is not firmly established, the second row uses $m_{B_{s1}} = 5.720$ GeV only as a lower-partner diagnostic, with

$$G_{\text{low}} = \sin \theta G_A + \cos \theta G_B. \quad (158)$$

The scan uses $8 \leq M^2 \leq 14$ GeV², $\theta = 35.3^\circ$, and $\sqrt{s_0} = 6.05, 6.15, 6.25$ GeV for the observed state (5.95, 6.05, 6.15 GeV for the lower diagnostic state).

Target	Photon input	G (GeV ⁻¹)	Γ (keV)
$B_{s1}(5830)$ high combo	legacy χ_s (Rohrwild/BBK)	0.0358 [0.0218, 0.0688]	0.27 [0.10, 1.01]
$B_{s1}(5830)$ high combo	lattice $f_{\gamma,s}^\perp$ (Bacchio et al.)	0.1496 [0.119, 0.216]	4.75 [3.02, 9.88]
lower 1^+ diagnostic	legacy χ_s (Rohrwild/BBK)	0.589 [0.502, 0.761]	33.8 [24.6, 56.5]
lower 1^+ diagnostic	lattice $f_{\gamma,s}^\perp$ (Bacchio et al.)	0.847 [0.725, 1.09]	69.9 [51.2, 116]

Table 5: First matched Stage-2 $B_{s1} \rightarrow B_s \gamma$ baseline. Brackets show the range over the Borel and continuum-threshold grid, not a full uncertainty interval. The B_{s1} decay constants are working heavy-quark-scaled inputs, so these values should be refined after dedicated B_{s1} hadronic inputs are fixed.

The qualitative behavior mirrors the D_{s1} calculation: the observed high-combination state is cancellation-sensitive, while the lower-combination diagnostic is much larger. The difference between the legacy and lattice photon-normalization rows is again substantial, so the bottom-sector analysis should keep both scenarios until the input convention is finalized. The machine-readable grid is `outputs/bs1_stage2_baseline.csv`.

The companion script

`scripts/bs1_uncertainty_and_plots.py`

performs a 500-point Monte Carlo scan for each state, photon-input scenario, and mixing-angle treatment. The observed $B_{s1}(5830)$ row samples $8 \leq M^2 \leq 14 \text{ GeV}^2$, $6.05 \leq \sqrt{s_0} \leq 6.25 \text{ GeV}$, the same quark and photon-DA inputs used in the D_{s1} scan, and assigns a conservative 20% width to the temporary heavy-quark-scaled B_{s1} axial and tensor decay constants. The lower-state diagnostic uses the lattice-inspired model mass $m_{B_{s1}} = 5.750 \pm 0.026 \text{ GeV}$ from Lang, Mohler, Prelovsek and Woloshyn.

Target	Photon input	Mixing input	implied χ_s	$G \text{ (GeV}^{-1}\text{)}$	$\Gamma \text{ (keV)}$
$B_{s1}(5830)$ high combo	legacy χ_s (Rohrwild/BBK)	$\theta = 35.3^\circ$	3.12 [2.83, 3.40]	+0.0273 [−0.0592, +0.133]	1.04 [0.068, 4.45]
$B_{s1}(5830)$ high combo	legacy χ_s (Rohrwild/BBK)	$25^\circ \leq \theta \leq 45^\circ$	3.14 [2.84, 3.47]	+0.0242 [−0.0992, +0.158]	1.51 [0.149, 7.76]
$B_{s1}(5830)$ high combo	lattice $f_{\gamma,s}^\perp$ (Bacchio et al.)	$\theta = 35.3^\circ$	5.03 [4.26, 6.06]	+0.143 [+0.0076, +0.298]	5.30 [0.454, 18.9]
$B_{s1}(5830)$ high combo	lattice $f_{\gamma,s}^\perp$ (Bacchio et al.)	$25^\circ \leq \theta \leq 45^\circ$	5.04 [4.22, 5.96]	+0.146 [−0.019, +0.337]	5.32 [0.588, 24.7]
lower 1^+ diagnostic	legacy χ_s (Rohrwild/BBK)	$\theta = 35.3^\circ$	3.16 [2.85, 3.44]	+0.609 [+0.487, +0.793]	45.8 [28.3, 78.7]
lower 1^+ diagnostic	lattice $f_{\gamma,s}^\perp$ (Bacchio et al.)	$\theta = 35.3^\circ$	4.97 [4.23, 6.00]	+0.885 [+0.738, +1.08]	97.0 [64.6, 156]

Table 6: Monte Carlo $B_{s1} \rightarrow B_s \gamma$ baseline. Entries are medians with 16–84 percentile intervals. The observed $B_{s1}(5830)$ amplitude is cancellation-sensitive, so the width should be quoted with asymmetric percentile intervals rather than a Gaussian error.




Figure 12: Monte Carlo width distributions for the observed $B_{s1}(5830) \rightarrow B_s \gamma$ high-combination state. The legacy and lattice-normalized photon inputs are shown separately for fixed $\theta = 35.3^\circ$ and for the mixing-angle scan.

The most important lesson from Table 6 is that the observed $B_{s1}(5830)$ amplitude can pass close to zero over the accepted input range. This makes the legacy scenario compatible with a sub-keV-to-few-keV width, while the lattice-normalized leading-twist input shifts the median to about 5 keV but leaves a long upper tail. A final quoted B_{s1} prediction therefore needs dedicated B_{s1} decay constants before we should advertise the central value too strongly. The machine-readable Monte Carlo outputs are `outputs/bs1_stage2_mc_scan.csv` and `outputs/bs1_stage2_mc_summary.csv`.

21 Combined D_{s1} and B_{s1} Summary

Table 7 collects the current headline width predictions in one place. The D_{s1} rows are the current final baseline of this note. The B_{s1} rows are working baseline results because the axial and tensor B_{s1} decay constants are still temporary heavy-quark-scaled inputs. All entries are medians with 16–84 percentile intervals.

The machine-readable version of Table 7 is `outputs/combined_recommended_results_table.csv`.




Figure 13: Central $B_{s1}(5830) \rightarrow B_s \gamma$ stability curves for the high-combination amplitude, width, and unmixed G_A, G_B components.

Sector	Decay	Photon input	Mixing input	G (GeV^{-1})	Γ (keV)
D_s final	$D_{s1}(2460) \rightarrow D_s \gamma$	legacy χ_s (Rohrwild/BBK)	$\theta = 35.3^\circ$	$-0.418 [-0.489, -0.357]$	$36.7^{+13.5}_{-9.9}$
D_s final	$D_{s1}(2536) \rightarrow D_s \gamma$	legacy χ_s (Rohrwild/BBK)	$\theta = 35.3^\circ$	$-0.134 [-0.188, -0.0908]$	$5.60^{+5.38}_{-3.04}$
D_s final	$D_{s1}(2460) \rightarrow D_s \gamma$	lattice $f_{\gamma,s}^\perp$ (Bacchio et al.)	$\theta = 35.3^\circ$	$-0.262 [-0.298, -0.233]$	$14.4^{+4.3}_{-3.0}$
D_s final	$D_{s1}(2536) \rightarrow D_s \gamma$	lattice $f_{\gamma,s}^\perp$ (Bacchio et al.)	$\theta = 35.3^\circ$	$-0.0468 [-0.0778, -0.0200]$	$0.68^{+1.20}_{-0.56}$
B_s working	$B_{s1}(5830) \rightarrow B_s \gamma$	legacy χ_s (Rohrwild/BBK)	$\theta = 35.3^\circ$	$+0.0273 [-0.0592, +0.133]$	$1.04^{+3.41}_{-0.97}$
B_s working	$B_{s1}(5830) \rightarrow B_s \gamma$	lattice $f_{\gamma,s}^\perp$ (Bacchio et al.)	$\theta = 35.3^\circ$	$+0.143 [+0.0076, +0.298]$	$5.30^{+13.6}_{-4.85}$
B_s working	$B_{s1}(5830) \rightarrow B_s \gamma$	legacy χ_s (Rohrwild/BBK)	$25^\circ \leq \theta \leq 45^\circ$	$+0.0242 [-0.0992, +0.158]$	$1.51^{+6.25}_{-1.36}$
B_s working	$B_{s1}(5830) \rightarrow B_s \gamma$	lattice $f_{\gamma,s}^\perp$ (Bacchio et al.)	$25^\circ \leq \theta \leq 45^\circ$	$+0.146 [-0.0190, +0.337]$	$5.32^{+19.4}_{-4.73}$

Table 7: Current combined summary of D_{s1} and B_{s1} radiative form factors and widths. The G column is the E1 coupling in the convention of Eq. (37). The B_s rows should be treated as working results until dedicated B_{s1} decay constants replace the heavy-quark-scaled temporary inputs.

22 Comparison With Experimental Information

The sum rule predicts a partial radiative width, $\Gamma_\gamma \equiv \Gamma(H_1 \rightarrow H\gamma)$, while experiments often quote branching fractions, total widths, or spectroscopy measurements in other decay channels. Therefore the comparison is not always one-to-one. In Table 8, the “interpretation” column has the following meaning:

- for $D_{s1}(2460)$, the measured branching fraction is used to infer the total-width scale that would be implied by our partial radiative width;
- for $D_{s1}(2536)$, the measured total width is used to infer the radiative branching fraction implied by our partial width;
- for $B_{s1}(5830)$, no direct radiative measurement is available, so the row is a prediction for future experimental searches rather than a numerical comparison.

Decay	Theory input	Experimental information	Interpretation
$D_{s1}(2460) \rightarrow D_s\gamma$	legacy χ_d : 36.7 [26.8, 50.2] keV	$\mathcal{B}(D_{s1}(2460) \rightarrow D_s\gamma) = (16 \pm 4 \pm 3)\%$ (BaBar)	implies $\Gamma_{\text{tot}} \sim 0.23$ MeV; compatible with a very narrow state
$D_{s1}(2460) \rightarrow D_s\gamma$	lattice $f_{D_s^*}^+$: 14.4 [11.4, 18.7] keV	$\mathcal{B}(D_{s1}(2460) \rightarrow D_s\gamma) = (16 \pm 4 \pm 3)\%$ (BaBar)	implies $\Gamma_{\text{tot}} \sim 0.09$ MeV; also compatible with a very narrow state
$D_{s1}(2536) \rightarrow D_s\gamma$	legacy χ_d : 5.60 [2.56, 10.98] keV	$\Gamma_{\text{tot}}(D_{s1}(2536)) = 0.92 \pm 0.05$ MeV (BaBar)	implies $\mathcal{B}_\gamma \sim 0.6\%$; experimentally small channel
$D_{s1}(2536) \rightarrow D_s\gamma$	lattice $f_{D_s^*}^+$: 0.68 [0.12, 1.88] keV	$\Gamma_{\text{tot}}(D_{s1}(2536)) = 0.92 \pm 0.05$ MeV (BaBar)	implies $\mathcal{B}_\gamma \sim 0.07\%$; strongly cancellation suppressed
$B_{s1}(5830) \rightarrow B_s\gamma$	legacy χ_d : 1.04 [0.068, 4.45] keV	no direct radiative measurement; observed mainly in $B^{(*)}K$ spectroscopy	prediction for future searches; comparison is qualitative
$B_{s1}(5830) \rightarrow B_s\gamma$	lattice $f_{B_s^*}^+$: 5.30 [0.454, 18.9] keV	no direct radiative measurement; observed mainly in $B^{(*)}K$ spectroscopy	prediction for future searches; branching fraction depends on the total width

Table 8: Comparison of the present radiative-width predictions with available experimental information. The BaBar $D_{s1}(2460)$ branching fraction is from the $B \rightarrow D^{(*)}D_{sJ}^{(*)}$ analysis. The $D_{s1}(2536)$ total width is the BaBar width measurement. The $B_{s1}(5830)$ has no direct radiative branching-fraction measurement at present; CMS and LHCb measurements are spectroscopy measurements in hadronic channels.

Numerically, the $D_{s1}(2460)$ rows say that the present partial-width prediction is consistent with a total width well below the MeV scale if the measured radiative branching fraction is used. The $D_{s1}(2536)$ rows say something sharper: with the measured total width near 0.92 MeV, the radiative branching fraction is expected to be below the percent level, especially in the lattice-normalized photon-input scenario. For $B_{s1}(5830)$ the table should be read as a target for future experimental searches, not as a test against an existing radiative measurement.

The machine-readable version of Table 8 is `outputs/experimental_comparison_table.csv`.

23 Missing Or Useful Paper Material

Before moving from the D_{s1} paper draft to the B_{s1} extension, the following items are useful to add or verify.

- A compact input table in the paper text: masses, decay constants, condensates, χ , DA parameters, Borel windows, and continuum thresholds. The final numerical scan already uses these values, but the paper needs a reader-facing table.
- A short paragraph on convention matching. The sign of χ , the Levi-Civita convention, and the tensor-current normalization $f_B = f_T m_{D_{s1}} / (m_c + m_s)$ should be stated explicitly because they move signs and absolute normalizations.
- A compact uncertainty table for the paper. The Monte Carlo and one-at-a-time outputs are now available, but the manuscript should present only the main quoted band and the dominant sources, not every diagnostic row.
- A physical light-quark counterpart requires new hadronic input tables. The OPE-level light- u and light- d diagnostics above are useful algebra checks, but physical $D_1 \rightarrow D\gamma$ or $B_1 \rightarrow B\gamma$ predictions need the corresponding hadron masses, decay constants, thresholds, Borel windows, condensates, and mixing convention.

- For the B_{s1} extension, the same code structure can be reused after replacing $m_c \rightarrow m_b$, $m_{D_s} \rightarrow m_{B_s}$, $m_{D_{s1}} \rightarrow m_{B_{s1}}$, and updating the tensor decay constant and threshold window. The hard-photon terms should be recomputed rather than scaled by hand, because the charm and bottom propagator denominators weight the heavy-line emission differently.