


UNIT 4 :-

Q.1) Solve using Backward multistage graph



Soln :- Backward Approach :-

$$bcost(i, j) = \min \{ c(l, j) + bcost(i-1, l) \}$$

* Stage 1 :-

$$bcost(1, 1) = 0$$

$$d(1, 1) =$$

$$bcost(1, 1)$$

Vertex	1	2	3	4	5	6	7	8	9
Cost	0	5	2	8	7	8	9	9	12
distance	0	1	1	1	3	3	2	4	5

* Stage 2 :-

$$bcost(2, 2) = \min \{ c(1, 2) + bcost(1, 1) \}$$

$$bcost(2, 2) = \min \{ 5 + 0 \} = 5$$

$$d(2, 2) = 1$$

$$b\text{Cost}(2,3) = \min \{ c(1,3) + b\text{Cost}(1,1), \dots \}$$

$$b\text{Cost}(2,3) = \min_2 \{ 2+0 \}$$

$$d(2,3) = 1$$

* Stage 3:-

$$b\text{Cost}(3,4) = \min \{ c(2,4) + b\text{Cost}(2,2), \\ c(3,4) + b\text{Cost}(2,3) \}$$

$$= \min \{ 3+5, 6+2 \}$$

$$b\text{Cost}(3,4) = \min \{ 8, 8 \} = 8$$

$$d(3,4) = 2 \text{ or } 3$$

$$b\text{Cost}(3,5) = \min \{ c(3,5) + b\text{Cost}(2,3), \dots \}$$

$$b\text{Cost}(3,5) = \min \{ 5+5, 5+2 \} = 7$$

$$d(3,5) = 3$$

$$b\text{Cost}(3,6) = \min \{ c(2,6) + b\text{Cost}(2,2), \dots \}$$

$$c(3,6) + b\text{Cost}(2,3) \}$$

$$= \min \{ 3+5, 8+2 \}$$

$$b\text{Cost}(3,6) = \min \{ 8, 10 \} = 8$$

$$d(3,6) = 2$$

*

Stage 4:-

$$\begin{aligned} b\text{cost}(4,7) &= \min \{ c(4,7) + b\text{cost}(3,4), \\ &\quad c(5,7) + b\text{cost}(3,5), \\ &\quad c(6,7) + b\text{cost}(3,6) \} \\ &= \min \{ 1+8, 6+7, 6+8 \} \end{aligned}$$

$$b\text{cost}(4,7) = \min \{ 9, 13, 14 \} = 9$$

$$d(4,7) = 4.$$

$$\begin{aligned} b\text{cost}(4,8) &= \min \{ c(4,8) + b\text{cost}(3,4), \\ &\quad c(5,8) + b\text{cost}(3,5), \\ &\quad c(6,8) + b\text{cost}(3,6) \} \\ &= \min \{ 4+8, 2+7, 2+8 \} \\ &= \min \{ 12, 9, 10 \} \\ b\text{cost}(4,8) &= 9 \\ d(4,8) &= 5. \end{aligned}$$

* Stages :-

$$\begin{aligned}
 b\text{cost}(5, 2) &= \min \{ c(7, 2) + b\text{cost}(4, 7), \\
 &\quad c(8, 2) + b\text{cost}(4, 8) \} \\
 &= \min \{ 7 + 9, 8 + 9 \} \\
 &= \min \{ 16, 12 \} \\
 b\text{cost}(5, 2) &= 12
 \end{aligned}$$

$$d(5, 2) = 8$$

Shortest path

$$\begin{aligned}
 d(5, 2) &= 8 \\
 d(4, 8) &= 5 \\
 d(3, 5) &= 3 \\
 d(2, 3) &= 1
 \end{aligned}$$


* Shortest Path :-

$$= 9 - 8 - 5 - 3 - 1$$

$$= 3 + 2 + 5 + 2 = \underline{\underline{12}}$$

Q. 2)

What is multistage graph? For the following graph find shortest path from source to destination. * (Forward Approach)



Solutn :-

$$\text{Cost}(i, j) = \min \{ c(j, l) + \text{Cost}(i+1, l) \}$$

Vertex	1	2	3	4	5	6	7	8	9	10	11	12
Cost	8	8	8	12	14	8	6	8	5	2	4	0
dist	2	6	7	8	7	10	10	10	12	12	12	12

$$\text{Cost}(5, 12) = 0$$

$$\text{Cost}(4, 9) = 5$$

$$\text{Cost}(4, 10) = 2$$

$$\text{Cost}(4, 11) = 4$$

$$\text{Cost}(3, 6) = \min \{ c(6, 2) + \text{Cost}(4, 2),$$

$$c(6, 10) + \text{Cost}(4, 10) \}$$

$$\text{Cost}(3, 6) = \min \{ 5 + 5, 6 + 2 \} = \min \{ 10, 8 \}$$

Shrikrupa

$$d(3, 6) = 10$$

$$\text{Cost}(3,7) = \min \{ c(7,2) + \text{cost}(4,2),$$

$$(c(5,2) + 20) + (c(7,10) + \text{cost}(4,10)) \}$$

$$= \min \{ 3+5, 4+2 \}$$

$$\text{Cost}(3,7) = \min \{ 8, 6 \}$$

$$\text{d}(3,7) = 10$$

$$\text{Cost}(3,8) = \min \{ c(8,10) + \text{cost}(4,10),$$

$$c(8,11) + \text{cost}(4,11) \}$$

$$= \min \{ 6+2, 5+4 \}$$

$$= \min \{ 8, 9 \}$$

$$\text{Cost}(3,8) = 8$$

$$\text{d}(3,8) = 10$$

$$\text{Cost}(2,2) = \min \{ c(2,6) + \text{cost}(3,6),$$

$$c(2,7) + \text{cost}(3,7),$$

$$c(2,8) + \text{cost}(3,8) \}$$

$$= \min \{ 1+18, 2+16, 4+18 \}$$

$$\text{Cost}(2,2) = \min \{ 9, 8, 12 \} = 8$$

$$\text{d}(2,2) = 7$$

$$\text{Cost}(2,3) = \min \{ c(3,6) + \text{cost}(3,6),$$

$$c(3,7) + \text{cost}(3,7), \}$$

$$= \min \{ 7 + 18, 2 + 16 \}$$

$$\text{Cost}(2,3) = \min \{ 15, 8 \} = 8$$

$$d(2,3) = 7$$

$$\text{Cost}(2,4) = \min \{ c(4,8) + \text{cost}(3,8) \}$$

$$= \min \{ 11 + 18 \}$$

$$= 19$$

$$d(2,4) = 8$$

$$\text{Cost}(2,5) = \min \{ c(5,7) + \text{cost}(3,7) \},$$

$$c(5,8) + \text{cost}(3,8) \}$$

$$= \min \{ 8 + 6, 11 + 8 \}$$

$$\text{Cost}(2,5) = \min \{ 14, 18 \} = 14$$

$$d(2,5) = 7$$

$$\text{Cost}(1,1) = \min \{c(1,2) + \text{cost}(2,2),$$

$$c(1,3) + \text{cost}(2,3),$$

$$c(1,4) + \text{cost}(2,4),$$

$$c(1,5) + \text{cost}(2,5)$$

$$= \min \{ 2+8, 3+8, 7+8, 9+14 \}$$

$$= \min \{ 8, 11, 15, 23 \}$$

$$\text{Cost}(1,1) = 10 = 10$$

$$d(1,1) = 2$$

→ shortest path.

$$d(2,2) = 2$$


$$d(2,7) = 7$$

$$d(3,7) = 10$$

$$d(4,10) = 2$$

if

path




Shortest dist. $\Rightarrow 2+2+4+2 = 10$

From source to destination

Date _____
Page _____

5) Find the minimum cost path from "s" to "t" in multistage graph shown. Using forward approach.



$\text{cost}(i, j) = \min \{ c(j, l) + \text{cost}(i+1, l) \}$

cutex	1	2	3	4	5	6	7	8	9	10	11	12
Cost	16	7	9	18	15	7	5	7	4	2	5	0
dist	2	7	6	8	8	10	10	10	12	12	12	12

$$\text{Cost}(5, 12) = 0$$

$$\text{Cost}(4, 9) = 4$$

$$\text{Cost}(4, 10) = 2$$

$$\text{Cost}(4, 11) = 5$$

$$\begin{aligned} \text{Cost}(3, 6) &= \min \{ c(6, 9) + \text{cost}(4, 9), \\ &\quad c(6, 10) + \text{cost}(4, 10) \} \end{aligned}$$

$$= \min \{ 6 + 4, 5 + 2 \}$$

$$\text{Cost}(3, 6) = \min \{ 10, 7 \} = 7$$

$$d(3, 6) = 10$$

$$\text{Cost}(3,7) = \min \{ c(7,5) + \text{Cost}(4,5),$$

$$c(7,10) + \text{Cost}(4,10) \}$$

$$= \min \{ 4+4, 3+2 \}$$

$$= \min \{ 8, 5 \}$$

$$\text{Cost}(3,7) = 5$$

$$d(3,7) = 10$$

$$\text{Cost}(3,8) = \min \{ c(8,10) + \text{Cost}(4,10),$$

$$c(8,11) + \text{Cost}(4,11) \}$$

$$= \{ 5+2, 6+5 \}$$

$$\text{Cost}(3,8) = \{ 7, 11 \}$$

$$\text{Cost}(3,8) = 7$$

$$d(3,8) = 10$$

$$\text{Cost}(2,2) = \min \{ c(2,6) + \text{Cost}(3,6),$$

$$c(2,7) + \text{Cost}(3,7),$$

$$c(2,8) + \text{Cost}(3,8) \}$$

$$= \min \{ 4+7, 2+5, 1+7 \}$$

$$= \min \{ 11, 7, 8 \}$$

$$\text{Cost}(2,2) = 7$$

$$d(2,2) = 7$$

Date _____
Page _____

$$\text{Cost}(2,3) = \min \{ c(3,6) + \text{Cost}(3,6), \\ c(3,7) + \text{Cost}(3,7) \}$$

$$= \min \{ 2+7+7+5 \}$$

$$\text{Cost}(2,3) = \min \{ 9, 9 \} = 9$$

$$d(2,3) = 6$$

$$\text{Cost}(2,4) = \min \{ c(4,8) + \text{Cost}(3,8) \}$$

$$= \min \{ 11+7 \} = \min \{ 18 \} = 18$$

$$\text{Cost}(2,5) = \min \{ c(5,7) + \text{Cost}(3,7), \\ c(5,8) + \text{Cost}(3,8) \}$$

$$= \min \{ 11+5, 8+7 \}$$

$$\text{Cost}(2,5) = \min \{ 16, 15 \} = 15$$

$$d(2,5) = 8$$

$$\text{Cost}(1,1) = \min \{ c(1,2) + \text{Cost}(2,2),$$

$$c(1,3) + \text{Cost}(2,3),$$

$$c(1,4) + \text{Cost}(2,4),$$

$$c(1,5) + \text{Cost}(2,5) \}$$

$$= \min \{ 9 + 7, 7 + 9, 3 + 18, 2 + 15 \}$$

$$= \min \{ 16, 16, 21, 17 \}$$

$$\text{Cost}(1, \dots) = 16$$

$$d(1, 1) = 2$$

Shortest path


$$d(1, 1) = 2$$

$$d(2, 2) = 7$$

$$d(3, 7) = 10$$

$$d(4, 10) = 2$$

path



Shortest dist $= 9 + 2 + 3 + 2 = \underline{\underline{16}}$
from s to t

Q.4) Differentiate b/w


Dynamic Programming	Greedy Strategy
1) Dynamic Programming is used to obtain the optimal solution.	1.) Greedy Method is also used to get the optimal solution.
2) Breaks down the problem into subproblems.	2.) Makes choices without considering the future.
3) Less efficient as compared to a greedy approach.	3) More efficient as compare to a greedy approach.
4) It is guaranteed that Dynamic Programming will generate an optimal sol'n using Principle of optimality.	4) In Greedy Method, there is no such guarantee of getting optimal solution.
5) Memorization is required.	5) No memorization is required.
6) Slow results comparatively.	6) Fast results.
7) Every problem can be solved by dynamic algorithm.	7) Every problem can't be solved by a greedy algorithm.

Date _____
Page _____


- | | |
|--------------------------|--|
| 8) LCS, knapsack problem | 8) Shortest path proble
m, Huffman Coding |
|--------------------------|--|

Q.5] Travelling Salesman Problem ? Implement
TSP for Following matrix multiplication

	0	10	15	20
2	5	0	9	10
3	6	13	0	12
4	8	8	9	0



Cost :-



$$g(i, \emptyset) = \min_{j \in S} \{ c_{ij} + g(j, \emptyset - \{j\}) \}$$

Let the starting vertex 1

* If $|S| = \phi$. \rightarrow Consider null set element

$$g(2, \emptyset) = C_{21} = 5$$

$$g(3, \emptyset) = C_{31} = 6$$

$$g(4, \emptyset) = C_{41} = 8$$

Page No. _____

Date / /

* If $|s| = 1 \rightarrow$ Consider 1 set element

$$g(2, \{3\}) = C_{23} + g(3, \emptyset) = 9 + 6 = 15$$

$$g(2, \{4\}) = C_{24} + g(4, \emptyset) = 10 + 8 = 18$$

$$g(3, \{2\}) = C_{32} + g(2, \emptyset) = 13 + 5 = 18$$

$$g(3, \{4\}) = C_{34} + g(4, \emptyset) = 12 + 8 = 20$$

$$g(4, \{3\}) = C_{43} + g(3, \emptyset) = 9 + 6 = 15$$

$$g(4, \{2\}) = C_{42} + g(2, \emptyset) = 8 + 5 = 13$$

* If $|s| = 2 \rightarrow$ Consider two set element.

$$g(2, \{3, 4\}) = \min \{ C_{23} + g(3, \{4\}), C_{24} + g(4, \{3\}) \}$$

$$= \min \{ 9 + 20, 10 + 15 \}$$

$$= \min \{ 29, 25 \}$$

$$g(2, \{3, 4\}) = \underline{\underline{25}}$$

$$g(2, \{4, 3\}) = \min \left\{ c(2, 4) + g(4, \{3\}), c(2, 3) + g(3, \{4\}) \right\}$$

$$= \min \left\{ 10 + 15, 12 + 20 \right\}$$

$$= \min \left\{ 25, 22 \right\}$$

$$g(2, \{4, 3\}) = 25$$

$$g(3, \{2, 4\}) = \min \left\{ c(3, 2) + g(2, \{4\}), c(3, 4) + g(4, \{2\}) \right\}$$

$$= \min \left\{ 13 + 18, 12 + 13 \right\}$$

$$= \min \left\{ 31, 25 \right\}$$

$$g(3, \{2, 4\}) = 25$$

$$g(3, \{4, 2\}) = \min \left\{ c(3, 4) + g(4, \{2\}), c(3, 2) + g(2, \{4\}) \right\}$$

$$= \min \left\{ 12 + 18, 18 + 12 \right\}$$

$$= \min \{ 30, 30 \}$$

$$g(3, \{4, 2\}) = 25$$

$$g(4, \{2, 3\}) = \min \{ C_{42} + g(2, \{3\}), \\ C_{43} + g(3, \{2\}) \} \\ = \min \{ 8 + 15, 9 + 20 \} \\ = \min \{ 23, 29 \} \\ g(4, \{2, 3\}) = \underline{\underline{23}}$$

$$g(4, \{3, 2\}) = \min \{ C_{43} + g(3, \{2\}), \\ C_{42} + g(2, \{3\}) \} \\ = \min \{ 9 + 18, 8 + 15 \} \\ = \min \{ 27, 23 \} \\ g(4, \{3, 2\}) = \underline{\underline{23}}$$

* If $|S| = 3$.

$$g(1, \{2, 3, 4\}) = \min \{ C_{12} + g(2, \{3, 4\}), \\ C_{13} + g(3, \{2, 4\}), \\ C_{14} + g(4, \{2, 3\}) \}$$

$$= \min \{ 10 + 25, 15 + 25, 20 + 23 \} \\ = \min \{ 35, 40, 43 \}$$

$$g(1, \{2, 3, 4\}) = \underline{\underline{35}}$$


path : $1 \xrightarrow{10} 2 \xrightarrow{10} 4 \xrightarrow{9} 3 \xrightarrow{6} 1$

And cost is 35

Q.10) Implement T. Salesman problem for following matrix.

	1	2	3	4
1	0	15	10	20
2	5	0	10	9
3	6	13	0	12
4	8	8	9	0

Soln :-



$$g(i, s) = \min \{ c_{ij} + g(j, s - \{j\}) \}$$

* If $|s| = \emptyset$

$$g(2, \emptyset) = C_{2,1} = 5$$

$$g(3, \emptyset) = C_{3,1} = 6$$

$$g(4, \emptyset) = C_{4,1} = 8$$

Page No. _____

Date / /

$$g_f(1\varphi) = 1$$

$$g(2, \{3\}) = \min \{ c_{23} + g(3, \varphi) \} = 10 + 6 = 16$$

$$g(2, \{4\}) = \min \{ c_{24} + g(4, \varphi) \} = 9 + 8 = 17$$

$$g(3, \{2\}) = \min \{ c_{32} + g(2, \varphi) \} = 13 + 5 = 18$$

$$g(3, \{4\}) = \min \{ c_{34} + g(4, \varphi) \} = 12 + 8 = 20$$

$$g(4, \{2\}) = \min \{ c_{42} + g(2, \varphi) \} = 8 + 5 = 13$$

$$g(4, \{3\}) = \min \{ c_{43} + g(3, \varphi) \} = 9 + 6 = 15$$

$$g_f(1\varphi) = 2$$

$$g(2, \{3, 4\}) = \min \{ c_{23} + g(3, \{4\}), \\ c_{24} + g(4, \{3\}) \}$$

$$= \min \{ 10 + 20, 9 + 15 \} \\ = \min \{ 30, 24 \}$$

$$g(2, \{3, 4\}) = 24$$

$$g(2, \{4, 3\}) = \min \{ c_{24} + g(4, \{3\}), \\ c_{23} + g(3, \{4\}) \}$$

$$= \min \{ 9 + 15, 10 + 20 \} \\ = \min \{ 24, 30 \}$$

Shrikrupa $g(2, \{4, 3\}) = 24$

$$g(4, \varphi) = c_{41} = 2$$

$$g(3, \{2, 4\}) = \min \{ c_{32} + g(2, \{4\}), \\ c_{34} + g(4, \{2\}) \}$$

$$= \min \{ 13 + 17, 12 + 13 \}$$

$$g(3, \{2, 4\}) = 25$$

$$g(3, \{4, 2\}) = \min \{ c_{34} + g(4, \{2\}), \\ c_{32} + g(2, \{4\}) \}$$

$$= \min \{ 12 + 13, 13 + 17 \}$$

$$g(3, \{4, 2\}) = \min \{ 25, 30 \}$$

$$g(4, \{2, 3\}) = \min \{ c_{42} + g(2, \{3\}), \\ c_{43} + g(3, \{2\}) \}$$

$$= \min \{ 8 + 16, 9 + 18 \}$$

$$= \min \{ 24, 27 \}$$

$$g(4, \{2, 3\}) = 24$$

$$\begin{aligned}
 g(4, \{3, 2\}) &= \min \{ C_{43} + g(3, \{2\}), \\
 &\quad C_{42} + g(2, \{3\}) \} \\
 &= \min \{ 9 + 18, 8 + 16 \} \\
 &= \min \{ 27, 24 \} \\
 g(4, \{3, 2\}) &= 24
 \end{aligned}$$

$$\begin{aligned}
 g(4, \{2, 3\}) &= \min \{ C_{42} + g(2, \{3\}), \\
 &\quad C_{43} + g(3, \{2\}) \} \\
 &= \min \{ 8 + 16, 9 + 18 \} \\
 &= \min \{ 24, 27 \} \\
 g(4, \{2, 3\}) &= 24
 \end{aligned}$$

* If $|S| = 3$

$$\begin{aligned}
 g(1, \{2, 3, 4\}) &= \min \{ C_{12} + g(2, \{3, 4\}), \\
 &\quad C_{13} + g(3, \{2, 4\}), \\
 &\quad C_{14} + g(4, \{2, 3\}) \} \\
 &= \min \{ 15 + 24, 10 + 25, 20 + 24 \} \\
 &= \min \{ 39, 35, 44 \} \\
 g(1, \{2, 3, 4\}) &= 35
 \end{aligned}$$


path is $1 \rightarrow 3 \rightarrow 4 \rightarrow 2 \rightarrow 1$




Date _____
Page _____

Q. 11). Travelling Salesman Problem ?

	1	2	3	4
1	0	9	8	7
2	10	0	6	5
3	12	13	0	8
4	2	3	5	0



Q. 11 :-



$$g(i, \emptyset) = \min_{j \in S} \{ c_{ij} + g(j, \emptyset - \{j\}) \}$$

Let the starting vertex 1

If $|S| = \phi \rightarrow$ Consider no set element.

$$g(2, \emptyset) = C_{21} = 10$$

$$g(3, \emptyset) = C_{31} = 12$$

$$g(4, \emptyset) = C_{41} = 2$$

$$g(1, \emptyset) = 1$$

$$g(2, \{3\}) = \min \{ C_{23} + g(3, \emptyset) \} \min \\ = 6 + 12 = 18$$

$$g(2, \{4\}) = \min \{ C_{24} + g(4, \emptyset) \} \\ = 5 + 2 = 7$$

$$g(3, \{2\}) = \min \{ C_{32} + g(2, \emptyset) \} \\ = 13 + 10 = 23$$

$$g(3, \{4\}) = \min \{ C_{34} + g(4, \emptyset) \} \\ = 8 + 2 = 10$$

$$g(4, \{3\}) = \min \{ C_{43} + g(3, \emptyset) \} \\ = 5 + 12 = 17$$

$$g(4, \{2\}) = \min \{ C_{42} + g(2, \emptyset) \} \\ = 9 + 10 = 19$$

Microsoft
Includes:
Office
Home & Student
Don't forget to activate

RYZEN
5000 SERIES

NVIDIA
GEFORCE
GTX

144Hz
REFRESH RATE



Date _____
Page _____

$$\text{If } |S| = 2$$

$$g(2, \{3, 4\}) = \min \{ C_{23} + g(3, \{4\}), \\ C_{24} + g(4, \{3\}) \} \\ = \min \{ 6 + 10, 5 + 17 \} \\ = \min \{ 16, 22 \} = 16$$

$$g(2, \{4, 3\}) = \min \{ C_{24} + g(4, \{3\}), \\ C_{23} + g(3, \{4\}) \} \\ = \min \{ 5 + 17, 6 + 10 \} \\ = \min \{ 22, 16 \} = 16$$

$$g(3, \{2, 4\}) = \min \{ C_{32} + g(2, \{4\}), \\ C_{34} + g(4, \{2\}) \} \\ = \min \{ 13 + 7, 8 + 13 \} \\ = \min \{ 20, 21 \} = 20$$

$$g(3, \{4, 2\}) = \min \{ C_{34} + g(4, \{2\}), \\ C_{32} + g(2, \{4\}) \} \\ = \min \{ 8 + 13, 13 + 7 \} \\ = \min \{ 21, 20 \} = 20$$

$$g(4, \{2, 3, 4\}) = \min \{ C_{42} + g(\{2, \{3, 4\}\}), \\ C_{43} + g(\{3, \{2, 4\}\}), \\ = \min \{ 3 + 18, 5 + 23 \}, \\ = \min \{ 21, 28 \} = 21.$$

$$g(4, \{3, 2, 4\}) = \min \{ C_{43} + g(\{3, \{2, 4\}\}), \\ C_{42} + g(\{2, \{3, 4\}\}), \\ = \min \{ 5 + 23, 3 + 18 \}, \\ = \min \{ 28, 21 \} = 21$$

* If $|S| = 3$

$$g(1, \{2, 3, 4\}) = \min \{ C_{12} + g(\{2, \{3, 4\}\}), \\ C_{13} + g(\{3, \{2, 4\}\}), \\ C_{14} + g(\{4, \{2, 3\}\}), \\ = \min \{ 9 + 16, 8 + 20, 7 + 21 \}, \\ = \min \{ 25, 28, 28 \} \\ = 25$$

path: $1 \xrightarrow{7} 4 \xrightarrow{3} 2 \xrightarrow{6} 3 \xrightarrow{12} 1$


$$\text{Cost} \Rightarrow 7 + 3 + 6 + 12 = 28$$

Date
Page

12)

	1	2	3	4
1	0	8	16	15
2	14	0	9	12
3	7	10	0	6
4	11	13	10	0

Soln:-



$$g(i, \{s\}) = \min_{j \in s} \{ c_{ij} + g(j, s - \{j\}) \}$$

* If $|s| = \phi$

$$g(2, \phi) = C_{21} = 14$$

$$g(3, \phi) = C_{31} = 7$$

$$g(4, \phi) = C_{41} = 11$$

* If $|s| = 1$

$$g(2, \{3\}) = \min \{ C_{23} + g(3, \phi) \}$$

$$= \min \{ \dots \}$$

$$g(2, \{3, 4\}) = \min \{ C_{24} + g(4, \emptyset) \} = 12$$

$$g(3, \{2\}) = \min \{ C_{32} + g(2, \emptyset) \} = 10$$

$$g(3, \{4\}) = \min \{ C_{34} + g(4, \emptyset) \} = 6$$

$$g(4, \{2\}) = \min \{ C_{42} + g(2, \emptyset) \} = 13$$

$$g(4, \{3\}) = \min \{ C_{43} + g(3, \emptyset) \} = 10$$

$$* \quad \text{If } |S| = 2$$

$$g(2, \{3, 4\}) = \min \{ C_{23} + g(3, \{4\}),$$

$$C_{24} + g(4, \{3\}) \}$$

$$= \min \{ 9 + 6, 12 + 10 \}$$

$$g(2, \{3, 4\}) = \min \{ 15, 22 \} = 15$$

$$g(3, \{2, 4\}) = \min \{ C_{32} + g(2, \{4\}),$$

$$C_{34} + g(4, \{2\}) \}$$

$$= \min \{ 10 + 12, 6 + 13 \}$$

$$= \min \{ 22, 19 \} = 19$$

Date _____
Page _____

$$g(3, \{4, 2\}) = \min \{ C_{34} + g(4, \{2\}), \\ C_{32} + g(2, \{4\}) \}$$

$$= \min \{ 6 + 13, 10 + 12 \}$$

$$g(3, \{4, 2\}) = \min \{ 19, 22 \} = 19$$

$$g(2, \{4, 3\}) = \min \{ C_{24} + g(4, \{3\}), \\ C_{23} + g(3, \{4\}) \}$$

$$= \min \{ 12 + 10, 9 + 6 \}$$

$$g(2, \{4, 3\}) = \min \{ 22, 15 \} = 15$$

$$g(4, \{2, 3\}) = \min \{ C_{42} + g(2, \{3\}), \\ C_{43} + g(3, \{2\}) \}$$

$$= \min \{ 13 + 9, 10 + 10 \}$$

$$g(4, \{2, 3\}) = \min \{ 22, 20 \} = 20$$

$$g(4, \{3, 2\}) = \min \{ C_{43} + g(3, \{2\}), \\ C_{42} + g(2, \{3\}) \}$$

$$= \min \{ 10 + 10, 13 + 9 \}$$

$$(4, \{3, 2\}) = \min \{ 20, 22 \} = 20.$$

$$* |f|_g = g$$

$$g(1, \{2, 3, 4\}) = \min \{ C_{12} + g(2, \{3, 4\}),$$

$$C_{13} + g(3, \{2, 4\}),$$

$$C_{14} + g(4, \{2, 3\}) \}$$

$$= \min \{ 8 + 15, 16 + 10, 15 + 20 \}$$

$$= \min \{ 23, 35, 35 \}$$

$$g(1, \{2, 3, 4\}) = 23$$

path i5.

$$1 - 2 - 3 - 4 - 1$$

$$\text{path cost i5} = 8 + 9 + 6 + 11 \\ = 34$$

Q. 16) Determine LCS; $X = \{A, G, T, A, B, Z\}$,
 $Y = \{G, X, T, X, A, Y, B\}$.

	X \ Y	G	X	T	A	Y	B
X	0	0	0	0	0	0	0
G	0	↑0	↖0	↖0	↖0	↑1	↖1
G	0	↑1	↖1	↖1	↖1	↖1	↖1
T	0	↑1	↖1	↖1	↖1	↖1	↖1
A	0	↑1	↖1	↖2	↖2	↖3	↖3
B	0	↑1	↖1	↖2	↖2	↖3	↖4
Z	0	↑1	↖1	↖2	↖2	↑3	↖3

We get the maximum length of common subsequence as 4.

So, Final string of
LCS \Rightarrow G T A B

Page No. _____

Date 1/1

Q.29)

Largest Common Subsequence.
Find optimal solution for given sequence.

$$X = (A, B, C, D, B, A, B) \quad \& \quad Y = (B, C, D, A, B, A)$$

$$X = (A, B, C, D, B, A, B)$$

$$Y = (B, C, D, A, B, A)$$

X	Y	B	C	D	A	B	A
	0	0	0	0	0	0	0
A	0	←0	←0	←0	↑1	←01	↑1
B	0	↑1	←1	←1	←1	↑2	←2
C	0	↑1	↖2	↖2	↖2	↖2	↖2
D	0	↑1	↑2	↖3	↖3	↖3	↖3
B	0	↑1	↑2	↑3	↖3	↖4	↖4
A	0	↑1	↑2	↑3	↖4	↖4	↖5
B	0	↖1	↖2	↖3	↖4	↖5	↖5

We get the maximum length of common subsequence as 5.

So, Final string of longest common subsequence. $\Rightarrow B C D A B$

Page No. _____

Date 1/1


Longest Common Subsequence.

X = a a b a a b a b a a
Y = b a b a a b a b

		b	a	b	a	a	a	b	a	b
		0	0	0	0	0	0	0	0	0
a	0	↖0	↑1	↖1	↖1	↖1	↖1	↖1	↖1	↖1
	0	↖0	↑1	↖1	↖2	↖2	↖2	↖2	↖2	↖2
b	0	↑1	↖1	↖2	↖2	↖2	↖3	↖3	↖3	↖3
	0	↑1	↖2	↖2	↖3	↖3	↖3	↖4	↖4	↖4
a	0	↑1	↑2	↖2	↖3	↖3	↖4	↖4	↖4	↖4
	0	↑1	↑2	↖2	↖3	↖4	↖4	↖4	↖4	↖4
b	0	↑1	↑2	↖3	↖3	↖4	↖4	↖5	↖5	↖5
	0	↑1	↑2	↑3	↑4	↖4	↖4	↖5	↖6	↖6
a	0	↑1	↑2	↑3	↖4	↖4	↖5	↖5	↖6	↖6
	0	↑1	↑2	↑3	↑4	↖4	↖5	↖6	↖7	↖7
a	0	↑1	↑2	↑3	↖4	↖5	↖5	↖6	↖6	↑7
	0	↑1	↑2	↑3	↖4	↖5	↖5	↖6	↖6	↑7

We get the maximum length of common subsequence as 7


So, Final string of \Rightarrow abaabab



Date _____
Page _____

Q.25)

Floyd Warshall.



Sol :-

1 2 3 4

	A	B	C	D
1 A	0	4	∞	5
2 B	∞	0	1	2
3 C	3	∞	0	∞
4 D	∞	∞	6	0

	1	2	3	4
1 A	0	4	∞	5
2 B	∞	0	1	2
3 C	3	7	0	8
4 D	∞	∞	6	0


	1	2	3	4
1	0	4	5	5
2	∞	0	1	2
3	3	7	0	8
4	∞	∞	6	0

$$D^3 = \begin{array}{c|cccc} & 1 & 2 & 3 & 4 \\ \hline 1 & 0 & 4 & 5 & 5 \\ 2 & 4 & 0 & 1 & 2 \\ 3 & 3 & 7 & 0 & 8 \\ 4 & 9 & 13 & 6 & 0 \end{array} \rightarrow$$
$$D^4 = \begin{array}{c|cccc} & A & B & C & D \\ \hline A & 0 & 4 & 5 & 5 \\ B & 4 & 0 & 1 & 2 \\ C & 3 & 7 & 0 & 8 \\ D & 9 & 13 & 6 & 0 \end{array}$$

→ D^4 gives the shortest path b/w each pair of vertices.

Q. 26

Find all shortest paths using
Floyd Warshall algorithm

Solutiⁿ:

	A	B	C	D
1 A	0	2	1	3
2 B	∞	0	4	∞
3 C	∞	∞	0	∞
4 D	∞	∞	8	0

	A	B	C	D
1 A	0	2	1	3
2 B	∞	0	4	∞
3 C	∞	∞	0	∞
4 D	∞	∞	8	0

	A	B	C	D
1 A	0	2	1	3
2 B	∞	0	4	∞
3 C	∞	∞	0	∞
4 D	∞	∞	6	0

Page No. _____

Date 7/1

$$D^3 = \begin{array}{c|ccccc} & A & B & C & D \\ \hline A & 0 & 2 & 1 & 3 \\ B & \infty & 0 & 4 & \infty \\ C & \infty & \infty & 0 & \infty \\ D & \infty & \infty & 6 & 0 \end{array}$$
$$D^4 = \begin{array}{c|ccccc} & A & B & C & D \\ \hline A & 0 & 2 & 1 & 3 \\ B & \infty & 0 & 4 & \infty \\ C & \infty & \infty & 0 & \infty \\ D & \infty & \infty & 6 & 0 \end{array}$$

→ D^4 gives the shortest path between each pair of vertices.