

# Intro to Modern Algebra: Homework #2

Due on November 2 at 9:30am

*Professor Lorenz 9:30-11:00*

Sam Cook

## Problem 6

Which of the following subsets of  $\mathbb{R}[x]$  are subrings of  $\mathbb{R}[x]$ .

- a All polynomials with constant term of  $0_R$
- b All polynomials of degree 2.

### Solution

a) This is a subring. We will show that it is a nonempty subset closed under subtraction and multiplication. First, zero is a polynomial with the constant term  $0_R$ , so it is a nonempty subset. Next, consider polynomials  $a, b \in \mathbb{R}[x]$ . Then if we subtract them and  $0_R$ , we get  $(a + 0_R) - (b + 0_R) = (a - b) + (0_R - 0_R) = a - b$ , and since  $a, b \in R$ , it is closed under subtraction. Similarly for multiplication, we get  $(a + 0_R) * (b + 0_R) = (ab) + (0_R * b) + (0_R * a) + 0_R = ab$ , so it is similarly closed under multiplication. Therefore, it is a subring of  $\mathbb{R}[x]$ .

b) This is not a subring because it is not closed under multiplication. Consider the polynomials  $x^2$  and  $x^2 + 1$ . When we multiply them,  $x^2 * (x^2 + 1) = x^4 + x^2$ , which is not a degree 2 polynomial, and therefore it is not closed under multiplication and cannot be a ring. Furthermore,  $0_R$  is not of degree 2 and therefore is not in the subset.

## Problem 20

Let  $D : \mathbb{R}[x] \rightarrow \mathbb{R}[x]$  be the derivative map defined by

$$D(a_0 + a_1x + a_2x^2 + \dots + a_nx^n) = a_1 + 2a_2x + 3a_3x^2 + \dots + na_nx^{n-1}$$

Is  $D$  a homomorphism of rings? An isomorphism?

### Solution

This is neither a homomorphism of rings nor an isomorphism. We will prove this by showing  $D$  does not preserve multiplication. Consider  $x^2, x^3 \in \mathbb{R}[x]$ . When we multiply them,  $x^2 * x^3 = x^5$ . Now, we will apply  $D$  and multiply them.  $D(x^2) * D(x^3) = 2x * 3x^2 = 6x^3 \neq x^5$ . Therefore,  $D$  does not preserve multiplication and cannot be a homomorphism or isomorphism.

## Problem 13

Prove Theorem 4.10.

### Theorem 4.10

Let  $\mathbb{F}$  be a field and  $a(x), b(x), c(x) \in \mathbb{F}[x]$ . If  $a(x) | b(x)c(x)$  and  $a(x)$  and  $b(x)$  are relatively prime, then  $a(x) | c(x)$ .

### Solution

Assume  $\mathbb{F}$  be a field and  $a(x), b(x), c(x) \in \mathbb{F}[x]$ ,  $a(x) | b(x)c(x)$ , and  $a(x)$  and  $b(x)$  are relatively prime. Then  $\gcd(a(x), b(x)) = 1$ , by the definition of relatively prime. But then, by Theorem 4.8, this means that for polynomials  $u(x), v(x) \in \mathbb{F}[x]$ ,

$$1 = a(x)v(x) + b(x)u(x).$$

We can multiply this equation by  $c(x)$  to get that

$$c(x)a(x)v(x) + c(x)b(x)u(x) = c(x)$$

Since  $a(x)|b(x)c(x)$ , there exists some  $z(x) \in (F)[x]$  such that  $a(x)z(x) = b(x)c(x)$ . We can substitute this into the above equation to get

$$c(x)a(x)v(x) + a(x)z(x)u(x) = c(x)$$

and we can then factor out  $a(x)$  to get

$$a(x)[c(x)v(x) + z(x)u(x)] = c(x)$$

Since  $c(x), v(x), z(x), u(x) \in \mathbb{F}[x]$ ,  $a(x)|c(x)$ .

## Problem 12

Express  $x^4 - 4$  as a product of irreducibles in  $\mathbb{Q}[x]$ ,  $\mathbb{R}[x]$ , and in  $\mathbb{C}[x]$ .

### Solution

First, we will factor this into a product of irreducibles in  $\mathbb{C}[x]$ . In  $\mathbb{C}[x]$ ,

$$x^4 - 4 = (x - i\sqrt{2})(x + i\sqrt{2})(x - 2)(x + 2)$$

These factors are all irreducible because they are all degree 1, both complex numbers and irrational numbers are in  $\mathbb{C}$

In  $\mathbb{R}[x]$ ,

$$x^4 - 4 = (x^2 + 2)(x + \sqrt{2})(x - \sqrt{2})$$

In this case,  $(x + \sqrt{2})$  and  $(x - \sqrt{2})$  are factors because they are degree 1 with coefficients in the real numbers.  $(x + \sqrt{2})$  is only irreducible in  $\mathbb{C}[x]$  because it has complex coefficients.

In  $\mathbb{Q}[x]$ ,

$$x^4 - 4 = (x^2 + 2)(x^2 - 2)$$

The first factor is irreducible because if it were not irreducible, it would be reducible in  $\mathbb{R}[x]$ . The second factor is irreducible because it only factors into polynomials with irrational coefficients, which, by definition, are not in the  $\mathbb{Q}$

## Problem 4

a For what value of  $k$  is  $x - 2$  a factor of  $x^4 - 5x^3 + 5x^2 + 3x + k$  in  $\mathbb{Q}[x]$

b For what value of  $k$  is  $x + 1$  a factor of  $x^4 + 2x^3 - 3x^2 + kx + 1$  in  $\mathbb{Z}_5[x]$

### Solution

a) For  $k = -2$  is  $(x - 2)$  a factor of  $x^4 - 5x^3 + 5x^2 + 3x + k$ . This is simply to solve for because  $(x - 2)$  is in the form of a root, so we can evaluate our polynomial at  $x = 2$  and solve for  $k$ .

$$2^4 - 5(2)^3 + 5(2)^2 + 3(2) - k = 0 \Rightarrow 16 - 40 + 20 + 6 = -k \Rightarrow -2 = k$$

b) For  $k = 3$  is  $x + 1$  a factor of  $x^4 + 2x^3 - 3x^2 + kx + 1$  in  $\mathbb{Z}_5[x]$ . Like in part a, since  $(x + 1)$  is in the form of a root, we can evaluate our polynomial at  $x = -1$  and solve for  $k$ .

$$(-1)^4 + 2(-1)^3 - 3(-1)^2 + k(-1) + 1 = 0 \Rightarrow 1 - 2 - 3 - k + 1 = 0 \Rightarrow 2 = k + 5 \Rightarrow k = 3$$

## Problem 5

Show that  $x - 1_F$  divides  $a_n x^n + \dots + a_2 x^2 + a_1 x + a_0$  in  $\mathbb{F}[x]$  if and only if  $a_0 + \dots + a_n = 0_F$ .

**Solution**

$\Rightarrow$  Assume that  $x - 1_F$  divides  $a_n x^n + \dots + a_2 x^2 + a_1 x + a_0$  in  $\mathbb{F}[x]$ . Then  $1_F$  is a root of  $a_n x^n + \dots + a_2 x^2 + a_1 x + a_0$ , so this means that  $a_n 1^n + \dots + a_2 1^2 + a_1 + a_0 = 0$ . Since 1 to any power is 1, this simplifies to  $a_n + \dots + a_2 + a_1 + a_0 = 0$

$\Leftarrow$  This proof is simply the above proof in reverse. Assume that  $a_0 + \dots + a_n = 0_F$  in  $\mathbb{F}[x]$ . This can be rewritten as  $a_0 1 + \dots + a_n 1 = 0_F$ , or  $a_0 1^0 + \dots + a_n 1^n = 0_F$ . This is the form of a polynomial root evaluated at 1, so we can rewrite this as  $a_n x^n + \dots + a_2 x^2 + a_1 x + a_0$

## Problem 14

a

## Problem 18

## Problem 19