

The Uniform (Co-Irreducible) Dimension of Rings and Modules

Supervisor: Prof. Dr Tiberiu Dumitrescu

Student: Sorin Bogde

Original thesis: University of Bucharest, Faculty of Mathematics, 1999

English translation: November 20, 2025

Contents

0	Generalities	1
0.1	Semisimple modules	1
0.2	Noetherian (Artinian) modules and Noetherian (Artinian) rings . . .	2
0.3	Finite length modules	3
0.4	The Jacobson radical	5
0.5	Semisimple rings	6
1	Essential submodules	7
2	Injective Modules	13
2.1	Injective Module	13
2.2	Injective envelopes	20

Chapter 0

Generalities

0.1 Semisimple modules

Definition 0.1.1. A non-zero R -module S is called *simple* if its only submodules are 0 and S .

Proposition 0.1.2. Let S be an R -module. The following statements are equivalent:

1. S is simple;
2. for every non-zero element $x \in S$ we have $S = xR$;
3. $S \cong R/I$, where I is a maximal right ideal.

Lemma 0.1.3 (Schur). Let S and S' be simple R -modules and let $f: S \rightarrow S'$ be a homomorphism of R -modules. Then $f = 0$ or f is an isomorphism. In particular $\text{End}_R(S)$ is a division ring.

Definition 0.1.4. Let M be an R -module and let $(S_i)_{i \in I}$ be the family of all simple submodules of M . If $M = \sum_{i \in I} S_i$, then M is called *semisimple*.

Proposition 0.1.5. Let M be a semisimple R -module and N a submodule of M . Then there exists a subset $J \subseteq I$ such that

1. the family $(S_j)_{j \in J}$ is independent;
2. $M = N \oplus (\bigoplus_{j \in J} S_j)$.

Corollary 0.1.6. With the above notation, for the semisimple module M there exists $J \subseteq I$ such that the family $(S_j)_{j \in J}$ is independent and

$$M = \bigoplus_{i \in I} M_i.$$

Corollary 0.1.7. If M is a semisimple R -module and N a submodule of M , then both N and M/N are semisimple.

Corollary 0.1.8. A direct sum of semisimple modules is a semisimple module.

Theorem 0.1.9. *Let M be an R -module. The following statements are equivalent:*

1. M is semisimple;
2. M is isomorphic to a direct sum of simple modules;
3. every submodule of M is a direct summand of M ;
4. every short exact sequence

$$0 \longrightarrow M' \xrightarrow{f} M \xrightarrow{g} M'' \longrightarrow 0$$

splits.

Definition 0.1.10. The sum of all simple submodules of M is called the *socle* of M and is denoted by $\text{soc}(M)$. If M has no simple submodule we put $\text{soc}(M) = 0$.

Proposition 0.1.11. *Let M and N be R -modules and $f: M \rightarrow N$ a homomorphism. Then $f(\text{soc}(M)) \subseteq \text{soc}(N)$.*

Proposition 0.1.12. *Let M be an R -module and N a submodule of M . Then*

$$\text{soc}(N) = \text{soc}(M) \cap N.$$

Proposition 0.1.13. *If $M = \bigoplus_{i \in I} M_i$, then*

$$\text{soc}(M) = \bigoplus_{i \in I} \text{soc}(M_i).$$

Proposition 0.1.14. *Let R be a ring. Then the socle $\text{soc}(R_R)$ is a two-sided ideal of R .*

0.2 Noetherian (Artinian) modules and Noetherian (Artinian) rings

Definition 0.2.1. Let R be a ring and M a right R -module. We say that M satisfies the *maximal condition* (resp. the *minimal condition*) if every non-empty set of submodules of M , ordered by inclusion, has a maximal (resp. minimal) element.

We say that M satisfies the *ascending* (resp. *descending*) *chain condition* if every ascending chain of submodules of M

$$M_1 \subseteq M_2 \subseteq \cdots \subseteq M_i \subseteq \cdots$$

(resp. every descending chain

$$M_1 \supseteq M_2 \supseteq \cdots \supseteq M_i \supseteq \cdots$$

) is stationary, that is, there exists $n \geq 1$ such that $M_n = M_{n+1} = \cdots$.

Proposition 0.2.2. *Let M be an R -module. The following statements are equivalent:*

1. M satisfies the maximal (minimal) condition;
2. M satisfies the ascending (descending) chain condition.

Definition 0.2.3. An R -module M is called *noetherian* (resp. *artinian*) if it satisfies the maximal (resp. minimal) condition. The ring R is called right noetherian (resp. right artinian) if the right module R_R is noetherian (resp. artinian).

Example 0.2.4.

1. \mathbb{Z} is a noetherian ring but not artinian.
2. Every finite group is a noetherian and artinian \mathbb{Z} -module.
3. Every finite ring is noetherian and artinian.
4. The ring $\mathbb{Z}[X_1, X_2, \dots, X_n, \dots]$ is neither noetherian nor artinian:

$$(X_1) \subsetneq (X_1, X_2) \subsetneq \cdots \subsetneq (X_1, \dots, X_n) \subsetneq \cdots$$

$$(X_1) \supsetneq (X_1^2) \supsetneq \cdots \supsetneq (X_1^k) \supsetneq \cdots$$

5. The Prüfer p -group \mathbb{Z}_{p^∞} is an artinian but not noetherian \mathbb{Z} -module.

Proposition 0.2.5. Let N and P be submodules of M such that $M = N + P$. Then M is noetherian (artinian) if and only if N and P are noetherian (artinian).

Proposition 0.2.6. For an R -module M the following statements are equivalent:

1. M is noetherian;
2. every submodule of M is finitely generated.

Proposition 0.2.7. For an R -module M the following statements are equivalent:

1. M is artinian;
2. for every family $(X_i)_{i \in I}$ of submodules of M there exists a finite subset $J \subseteq I$ such that

$$\bigcap_{i \in I} X_i = \bigcap_{j \in J} X_j.$$

0.3 Finite length modules

Definition 0.3.1. Let M be a non-zero right R -module. A *composition series* or *Jordan–Hölder series* of M is a finite strictly ascending chain of submodules

$$0 = X_0 \subset X_1 \subset \cdots \subset X_n = M$$

such that X_{i+1}/X_i is a simple module for $0 \leq i \leq n-1$. The integer n is called the *length* of the series, and the modules X_{i+1}/X_i are called its *factors*.

Proposition 0.3.2. Let M be an R -module. The following statements are equivalent:

1. M has a composition series;
2. M is noetherian and artinian.

Proposition 0.3.3. *Let*

$$0 \longrightarrow M' \longrightarrow M \longrightarrow M'' \longrightarrow 0$$

be a short exact sequence of right R -modules. Then M has a composition series if and only if both M' and M'' have composition series.

If

$$0 = M_0 \subseteq M_1 \subseteq \cdots \subseteq M_n = M, \quad 0 = N_0 \subseteq N_1 \subseteq \cdots \subseteq N_p = M$$

are two composition series of M , we say that they are *equivalent* if $n = p$ and there exists a bijection $\sigma : \{0, \dots, n-1\} \rightarrow \{0, \dots, n-1\}$ such that

$$M_{i+1}/M_i \cong M_{\sigma(i)+1}/M_{\sigma(i)} \quad (0 \leq i \leq n-1).$$

Theorem 0.3.4 (Jordan–Hölder). *If an R -module M has two composition series*

$$0 = M_0 \subseteq M_1 \subseteq \cdots \subseteq M_n = M, \quad 0 = N_0 \subseteq N_1 \subseteq \cdots \subseteq N_p = M,$$

then these two series are equivalent.

Definition 0.3.5. An R -module M which admits a composition series is called a *module of finite length*. The length of its composition series is called the *length* of M and is denoted by $l(M)$. If M admits no composition series, we say that M has *infinite length* and we write $l(M) = \infty$.

Proposition 0.3.6. *Let*

$$0 \longrightarrow M' \longrightarrow M \longrightarrow M'' \longrightarrow 0$$

be a short exact sequence of R -modules of finite length. Then

$$l(M) = l(M') + l(M'').$$

Corollary 0.3.7. *Let M be an R -module of finite length and let N, L be submodules of M . Then:*

1. $l(M) = l(N) + l(M/N)$;
2. $l(N + L) + l(N \cap L) = l(N) + l(L)$.

Corollary 0.3.8. *Let M be an R -module of finite length and let M_1, M_2, \dots, M_n be submodules of M such that*

$$M = M_1 \oplus M_2 \oplus \cdots \oplus M_n.$$

Then

$$l(M) = \sum_{i=1}^n l(M_i).$$

0.4 The Jacobson radical

The Jacobson radical of a module

Definition 0.4.1. Let M be an R -module. The intersection of all maximal submodules of M is called the *Jacobson radical* of M and is denoted by $\text{Rad}(M)$. If M has no maximal submodules, we adopt the convention $\text{Rad}(M) = M$.

Remark 0.4.2. If M is a finitely generated R -module, then $\text{Rad}(M) \neq M$.

Proposition 0.4.3. Let M be an R -module. Then

$$\text{Rad}(M) = \bigcap_{\substack{f: M \rightarrow S \\ S \text{ simple}}} \ker(f) = \bigcap_{\substack{f: M \rightarrow X \\ X \text{ semisimple}}} \ker(f).$$

Proposition 0.4.4. Let $f : M \rightarrow N$ be a homomorphism of R -modules. Then $f(\text{Rad}(M)) \subseteq \text{Rad}(N)$. If, in addition, f is an epimorphism and $\ker(f) \subseteq \text{Rad}(M)$, then $f(\text{Rad}(M)) = \text{Rad}(N)$.

Corollary 0.4.5. For every R -module M we have $\text{Rad}(M/\text{Rad}(M)) = 0$.

Corollary 0.4.6. If M is a semisimple R -module, then $\text{Rad}(M) = 0$.

Corollary 0.4.7. If $M = \bigoplus_{i \in I} M_i$, then

$$\text{Rad}(M) = \bigoplus_{i \in I} \text{Rad}(M_i).$$

Proposition 0.4.8. Let M be an R -module with $\text{Rad}(M) \neq M$. Then

$$\text{Rad}(M) = \bigcap \{ L \leq M \mid L \text{ is a superfluous submodule} \}.$$

Proposition 0.4.9 (Nakayama's Lemma). Let M be a finitely generated R -module and N a submodule of M . If $N + \text{Rad}(M) = M$, then $N = M$. (In other words, $\text{Rad}(M)$ is the largest superfluous submodule of M .)

The Jacobson radical of a ring

Let R be a ring. We consider the left ideal $\text{Rad}({}_R R)$, the intersection of all maximal left ideals of R , and the right ideal $\text{Rad}(R_R)$, the intersection of all maximal right ideals of R .

Proposition 0.4.10.

1. $\text{Rad}(R_R)$ is a two-sided ideal.
2. $\text{Rad}(R_R) = \{ r \in R \mid 1 - ar \in U(R) \text{ for all } a \in R \}$.
3. $\text{Rad}(R_R) = \text{Rad}({}_R R)$.

Definition 0.4.11. The two-sided ideal $\text{Rad}(R_R) = \text{Rad}({}_R R)$ is called the *Jacobson radical* of the ring R and is denoted by $\text{Rad}(R)$.

Proposition 0.4.12.

1. If J is a left (resp. right or two-sided) ideal such that $1 - x$ is invertible for every $x \in J$, then $J \subseteq \text{Rad}(R)$.
2. If J is a nil left (resp. right or two-sided) ideal, then $J \subseteq \text{Rad}(R)$.

Proposition 0.4.13. Let $\varphi : R \rightarrow S$ be a surjective ring homomorphism. Then $\varphi(\text{Rad}(R)) \subseteq \text{Rad}(S)$. If $\ker(\varphi) \subseteq \text{Rad}(R)$, then $\varphi(\text{Rad}(R)) = \text{Rad}(S)$.

Proposition 0.4.14. If $(R_i)_{i \in I}$ is a family of rings, then

$$\text{Rad}\left(\prod_{i \in I} R_i\right) = \prod_{i \in I} \text{Rad}(R_i).$$

Proposition 0.4.15. Let M be an R -module. Then $M \text{Rad}(R) \subseteq \text{Rad}(M)$.

Theorem 0.4.16. If R is an artinian ring, then $\text{Rad}(R)$ is nilpotent.

0.5 Semisimple rings

Theorem 0.5.1. For a ring R the following statements are equivalent:

1. every non-zero right R -module is semisimple;
2. R as a right R -module is semisimple;
3. R is artinian and $\text{Rad}(R) = 0$.

Definition 0.5.2. A ring R satisfying any (hence all) of the above conditions is called a *semisimple ring*.

Proposition 0.5.3. Let R be a semisimple ring and M a non-zero R -module. The following statements are equivalent:

1. M has finite length;
2. M is noetherian;
3. M is artinian.

Theorem 0.5.4. Let R be a right artinian ring and M a non-zero right R -module. The following statements are equivalent:

1. M has finite length;
2. M is noetherian;
3. M is artinian.

Corollary 0.5.5 (Hopkins). A right artinian ring (respectively a left artinian ring) is right noetherian (respectively left noetherian).

Chapter 1

Essential submodules

Definition 1.0.1. Let M be a right R -module. A submodule N of M is called *essential* (or we say that M is an essential extension of N) if $N \cap N' \neq 0$ for every non-zero submodule N' of M . In this case we write $N \trianglelefteq M_R$.

A homomorphism of right R -modules $f : M \rightarrow N$ is called *essential* if $\text{Im } f$ is an essential submodule of N (i.e. $\text{Im } f \trianglelefteq N_R$).

Example.

1. $n\mathbb{Z} \trianglelefteq \mathbb{Z}_{\mathbb{Z}}$ for every $n \geq 1$.
2. Every submodule of the Prüfer group \mathbb{Z}_{p^∞} is essential.

Remark 1.0.2. Let M be a right R -module and N a submodule of M . Then $N \trianglelefteq M_R$ if and only if for every $x \in M$, $x \neq 0$, there exists $r \in R$ such that $xr \in N \setminus \{0\}$.

Proof. “ \Rightarrow ” Let $x \in M \setminus \{0\}$. Since $0 \neq xR \leq M_R$ and $N \trianglelefteq M_R$, we have $xR \cap N \neq 0$; hence there exists $r \in R$ with $xr \in N \setminus \{0\}$.

“ \Leftarrow ” Let $N' \leq M_R$, $N' \neq 0$. For every $x \in N' \setminus \{0\}$ there exists $r \in R$ such that $xr \in N \setminus \{0\}$, so $N \cap N' \neq 0$. Thus $N \trianglelefteq M_R$. \square

Definition 1.0.3. A monomorphism of right R -modules $f : N_R \rightarrow M_R$ is called *essential* if $\text{Im } f \trianglelefteq M_R$. It is immediate that, if N is a submodule of M , then the canonical inclusion $i_N : N \rightarrow M$ is an essential monomorphism if and only if $N \trianglelefteq M_R$.

Proposition 1.0.4. A monomorphism $f : N_R \rightarrow M_R$ is essential if and only if for every right R -module M' and every $g \in \text{Hom}(M, M')$, the fact that $g \circ f$ is a monomorphism implies that g is a monomorphism.

Proof. “ \Rightarrow ” Let g be as in the statement and suppose that $g \circ f$ is a monomorphism. Assume $\text{Ker } g \neq 0$. Take $x \in \text{Ker } g \cap \text{Im } f \setminus \{0\}$. Then $x = f(x')$ for some $x' \in N$ and $g(x) = 0$, so $g(f(x')) = 0$. Since $g \circ f$ is a monomorphism, it follows that $x' = 0$ and hence $x = 0$, a contradiction.

“ \Leftarrow ” If f is not an essential monomorphism, then there exists $N' \leq M_R$, $N' \neq 0$, such that $N' \cap \text{Im } f = 0$. Consider the canonical projection $\pi_{N'} : M \rightarrow M/N'$. If $x \in \text{Ker}(\pi_{N'} \circ f)$, then $f(x) \in N'$, so $f(x) = 0$, hence $x = 0$. Thus $\pi_{N'} \circ f$ is injective. By the hypothesis in the statement, this implies that $\pi_{N'}$ is injective, so $N' = 0$, a contradiction. \square

Corollary 1.0.5. *Let M be a right R -module and $N \leq M_R$. Then the following statements are equivalent:*

1. $N \trianglelefteq M_R$;
2. the inclusion $i_N : N \rightarrow M$ is an essential monomorphism;
3. for every $f \in \text{Hom}(M, M')$, with M' an arbitrary R -module, the fact that $f \circ i_N$ is a monomorphism implies that f is a monomorphism.

Proposition 1.0.6. *Let $f : N_R \rightarrow M_R$ and $g : M_R \rightarrow P_R$ be monomorphisms. Then $g \circ f$ is essential if and only if both g and f are essential.*

Proof. “ \Leftarrow ” Let $z \in P \setminus \{0\}$. Since g is essential, there exists $r \in R$ such that $zr \in \text{Im } g \setminus \{0\}$. Thus there exists $y \in M \setminus \{0\}$ with $zr = g(y)$. As f is essential, there exists $r' \in R$ such that $yr' \in \text{Im } f \setminus \{0\}$. Hence there exists $x \in N \setminus \{0\}$ with $yr' = f(x)$. But $zr' = g(y)r' = g(yr') = g(f(x))$. If $zr' = 0$, then $g(f(x)) = 0$, hence $x = 0$, a contradiction. Therefore $zr' \in \text{Im}(g \circ f)$ and $zr' \neq 0$, which shows that $g \circ f$ is essential.

“ \Rightarrow ” Let $y \in M \setminus \{0\}$. Since g is a monomorphism, $g(y) \neq 0$. Thus there exists $r \in R$ such that $g(yr) \in \text{Im } g \setminus \{0\}$ and $g(yr) \neq 0$. Hence there exists $x \in N \setminus \{0\}$ such that $g(yr) = g(f(x))$, therefore $yr = f(x) \in \text{Im } f$, which shows that f is an essential monomorphism.

If $z \in P \setminus \{0\}$, there exists $r \in R$ such that $zr \in \text{Im}(g \circ f)$ and $zr \neq 0$. Since $\text{Im}(g \circ f) \subseteq \text{Im } g$, we obtain $zr \in \text{Im } g$, so g is essential. \square

Proposition 1.0.7. *Let M be a right R -module and L_1, L_2, \dots, L_n submodules of M . Then:*

- 1) $\bigcap_{i=1}^n L_i$ is essential in M iff each L_i is essential in M .
- 2) If $L_1 \subseteq L_2$ and L_1 is essential in M , then L_2 is essential in M .

The proof is obvious.

Proposition 1.0.8. *Let K and L be submodules of M .*

- 1) If $K \subseteq L \subseteq M$, then $K \trianglelefteq M$ iff $K \trianglelefteq L$ and $L \trianglelefteq M$.
- 2) If $h : K_R \rightarrow M_R$ is a module morphism and $L \trianglelefteq M$, then $h^{-1}(L) \trianglelefteq K$.
- 3) If $L_1, L_2 \leq M_R$ with $K_1 \trianglelefteq L_1$ and $K_2 \trianglelefteq L_2$, then $K_1 \cap K_2 \trianglelefteq L_1 \cap L_2$.

Proof.

1) Apply 1.5 and 1.6.

2) Let U be a non-zero submodule of K .

(i) If $h(U) = 0$, then $U \subseteq \ker h \subseteq h^{-1}(L)$, so $U \cap h^{-1}(L) \neq 0$.

(ii) If $h(U) \neq 0$, then $h(U) \cap L \neq 0$. Hence there exists $u \in U$ with $h(u) \in L$, $h(u) \neq 0$, so $u \in U \cap h^{-1}(L)$ and $u \neq 0$.

Thus $h^{-1}(L) \trianglelefteq K$.

3) Let $0 \neq X \leq L_1 \cap L_2$. Then $X \subseteq L_1$, so $0 \neq X \cap K_1 \leq L_1$. But $X \subseteq L_2$ implies $0 \neq (X \cap K_1) \cap L_2 = X \cap (K_1 \cap K_2)$. Hence $K_1 \cap K_2 \trianglelefteq L_1 \cap L_2$.

Proposition 1.0.9. *Let $(K_\lambda)_{\lambda \in \Lambda}$, $(L_\lambda)_{\lambda \in \Lambda}$ be families of submodules of M . If (K_λ) is independent in M and $K_\lambda \leq L_\lambda$ for all $\lambda \in \Lambda$, then (L_λ) is independent in M and*

$$\left(\bigoplus_{\lambda \in \Lambda} K_\lambda \right) \leq \left(\bigoplus_{\lambda \in \Lambda} L_\lambda \right).$$

Proof.

Let $K_1 \leq L_1$, $K_2 \leq L_2$ with $K_1 \cap K_2 = 0$. By 1.8(3), $0 \leq L_1 \cap L_2$, hence $L_1 \cap L_2 = 0$.

Let $\pi_1 : L_1 \oplus L_2 \rightarrow L_1$, $\pi_2 : L_1 \oplus L_2 \rightarrow L_2$ be the canonical projections. Since $K_1 \leq L_1$ and $K_2 \leq L_2$,

$$\pi_1^{-1}(K_1) = K_1 \oplus 0 \leq L_1 \oplus L_2,$$

and

$$\pi_2^{-1}(K_2) = 0 \oplus K_2 \leq L_1 \oplus L_2.$$

Hence

$$K_1 \oplus K_2 = \pi_1^{-1}(K_1) \cap \pi_2^{-1}(K_2) \leq L_1 \oplus L_2.$$

Induction gives the finite case. For the general case, let $0 \neq m \in \bigoplus_{\lambda \in \Lambda} L_\lambda$. Then m lies in a finite direct sum $\bigoplus_{\lambda \in \Lambda_0} L_\lambda$ for some finite $\Lambda_0 \subseteq \Lambda$. Since $(\bigoplus_{\lambda \in \Lambda_0} K_\lambda) \leq (\bigoplus_{\lambda \in \Lambda_0} L_\lambda)$, there exists $r \in R$ with $rm \in (\bigoplus_{\lambda \in \Lambda_0} K_\lambda) \setminus \{0\} \subseteq (\bigoplus_{\lambda \in \Lambda} K_\lambda)$. Thus

$$\left(\bigoplus_{\lambda \in \Lambda} K_\lambda \right) \leq \left(\bigoplus_{\lambda \in \Lambda} L_\lambda \right).$$

Proposition 1.0.10. *Let N be a submodule of M . Then there exists a submodule Q with $N \subseteq Q \subseteq M$ such that Q is a maximal essential extension of N inside M .*

Proof.

Let

$$\mathfrak{S} = \{L \leq M ; N \subseteq L \subseteq M, N \leq L\},$$

ordered by inclusion. Clearly $\mathfrak{S} \neq \emptyset$, since $N \in \mathfrak{S}$.

Let $(L_\lambda)_{\lambda \in \Lambda}$ be a totally ordered family of elements of \mathfrak{S} and put

$$L := \bigcup_{\lambda \in \Lambda} L_\lambda.$$

Clearly $L \leq M_R$.

Let $x \in L \setminus \{0\}$. Then there exists $\lambda_0 \in \Lambda$ with $x \in L_{\lambda_0}$. Since N is essential in L_{λ_0} , there exists $r \in R$ such that $xr \in N$ and $xr \neq 0$, hence L is an essential extension of N . Thus \mathfrak{S} is inductive and, by Zorn's lemma, \mathfrak{S} admits a maximal element Q which satisfies the required conditions.

Definition 1.0.11. Let M be a right R -module and $N \leq M_R$. A submodule $K \leq M_R$ is called a *complement* of N in M if K is a maximal submodule of M with the property that $K \cap N = 0$. A submodule $K \leq M_R$ is called a *complement submodule* of M if there exists $N \leq M_R$ such that K is a complement of N in M .

Remark 1.0.12. The set

$$\tilde{\mathfrak{S}} = \{ L \leq M_R \mid N \cap L = 0 \}$$

is inductive and, by applying Zorn's lemma, it follows that there exists a complement of N in M . In particular, 0 and M are complement submodules of M .

Proposition 1.0.13. *Let M_R , $N \leq M_R$ and $K \leq M_R$, with K a complement of N in M . There exists a complement Q of K in M such that $N \subseteq Q$. Moreover, Q is a maximal essential extension of N in M .*

Proof. It is easy to see that the set

$$\tilde{\mathfrak{S}} = \{ L \leq M_R \mid K \cap L = 0, N \subseteq L \}$$

is inductive, and Zorn's lemma guarantees the existence of Q .

Let L be a non-zero submodule of Q such that $L \cap N = 0$. Put $K_1 = L + K$. Clearly $K \subseteq K_1$. If $x \in N \cap (L + K)$, then $x = y + z$ with $y \in L$, $z \in K$. But $z = x - y \in Q$. Since $Q \cap K = 0$, we obtain $z = 0$ and hence $x = y$. From $L \cap N = 0$ it follows that $x = y = 0$, and therefore $N \cap (L + K) = 0$, which contradicts the fact that K is a complement of N in M . Thus $L \cap N \neq 0$ for every $0 \neq L \leq Q$, so Q is an essential extension of N .

Suppose that there exists $Q' \leq M_R$ with $N \leq Q'$ and $Q \subsetneq Q'$. Since Q' is a complement of K , we have $Q' \cap K \neq 0$. But $N \cap (Q' \cap K) = 0$ and $0 \neq Q' \cap K \leq Q'$, contradicting $N \leq Q'$. Hence Q is a maximal essential extension of N in M . \square

Definition 1.0.14. A submodule N of M_R is called *closed* if N has no proper (meaning different from N) essential extension in M .

Corollary 1.0.15. *Let M_R be a right R -module. The complement submodules of M coincide with the closed submodules of M .*

Proof. From 1.13 it follows immediately that every closed submodule of M is a complement submodule of M .

Conversely, let K be a complement submodule of M_R . Then there exists $N \leq M_R$ such that K is a complement of N in M . Assume that K has a proper essential extension in M ; that is, there exists $K' \leq M_R$ with $K \leq K'$ and $K \subsetneq K'$. By the maximality of K we have $K' \cap N \neq 0$, and since $K \leq K'$, it follows that

$$K \cap K' \cap N \neq 0,$$

contradiction. \square

Corollary 1.0.16. *Let N be a submodule of M_R . If K is a complement of N in M , then:*

1. $(N + K) \leq M_R$.
2. The canonical morphism $\pi_K \circ i_N : N \rightarrow M/K$ is an essential monomorphism.

Proof. (1) Let $x \in M \setminus \{0\}$. If $x \notin K$, then $K + Rx \neq K$ and hence $N \cap (K + Rx) \neq 0$. Let $y \in N \cap (K + Rx)$, $y \neq 0$. There exist $z \in K$ and $r \in R$ such that $y = z + rx$. If $rx = 0$, then $y = z$ and, since $N \cap K = 0$, we obtain $y = 0$, a contradiction. Thus $rx \neq 0$ and, because $rx = y - z$, we have $rx \in N + K$, which shows that $(N + K) \trianglelefteq M_R$.

(2) We have $\text{Im}(\pi_K \circ i_N) = (N + K)/K$. Let L/K be a non-zero submodule of M/K . Then

$$\frac{N + K}{K} \cap \frac{L}{K} = \frac{(N + K) \cap L}{K} = \frac{N \cap L + K}{K}.$$

Since K is a complement of N , we have $N \cap L \neq 0$, and hence

$$\frac{N \cap L + K}{K} \neq 0,$$

which shows that $\pi_K \circ i_N$ is an essential monomorphism. □

Chapter 2

Injective Modules

2.1 Injective Module

Let Q and M be two right R -modules. We say that Q is M -*injective* if for every monomorphism $u : M' \rightarrow M$ and every morphism $f : M' \rightarrow Q$, there exists $g : M \rightarrow Q$ such that $g \circ u = f$; that is, the diagram

$$\begin{array}{ccccc} 0 & \longrightarrow & M' & \xrightarrow{u} & M \\ & & \downarrow f & \swarrow g & \\ & & Q & & \end{array}$$

is commutative.

This property is equivalent to the condition that the map

$$\text{Hom}(u, Q) : \text{Hom}(M, Q) \longrightarrow \text{Hom}(M', Q)$$

is surjective for every monomorphism $u : M' \rightarrow M$. Since the functor $\text{Hom}(-, Q)$ is left exact, it follows that Q is M -injective if and only if $\text{Hom}(-, Q)$ is exact with respect to every short exact sequence of the form

$$0 \longrightarrow M' \longrightarrow M \longrightarrow M'' \longrightarrow 0.$$

The R -module Q is called *quasi-injective* (or *self-injective*) if it is Q -injective. If Q is M -injective for every R -module M , then Q is called *injective*.

2.1.1 Proposition

Proposition 2.1.1. *Let Q and M be two R -modules. The following statements are equivalent:*

1. Q is M -injective.
2. For every submodule N of M and every morphism $f : N \rightarrow Q$, there exists $g : M \rightarrow Q$ such that $g|_N = f$.
3. For every essential submodule N of M and every morphism $f : N \rightarrow Q$, there exists $g : M \rightarrow Q$ such that $g|_N = f$.

Proof. Implications (1) \Rightarrow (2) and (2) \Rightarrow (3) are obvious.

(2) \Rightarrow (1). Let M'_R , $0 \longrightarrow M' \xrightarrow{u} M$ and $f : M' \rightarrow Q$. Then $u(M') \leq M$. Consider $i : u(M') \rightarrow M$ the canonical injection and $\bar{u} : M' \rightarrow u(M')$ the isomorphism induced by u . There exists $g : M \rightarrow Q$ such that $g \circ i = f \circ \bar{u}^{-1}$. Hence

$$g \circ i \circ \bar{u} = f \quad \text{and therefore} \quad g \circ u = f.$$

$$\begin{array}{ccccccc} 0 & \longrightarrow & M' & \xrightarrow{\bar{u}} & u(M') & \xhookrightarrow{i} & M \\ & & & \searrow f & & \swarrow g & \\ & & & & Q & & \end{array}$$

(3) \Rightarrow (2). Let N be a submodule of M and K a complement of N in M . Then $(N \oplus K) \leq M$. Define $h : N \oplus K \rightarrow Q$ by $h(n + k) = f(n)$ for all $n \in N$, $k \in K$. Since $N \cap K = 0$, the map h is well-defined. There exists $g : M \rightarrow Q$ such that $g|_{N \oplus K} = h$, and hence $g|_N = h|_N = f$.

$$\begin{array}{ccccccc} 0 & \longrightarrow & N & \hookrightarrow & N \oplus K & \hookrightarrow & M \\ & & & \searrow f & \downarrow h & \swarrow g & \\ & & & & Q & & \end{array}$$

□

Proposition 2.1.2. *Let $(M_\alpha)_{\alpha \in \Lambda}$ be a family of R -modules and M an R -module. Then $\prod_{\alpha \in \Lambda} M_\alpha$ is M -injective if and only if each M_α is M -injective.*

Proof. Let N be a submodule of M . Put $P = \prod_{\alpha \in \Lambda} M_\alpha$ and let $\pi_\alpha : P \rightarrow M_\alpha$ be the canonical projections for all $\alpha \in \Lambda$.

" \Leftarrow " Given a morphism $f : N \rightarrow P$, the morphisms $\pi_\alpha \circ f : N \rightarrow M_\alpha$ can be extended to $g_\alpha : M \rightarrow M_\alpha$. There exists $g : M \rightarrow P$ such that $g|_N = f$.

$$\begin{array}{ccccc} 0 & \longrightarrow & N & \longrightarrow & M \\ & & \downarrow f & \swarrow g & \searrow g_\alpha \\ & & P & \xrightarrow{\pi_\alpha} & M_\alpha \end{array}$$

" \Rightarrow " Let $\forall \alpha \in \Lambda$ and $f : N \rightarrow M_\alpha$. Considering the canonical inclusion $\varepsilon_\alpha : M_\alpha \rightarrow P$, since P is M -injective, there exists $g : M \rightarrow P$ which extends $\varepsilon_\alpha \circ f : N \rightarrow P$. Then $\varepsilon_\alpha : M_\alpha \rightarrow P$ extends f and hence M_α is M -injective.

$$\begin{array}{ccccc}
0 & \longrightarrow & N & \longrightarrow & M \\
& & \downarrow f & & \downarrow g \\
& & M_\alpha & & \\
& & \downarrow \varepsilon_\alpha & & \downarrow \pi_\alpha g \\
& & P & & \\
& & \downarrow \pi_\alpha & & \downarrow \pi_\alpha \\
& & M_\alpha & &
\end{array}$$

□

Corollary 2.1.3.

1. Let $(Q_\alpha)_{\alpha \in \Lambda}$ be a family of R -modules. Then $\prod_{\alpha \in \Lambda} Q_\alpha$ is injective if and only if each Q_α is injective for every $\alpha \in \Lambda$.
2. The module $Q_1 \oplus Q_2$ is an injective R -module if and only if each Q_i is injective for $i = 1, 2$. In particular, a direct summand of an injective module is injective.

Proposition 2.1.4. *Let Q be an R -module.*

1. If $0 \rightarrow M' \xrightarrow{f} M \xrightarrow{g} M'' \rightarrow 0$ is an exact sequence of R -modules and Q is M -injective, then Q is M' -injective and M'' -injective.
2. If $(M_\alpha)_{\alpha \in \Lambda}$ is a family of submodules of M such that $M = \sum_{\alpha \in \Lambda} M_\alpha$ and Q is M_α -injective for every α , then Q is M -injective.
3. Let $(N_\alpha)_{\alpha \in \Lambda}$ be a family of R -modules. Then Q is $\bigoplus_{\alpha \in \Lambda} N_\alpha$ -injective if and only if Q is N_α -injective for every $\alpha \in \Lambda$.

Proof. To show that Q is M' -injective, we consider N a submodule of M' and $\varphi : N \rightarrow Q$ a morphism of R -modules. Since Q is M -injective, there exists $\phi : M \rightarrow Q$ such that $\phi \circ f|_N = \varphi$, and hence $\psi \circ f : M' \rightarrow Q$ is a morphism which extends φ .

$$\begin{array}{ccccccc}
0 & \longrightarrow & N & \longrightarrow & M' & \xrightarrow{f} & M \\
& & \downarrow \varphi & & \swarrow & & \downarrow \psi \\
& & Q & & & &
\end{array}$$

Let $h : L \rightarrow M''$ be a monomorphism. We may assume, without loss of generality, that $M' \leq M$ and $M'' = M/M'$. Since $L \cong h(L) \leq M''$, there exist $P \leq M$, $M' \subseteq P$ such that $h(L) = P/M'$ and hence $L \cong P/M'$. We obtain the commutative diagram:

$$\begin{array}{ccccccccc}
& & & & 0 & & 0 & & \\
& & & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & M' & \longrightarrow & P & \longrightarrow & L & \longrightarrow & 0 \\
& & \downarrow & & \downarrow & & \downarrow h & & \\
0 & \longrightarrow & M' & \longrightarrow & M & \longrightarrow & M'' & \longrightarrow & 0
\end{array}$$

Since Q is M -injective, applying the functor $\text{Hom}(-, Q)$ we obtain the commutative diagram:

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \text{Hom}(M'', Q) & \longrightarrow & \text{Hom}(M, Q) & \longrightarrow & \text{Hom}(M', Q) \longrightarrow 0 \\
 & & \downarrow h^* & & \downarrow & & \parallel \\
 0 & \longrightarrow & \text{Hom}(L, Q) & \longrightarrow & \text{Hom}(P, Q) & \longrightarrow & \text{Hom}(M', Q) \\
 & & & & \downarrow & & \\
 & & & & 0 & &
 \end{array}$$

We obtain that $h^* = \text{Hom}(h, Q)$ is an epimorphism, which shows that Q is M'' -injective.

2) Let N be a submodule of M and $f : N \rightarrow Q$ a morphism of R -modules. Consider the set

$$\mathfrak{S} = \{(L, h) \mid N \leq L \leq M, h : L \rightarrow Q, h|_N = f\}.$$

Since $(N, f) \in \mathfrak{S}$, we have $\mathfrak{S} \neq \emptyset$. Define on \mathfrak{S} the order relation $(L_1, h_1) \preceq (L_2, h_2)$ if and only if $L_1 \leq L_2$ and $h_2|_{L_1} = h_1$. One checks that \mathfrak{S} is inductive and, by Zorn's lemma, there exists a maximal element (L_0, g_0) of \mathfrak{S} . To show that $L_0 = M$ it is enough to prove that $M_\alpha \leq L_0$ for every $\alpha \in \Lambda$.

Consider the diagram

$$\begin{array}{ccccc}
 0 & \longrightarrow & L_0 \cap M_\alpha & \xrightarrow{i_\alpha} & M_\alpha \\
 & & \downarrow i_0 & & \swarrow h_\alpha \\
 & & L_0 & & \\
 & & \downarrow g_0 & & \\
 & & Q & &
 \end{array}$$

from which it follows that there exists $h_\alpha : M_\alpha \rightarrow Q$ such that $h_\alpha \circ i_\alpha = g_0 \circ i_\alpha$. Define $h^* : L_0 + M_\alpha \rightarrow Q$ by $h^*(l + m_\alpha) = g_0(l) + h_\alpha(m_\alpha)$, for all $l \in L_0, m_\alpha \in M_\alpha$. If $l + m_\alpha = 0$, then $l = -m_\alpha \in L_0 \cap M_\alpha$ and hence $h^*(l + m_\alpha) = g_0(l) + h_\alpha(l)$, which shows that h^* is well defined. Thus $(L_0 + M_\alpha, h^*) \in \mathfrak{S}$, and since $(L_0, g_0) \preceq (L_0 + M_\alpha, h^*)$, by the maximality of (L_0, g_0) we obtain $L_0 = L_0 + M_\alpha$, i.e. $M_\alpha \leq L_0$ for every $\alpha \in \Lambda$.

3) “ \Rightarrow ” Since $N_\alpha \leq N$ and Q is N -injective, it follows that Q is N_α -injective for every $\alpha \in \Lambda$.

“ \Leftarrow ” Let $N'_\alpha = i_\alpha(N_\alpha)$. Since Q is N_α -injective and $N'_\alpha \cong N_\alpha$, we see that Q is N'_α -injective. Now apply (2). \square

Corollary 2.1.5.

1. The module $Q_1 \oplus Q_2$ is a quasi-injective R -module if and only if each Q_i is Q_j -injective for all $i, j = 1, 2$. In particular, a direct summand of a quasi-injective module is quasi-injective.

2. The module Q^n is quasi-injective over R if and only if Q is quasi-injective.

Corollary 2.1.6. *Let Q and M be two R -modules. Then Q is M -injective if and only if Q is mR -injective for every $m \in M$.*

Proof. The implication “ \Rightarrow ” is clear.

For “ \Leftarrow ”, since $M = \sum_{m \in M} mR$, it follows from 2.1.4(2) that Q is M -injective. \square

Theorem 2.1.7 (Baer’s criterion). *For an R -module Q the following statements are equivalent:*

1. Q is injective.
2. Q is R -injective.
3. For every right ideal I of R and every morphism $f : I \rightarrow Q$ there exists $x \in Q$ such that $f(a) = xa$ for all $a \in I$.

Proof. The implication (1) \Rightarrow (2) is obvious.

(2) \Rightarrow (1). Let M be an R -module and $x \in M$. Since $\varphi_x : R \rightarrow xR$, $\varphi_x(a) = xa$ for all $a \in R$, is a surjective morphism of R -modules, it follows that $R/\text{Ker } \varphi_x \cong xR$. As Q is R -injective, 2.1.4(1) implies that Q is $R/\text{Ker } \varphi_x$ -injective, and hence Q is xR -injective for every $x \in M$. Therefore, by 2.1.6 we obtain that Q is M -injective.

(2) \Rightarrow (3). Let I be a right ideal of R and $f : I \rightarrow Q$. There exists $g : R \rightarrow Q$ such that $g|_I = f$. Put $x = g(1) \in Q$. Then $f(a) = g(a) = ag(1) = xa$ for all $a \in I$.

(3) \Rightarrow (2). Suppose that for a morphism $f : I \rightarrow Q$ there exists $x \in Q$ with $f(a) = xa$ for all $a \in I$. Define $g : R \rightarrow Q$ by $g(r) = xr$ for all $r \in R$. Then clearly $g|_I = f$. \square

Definition 2.1.8.

1. An R -module Q is called *divisible* if for every $y \in Q$ and every $a \in R$ which is not a zero divisor, there exists $x \in Q$ such that $ax = y$. It is easily seen that any factor module of a divisible module is divisible.
2. A commutative integral domain is called a *PID ring* (principal ideal domain) if every ideal of it is principal.

Proposition 2.1.9.

1. Every injective module is divisible.
2. Let R be a PID.
 - (i) If Q is an R -module, then Q is injective if and only if it is divisible.
 - (ii) If I is a non-zero ideal of R , then R/I is a quasi-injective R -module. In particular, \mathbb{Z}_n is a quasi-injective \mathbb{Z} -module for every $n \geq 1$.

Proof.

1) Let Q be a divisible R -module, $y \in Q$ and $a \in R$ a non-zero divisor. Define $f : aR \rightarrow Q$ by $f(ax) = yx$ for all $x \in R$. Since a is not a zero divisor, f is well defined. By Baer's criterion there exists $x \in Q$ such that $f(\lambda) = x\lambda$ for all $\lambda \in aR$. In particular $y = f(a) = f(a \cdot 1) = xa$; hence Q is divisible.

2)(i) The implication " \Rightarrow " follows immediately from (1).

" \Leftarrow " Let Q be a divisible R -module and I a right ideal of R . Then there exists $a \in R$ with $I = aR$. Given a morphism of R -modules $f : I \rightarrow Q$, choose $x \in Q$ with $f(a) = xa$. For any $r \in R$, $f(ar) = f(a)r = xar$, so by Baer's criterion Q is injective.

2)(ii) Let $I = aR$ and $J = bR$ be non-zero ideals of R with $I \subseteq J$, and let $f : J/I \rightarrow R/I$ be a morphism of R -modules. There exists $c \in R$ such that $a = bc$. Write $f(\widehat{b}) = \widehat{x} \in R/I$. Then $xc \in I = aR$, so there is $a_1 \in R$ with $xc = aa_1$, and hence $x = ba_1$.

Define $g : R/I \rightarrow R/I$ by $g(\widehat{r}) = \widehat{a_1 r}$ for all $r \in R$. Then g is an R -module homomorphism and $g(\widehat{b}) = \widehat{x}$, so $g|_{J/I} = f$. Thus R/I is quasi-injective. In particular, for $R = \mathbb{Z}$ and $I = n\mathbb{Z}$ we obtain that \mathbb{Z}_n is a quasi-injective \mathbb{Z} -module for every $n \geq 1$. \square

Corollary 2.1.10. *An abelian group G is an injective \mathbb{Z} -module if and only if G is divisible.*

Corollary 2.1.11.

1. \mathbb{Q} and \mathbb{Z}_{p^∞} are injective \mathbb{Z} -modules.
2. Any direct sum of injective \mathbb{Z} -modules is an injective \mathbb{Z} -module.
3. Any factor group of an injective \mathbb{Z} -module is injective.

Lemma 2.1.12. *Let A, S, T be rings and let ${}_A M_S, {}_A N_T$ be bimodules. Then $\text{Hom}_A(M, N)$ has a structure of left S -module and right T -module given by*

$$(s \cdot f)(x) = f(xs), \quad (f \cdot t)(x) = f(x)t,$$

for $s \in S, t \in T, x \in M, f \in \text{Hom}_A(M, N)$.

Proof. Let $a, b \in A$ and $x, y \in M$. Then

$$(s \cdot f)(ax + by) = f((ax + by)s) = f((ax)s) + f((by)s) = af(xs) + bf(ys) = a(s \cdot f)(x) + b(s \cdot f)(y),$$

so $s \cdot f \in \text{Hom}_A(M, N)$. Similarly $f \cdot t \in \text{Hom}_A(M, N)$.

For $s, s' \in S$ and $f, g \in \text{Hom}_A(M, N)$ we have

$$(s \cdot (f + g))(x) = (f + g)(xs) = f(xs) + g(xs) = (s \cdot f)(x) + (s \cdot g)(x), \quad (1)$$

$$((s + s') \cdot f)(x) = f(x(s + s')) = f(xs + xs') = f(xs) + f(xs') = (s \cdot f)(x) + (s' \cdot f)(x), \quad (2)$$

$$((ss') \cdot f)(x) = f(xss') = f((xs)s') = (s' \cdot f)(xs) = (s \cdot (s' \cdot f))(x), \quad (3)$$

$$(1_S \cdot f)(x) = f(x1_S) = f(x). \quad (4)$$

From (1)–(4) we see that $\text{Hom}_A(M, N)$ is a left S -module. A similar computation shows that $\text{Hom}_A(M, N)$ is a right T -module. Moreover, $((s \cdot f) \cdot t)(x) = (s \cdot f)(xt) = f(xts)t$ and $(s \cdot (f \cdot t))(x) = f(xts)t$, so the two actions commute and $\text{Hom}_A(M, N)$ is an S – T bimodule. \square

Proposition 2.1.13 (Eckmann–Schopf). *Let Q be a divisible abelian group. Then the left R -module $\text{Hom}_{\mathbb{Z}}(R, Q)$ is injective.*

Proof. By the previous lemma, $\text{Hom}_{\mathbb{Z}}(R, Q)$ carries a structure of left R -module given by

$$(r \cdot f)(a) = f(ar), \quad \forall a, r \in R, f \in \text{Hom}_{\mathbb{Z}}(R, Q).$$

Let I be a left ideal of R and $h : I \rightarrow \text{Hom}_{\mathbb{Z}}(R, Q)$ a morphism of left R -modules. Define

$$\gamma : \mathbb{Z}I \longrightarrow \mathbb{Z}Q, \quad \gamma(a) = h(a)(1).$$

Then γ is a morphism of \mathbb{Z} -modules. Since Q is \mathbb{Z} -injective, there exists $\tilde{\gamma} : \mathbb{Z}R \rightarrow \mathbb{Z}Q$ such that $\tilde{\gamma}|_I = \gamma$. For $a \in I$ and $r \in R$ we have

$$(a \cdot \tilde{\gamma})(r) = \tilde{\gamma}(ra) = h(ra)(1) = (r \cdot h(a))(1) = h(a)(r),$$

hence $h(a) = a \cdot \tilde{\gamma}$ for all $a \in I$. By Baer's criterion, $\text{Hom}_{\mathbb{Z}}(R, Q)$ is an injective left R -module. \square

Proposition 2.1.14. *Every left R -module M can be embedded into an injective left R -module.*

Proof. There exists a free abelian group $\mathbb{Z}^{(A)}$ and a surjective \mathbb{Z} -morphism $f : \mathbb{Z}^{(A)} \rightarrow M$. Hence

$$\mathbb{Z}M \cong \mathbb{Z}^{(A)} / \ker f \subseteq \mathbb{Q}^{(A)} / \ker f,$$

and therefore there is a divisible abelian group G with $\mathbb{Z}M \subseteq \mathbb{Z}G$. Applying the functor $\text{Hom}_{\mathbb{Z}}(R, -)$ we obtain a monomorphism

$${}_R M \cong \text{Hom}_{\mathbb{Z}}(R, M) \hookrightarrow \text{Hom}_{\mathbb{Z}}(R, G).$$

Since G is divisible, Proposition 2.1.13 implies that $\text{Hom}_{\mathbb{Z}}(R, G)$ is an injective left R -module. Thus M embeds into an injective left R -module. \square

Proposition 2.1.15. *Let Q be an R -module. Then Q is injective if and only if every short exact sequence*

$$0 \longrightarrow Q \xrightarrow{f} M \xrightarrow{g} M' \longrightarrow 0$$

splits.

Proof. “ \Rightarrow ”. Assume Q is injective. Exactness implies that f is a monomorphism. By injectivity of Q there exists $h : M \rightarrow Q$ with $hf = \text{id}_Q$, hence the sequence splits.

“ \Leftarrow ”. By Proposition 2.1.14 there is an injective R -module Q' and a monomorphism $i : Q \rightarrow Q'$. Consider the exact sequence

$$0 \longrightarrow Q \xrightarrow{i} Q' \longrightarrow Q'/i(Q) \longrightarrow 0.$$

By the hypothesis this sequence splits, so Q is a direct summand of Q' . A direct summand of an injective module is injective, therefore Q is injective. \square

2.2 Injective envelopes

Definition 2.2.1. Let M be an R -module. A pair (E, i) is called an *injective envelope* of M if E is injective and $i : M \rightarrow E$ is an essential monomorphism.

Proposition 2.2.2. Let Q be an injective R -module. Then every complement submodule of Q is a direct summand of Q .

Proof. Let K be a submodule of Q and N a complement of K in Q , that is, $K \cap N = 0$ and $K + N$ is an essential submodule of Q . Then $(K + N)/N \cong Q/N$. Define $g : (K + N)/N \rightarrow Q$ by

$$g((x + y) + N) = x, \quad x \in K, y \in N.$$

Since $K \cap N = 0$, g is well defined and injective. As Q is injective, there exists $h : Q/N \rightarrow Q$ with $h|_{(K+N)/N} = g$. Because $(K + N)/N \cong Q/N$ and g is a monomorphism, h is also a monomorphism. We have $K = \text{Im } g = h((K + N)/N) \subseteq h(Q/N)$. As K is a closed submodule, it follows that $K = h(Q/N)$. Since h is a monomorphism, $(K + N)/N = Q/N$, hence $K + N = Q$. Thus K is a direct summand of Q . \square

Theorem 2.2.3 (Eckmann–Schopf). Every R -module M has an injective envelope, unique up to isomorphism.

Proof. By Proposition 2.1.14 there exists an injective R -module Q with $M \leq Q$. Let E be a maximal essential extension of M in Q . Then E is a complement submodule of Q , and by the previous proposition E is injective. Hence (E, i) , where $i : M \hookrightarrow E$ is the inclusion, is an injective envelope of M .

For uniqueness, let (E_1, i_1) and (E_2, i_2) be two injective envelopes of M . Since E_2 is injective, there exists $f : E_1 \rightarrow E_2$ with $f i_1 = i_2$. The map i_2 is a monomorphism and i_1 is an essential monomorphism, so (using 1.4) f is a monomorphism. Thus $E_1 \cong f(E_1)$ and $E_2 = f(E_1) \oplus E_3$ for some submodule E_3 . But $i_2(M) \subseteq f(E_1)$, hence $i_2(M) \cap E_3 = 0$. Since i_2 is essential, we must have $E_3 = 0$, so $E_2 = f(E_1)$ and f is an isomorphism. \square

In practice we fix one representative of this isomorphism class and denote it by $E(M)$, with $M \leq E(M)$.

Proposition 2.2.4. Let M be an R -module and $i : M \rightarrow Q$ a monomorphism with Q_R injective. The following statements are equivalent:

1. (Q, i) is an injective envelope of M ;
2. for every monomorphism $f : M \rightarrow Q'$ with Q' injective, there exists a monomorphism $g : Q \rightarrow Q'$ such that $gi = f$.

Proof. (1) \Rightarrow (2). Let $f : M \rightarrow Q'$ be a monomorphism with Q' injective. By injectivity of Q' there is $u : Q \rightarrow Q'$ with $ui = f$. Because i is an essential monomorphism and Q' is injective, the image $u(Q)$ is a complement of $f(M)$; by the definition of injective envelope this forces u to be a monomorphism. Set $g = u$.

(2) \Rightarrow (1). Let $(E(M), j)$ be an injective envelope of M . Applying (2) to $f = j$ we obtain a monomorphism $g : Q \rightarrow E(M)$ with $gi = j$. Since j is an essential monomorphism, it follows that i is also essential; hence (Q, i) is an injective envelope of M . \square

Proposition 2.2.5. *For any family of right R -modules M_1, M_2, \dots, M_n we have*

$$E\left(\bigoplus_{i=1}^n M_i\right) \cong \bigoplus_{i=1}^n E(M_i).$$

Proof. By 1.9, $\bigoplus_{i=1}^n E(M_i)$ is an essential extension of $\bigoplus_{i=1}^n M_i$. Moreover,

$$\bigoplus_{i=1}^n E(M_i) \cong \prod_{i=1}^n E(M_i),$$

so by 2.1.3 the module $\bigoplus_{i=1}^n E(M_i)$ is injective. By uniqueness of injective envelopes we obtain

$$E\left(\bigoplus_{i=1}^n M_i\right) \cong \bigoplus_{i=1}^n E(M_i).$$

\square

Theorem 2.2.6. *Let Q and M be two R -modules. Then Q is M -injective if and only if $f(M) \subseteq Q$ for every $f \in \text{Hom}(E(M), E(Q))$.*

Proof. “ \Rightarrow ” Let $f \in \text{Hom}(E(M), E(Q))$ and set

$$K := \{m \in M \mid f(m) \in Q\}.$$

Since Q is M -injective, there exists a morphism $\bar{f} : M \rightarrow Q$ such that $\bar{f}|_K = f|_K$. We claim that

$$Q \cap (\bar{f} - f)(M) = 0.$$

Take $x \in Q$ and $m \in M$ with $x = (\bar{f} - f)(m)$. Then

$$f(m) = \bar{f}(m) - x \in Q,$$

so $m \in K$. Hence

$$x = \bar{f}(m) - f(m) = f(m) - f(m) = 0,$$

and therefore $Q \cap (\bar{f} - f)(M) = 0$. Since Q is an essential submodule of $E(Q)$, it follows that $(\bar{f} - f)(M) = 0$. Thus $f(M) = \bar{f}(M) \subseteq Q$.

“ \Leftarrow ” As $E(Q)$ is injective, it is enough to work with $f \in \text{Hom}(M, E(Q))$. Let N be a submodule of M and $g : N \rightarrow Q$ a morphism of R -modules. Because $E(Q)$ is injective, there exists $\tilde{g} : M \rightarrow E(Q)$ such that $\tilde{g}|_N = i \circ g$, where $i : Q \rightarrow E(Q)$ is the canonical injection. By hypothesis, $\tilde{g}(M) \subseteq Q$, so identifying \tilde{g} with its corestriction to Q we obtain a morphism $h : M \rightarrow Q$ with $h|_N = g$. Therefore Q is M -injective. \square

Corollary 2.2.7. *An R -module Q is quasi-injective if and only if $f(Q) \subseteq Q$ for every $f \in \text{End}(E(Q))$.*

Theorem 2.2.8 (Matlis–Bass). *Let R be a ring. Then R is right noetherian if and only if, for every simple right R -module S_i ($i \geq 1$),*

$$Q := \bigoplus_{i=1}^{\infty} E(S_i)$$

is an injective right R -module.

Proof. “ \Rightarrow ” Let L be a right ideal of R ,

$$Q = \bigoplus_{i=1}^{\infty} E(S_i)$$

and let $f : L \rightarrow Q$ be a morphism of right R -modules. There exist elements $a_1, \dots, a_n \in L$ such that

$$L = a_1 R + a_2 R + \dots + a_n R.$$

Clearly, there is $m \geq 1$ with $f(a_k) \in \bigoplus_{j=1}^m E(S_j)$ for all $k = 1, \dots, n$, hence

$$\Im f \subseteq \bigoplus_{j=1}^m E(S_j).$$

Since $\bigoplus_{j=1}^m E(S_j)$ is injective, there exists $g : R \rightarrow \bigoplus_{j=1}^m E(S_j)$ such that $g|_L = f$. Let $\bar{f} = i \circ g$, where $i : \bigoplus_{j=1}^m E(S_j) \rightarrow Q$ is the canonical injection. Then $\bar{f}|_L = f$, so Q is injective.

“ \Leftarrow ” Suppose that R is not right noetherian. Then there exists a strictly ascending chain of finitely generated right ideals:

$$L_1 \subsetneq L_2 \subsetneq \dots \subsetneq L_n \subsetneq \dots$$

By Krull’s lemma, for every $n \geq 1$ there exists a maximal submodule $M_n \subsetneq L_n$ such that

$$L_{n-1} \subseteq M_n \quad \text{for all } n \geq 2.$$

Set

$$L := \bigcup_{k=1}^{\infty} L_k, \quad \pi_k : L_k \longrightarrow L_k/M_k$$

for the canonical projections, and put

$$E_k := E(L_k/M_k) \quad \text{for each } k \geq 1.$$

Then

$$E := \bigoplus_{k=1}^{\infty} E_k$$

is injective, and

$$f : L \longrightarrow E, \quad f(a) = \sum_{k=1}^{\infty} \pi_k(a)$$

is well defined. There exists an element $x \in E_1 \oplus \cdots \oplus E_n$ such that $f(a) = xa$ for all $a \in L$. It follows that $\pi_k(a) = 0$ for every $k \geq n + 1$, that is, $a \in M_k$ for all $k \geq n + 1$. Hence

$$L \subseteq M_{n+1} \subsetneq L_{n+1} \subseteq M_{n+2} \subsetneq L_{n+2} \subseteq \cdots \subseteq L,$$

a contradiction. Therefore R is right noetherian. □

