

# November 17, 2023 Class 14: RNAseq Mini Project

Savannah Bogus A69027475

## Differential Expression Analysis

```
library(DESeq2)
```

```
Loading required package: S4Vectors
```

```
Loading required package: stats4
```

```
Loading required package: BiocGenerics
```

```
Attaching package: 'BiocGenerics'
```

```
The following objects are masked from 'package:stats':
```

```
IQR, mad, sd, var, xtabs
```

```
The following objects are masked from 'package:base':
```

```
anyDuplicated, aperm, append, as.data.frame, basename, cbind,  
colnames, dirname, do.call, duplicated, eval, evalq, Filter, Find,  
get, grep, grepl, intersect, is.unsorted, lapply, Map, mapply,  
match, mget, order, paste, pmax, pmax.int, pmin, pmin.int,  
Position, rank, rbind, Reduce, rownames, sapply, setdiff, sort,  
table, tapply, union, unique, unsplit, which.max, which.min
```

Attaching package: 'S4Vectors'

The following object is masked from 'package:utils':

findMatches

The following objects are masked from 'package:base':

expand.grid, I, unname

Loading required package: IRanges

Loading required package: GenomicRanges

Loading required package: GenomeInfoDb

Warning: package 'GenomeInfoDb' was built under R version 4.3.2

Loading required package: SummarizedExperiment

Warning: package 'SummarizedExperiment' was built under R version 4.3.2

Loading required package: MatrixGenerics

Loading required package: matrixStats

Attaching package: 'MatrixGenerics'

The following objects are masked from 'package:matrixStats':

colAlls, colAnyNAs, colAnys, colAvgsPerRowSet, colCollapse,  
colCounts, colCummaxs, colCummins, colCumprods, colCumsums,  
colDiffs, colIQRDiffs, colIQRs, colLogSumExps, colMadDiffs,  
colMads, colMaxs, colMeans2, colMedians, colMins, colOrderStats,  
colProds, colQuantiles, colRanges, colRanks, colSdDiffs, colSds,  
colSums2, colTabulates, colVarDiffs, colVars, colWeightedMads,  
colWeightedMeans, colWeightedMedians, colWeightedSds,

```
colWeightedVars, rowAlls, rowAnyNAs, rowAnys, rowAvgsPerColSet,
rowCollapse, rowCounts, rowCummaxs, rowCummins, rowCumprods,
rowCumsums, rowDiffs, rowIQRDiffs, rowIQRs, rowLogSumExps,
rowMadDiffs, rowMads, rowMaxs, rowMeans2, rowMedians, rowMins,
rowOrderStats, rowProds, rowQuantiles, rowRanges, rowRanks,
rowSdDiffs, rowSds, rowSums2, rowTabulates, rowVarDiffs, rowVars,
rowWeightedMads, rowWeightedMeans, rowWeightedMedians,
rowWeightedSds, rowWeightedVars
```

Loading required package: Biobase

Welcome to Bioconductor

```
Vignettes contain introductory material; view with
'browseVignettes()'. To cite Bioconductor, see
'citation("Biobase")', and for packages 'citation("pkgname")'.
```

Attaching package: 'Biobase'

The following object is masked from 'package:MatrixGenerics':

```
rowMedians
```

The following objects are masked from 'package:matrixStats':

```
anyMissing, rowMedians
```

```
metaFile <- "GSE37704_metadata.csv"
countFile <- "GSE37704_featurecounts.csv"

# Import metadata
colData = read.csv(metaFile, row.names=1)
head(colData)
```

```
              condition
SRR493366 control_sirna
SRR493367 control_sirna
SRR493368 control_sirna
SRR493369      hoxa1_kd
SRR493370      hoxa1_kd
SRR493371      hoxa1_kd
```

```
# Import countdata
countData = read.csv(countFile, row.names=1)
head(countData)
```

	length	SRR493366	SRR493367	SRR493368	SRR493369	SRR493370
ENSG00000186092	918	0	0	0	0	0
ENSG00000279928	718	0	0	0	0	0
ENSG00000279457	1982	23	28	29	29	28
ENSG00000278566	939	0	0	0	0	0
ENSG00000273547	939	0	0	0	0	0
ENSG00000187634	3214	124	123	205	207	212

	SRR493371
ENSG00000186092	0
ENSG00000279928	0
ENSG00000279457	46
ENSG00000278566	0
ENSG00000273547	0
ENSG00000187634	258

I need to remove the weird column though.

```
countData <- as.matrix(countData[,-1])
head(countData)
```

	SRR493366	SRR493367	SRR493368	SRR493369	SRR493370	SRR493371
ENSG00000186092	0	0	0	0	0	0
ENSG00000279928	0	0	0	0	0	0
ENSG00000279457	23	28	29	29	28	46
ENSG00000278566	0	0	0	0	0	0
ENSG00000273547	0	0	0	0	0	0
ENSG00000187634	124	123	205	207	212	258

Now I have to get rid of the zeros.

```
countData = countData[-which(rowSums(countData)==0), ]
head(countData)
```

	SRR493366	SRR493367	SRR493368	SRR493369	SRR493370	SRR493371
ENSG00000279457	23	28	29	29	28	46
ENSG00000187634	124	123	205	207	212	258

ENSG00000188976	1637	1831	2383	1226	1326	1504
ENSG00000187961	120	153	180	236	255	357
ENSG00000187583	24	48	65	44	48	64
ENSG00000187642	4	9	16	14	16	16

```
dds = DESeqDataSetFromMatrix(countData=countData,
                              colData=colData,
                              design=~condition)
```

Warning in DESeqDataSet(se, design = design, ignoreRank): some variables in design formula are characters, converting to factors

```
dds = DESeq(dds)
```

estimating size factors

estimating dispersions

gene-wise dispersion estimates

mean-dispersion relationship

final dispersion estimates

fitting model and testing

```
res=results(dds)
summary(res)
```

out of 15975 with nonzero total read count

adjusted p-value < 0.1

LFC > 0 (up) : 4349, 27%

LFC < 0 (down) : 4396, 28%

outliers [1] : 0, 0%


low counts [2] : 1237, 7.7%

(mean count < 0)

[1] see 'cooksCutoff' argument of ?results

[2] see 'independentFiltering' argument of ?results

```
plot( res$log2FoldChange, -log(res$padj),col="darkcyan" )
```




```
# Make a color vector for all genes
mycolors <- rep("cyan", nrow(res) )

#coloring genes with fold change above 2
mycolors[ abs(res$log2FoldChange) > 2 ] <- "violet"

# coloring genes w adjusted p-value less than 0.01
# and absolute fold change more than 2
inds <- (res$padj<0.01) & (abs(res$log2FoldChange) > 2 )
mycolors[ inds ] <- "darkcyan"

plot( res$log2FoldChange, -log(res$padj), col=mycolors, xlab="Log2(FoldChange)", ylab="-Log10(adjusted p-value)" )
```



```
library("AnnotationDbi")
library("org.Hs.eg.db")
```

```
columns(org.Hs.eg.db)
```

[1]	"ACCNUM"	"ALIAS"	"ENSEMBL"	"ENSEMBLPROT"	"ENSEMBLTRANS"
[6]	"ENTREZID"	"ENZYME"	"EVIDENCE"	"EVIDENCEALL"	"GENENAME"
[11]	"GENETYPE"	"GO"	"GOALL"	"IPI"	"MAP"
[16]	"OMIM"	"ONTOLOGY"	"ONTOLOGYALL"	"PATH"	"PFAM"
[21]	"PMID"	"PROSITE"	"REFSEQ"	"SYMBOL"	"UCSCKG"
[26]	"UNIPROT"				

```
res$symbol = mapIds(org.Hs.eg.db,
  keys=row.names(res),
  keytype="ENSEMBL",
  column="SYMBOL",
  multiVals="first")
```

'select()' returned 1:many mapping between keys and columns

```
res$entrez = mapIds(org.Hs.eg.db,  
                    keys=row.names(res),  
                    keytype="ENSEMBL",  
                    column="ENTREZID",  
                    multiVals="first")
```

'select()' returned 1:many mapping between keys and columns

```
res$name = mapIds(org.Hs.eg.db,  
                  keys=row.names(res),  
                  keytype="ENSEMBL",  
                  column="GENENAME",  
                  multiVals="first")
```

'select()' returned 1:many mapping between keys and columns

```
head(res, 10)
```

log2 fold change (MLE): condition hoxa1 kd vs control sirna

Wald test p-value: condition hoxa1 kd vs control sirna

DataFrame with 10 rows and 9 columns

	baseMean	log2FoldChange	lfcSE	stat	pvalue
	<numeric>	<numeric>	<numeric>	<numeric>	<numeric>
ENSG00000279457	29.913579	0.1792571	0.3248216	0.551863	5.81042e-01
ENSG00000187634	183.229650	0.4264571	0.1402658	3.040350	2.36304e-03
ENSG00000188976	1651.188076	-0.6927205	0.0548465	-12.630158	1.43989e-36
ENSG00000187961	209.637938	0.7297556	0.1318599	5.534326	3.12428e-08
ENSG00000187583	47.255123	0.0405765	0.2718928	0.149237	8.81366e-01
ENSG00000187642	11.979750	0.5428105	0.5215599	1.040744	2.97994e-01
ENSG00000188290	108.922128	2.0570638	0.1969053	10.446970	1.51282e-25
ENSG00000187608	350.716868	0.2573837	0.1027266	2.505522	1.22271e-02
ENSG00000188157	9128.439422	0.3899088	0.0467163	8.346304	7.04321e-17
ENSG00000237330	0.158192	0.7859552	4.0804729	0.192614	8.47261e-01
	padj	symbol	entrez		name
	<numeric>	<character>	<character>		<character>
ENSG00000279457	6.86555e-01	NA	NA		NA



ENSG00000187634	5.15718e-03	SAMD11	148398	sterile alpha motif ..
ENSG00000188976	1.76549e-35	NOC2L	26155	NOC2 like nucleolar ..
ENSG00000187961	1.13413e-07	KLHL17	339451	kelch like family me..
ENSG00000187583	9.19031e-01	PLEKHN1	84069	pleckstrin homology ..
ENSG00000187642	4.03379e-01	PERM1	84808	PPARGC1 and ESRR ind..
ENSG00000188290	1.30538e-24	HES4	57801	hes family bHLH tran..
ENSG00000187608	2.37452e-02	ISG15	9636	ISG15 ubiquitin like..
ENSG00000188157	4.21963e-16	AGRN	375790	agrin
ENSG00000237330	NA	RNF223	401934	ring finger protein ..

```
res = res[order(res$pvalue),]
write.csv(res, file="deseq_results.csv")
```

## Pathway Analysis

```
library(pathview)
```

#####

Pathview is an open source software package distributed under GNU General Public License version 3 (GPLv3). Details of GPLv3 is available at <http://www.gnu.org/licenses/gpl-3.0.html>. Particullary, users are required to formally cite the original Pathview paper (not just mention it) in publications or products. For details, do citation("pathview") within R.

The pathview downloads and uses KEGG data. Non-academic uses may require a KEGG license agreement (details at <http://www.kegg.jp/kegg/legal.html>).

#####

```
library(gage)
```

```
library(gageData)
data(kegg.sets.hs)
data(sigmet.idx.hs)
```

Now, I'm going to limit it to signaling and metabolic pathways.

```
kegg.sets.hs = kegg.sets.hs[sigmet.idx.hs]
head(kegg.sets.hs, 3)
```

```
$`hsa00232 Caffeine metabolism`
```

```
[1] "10" "1544" "1548" "1549" "1553" "7498" "9"
```

```
$`hsa00983 Drug metabolism - other enzymes`
```

```
[1] "10" "1066" "10720" "10941" "151531" "1548" "1549" "1551"
[9] "1553" "1576" "1577" "1806" "1807" "1890" "221223" "2990"
[17] "3251" "3614" "3615" "3704" "51733" "54490" "54575" "54576"
[25] "54577" "54578" "54579" "54600" "54657" "54658" "54659" "54963"
[33] "574537" "64816" "7083" "7084" "7172" "7363" "7364" "7365"
[41] "7366" "7367" "7371" "7372" "7378" "7498" "79799" "83549"
[49] "8824" "8833" "9" "978"
```

```
$`hsa00230 Purine metabolism`
```

```
[1] "100" "10201" "10606" "10621" "10622" "10623" "107" "10714"
[9] "108" "10846" "109" "111" "11128" "11164" "112" "113"
[17] "114" "115" "122481" "122622" "124583" "132" "158" "159"
[25] "1633" "171568" "1716" "196883" "203" "204" "205" "221823"
[33] "2272" "22978" "23649" "246721" "25885" "2618" "26289" "270"
[41] "271" "27115" "272" "2766" "2977" "2982" "2983" "2984"
[49] "2986" "2987" "29922" "3000" "30833" "30834" "318" "3251"
[57] "353" "3614" "3615" "3704" "377841" "471" "4830" "4831"
[65] "4832" "4833" "4860" "4881" "4882" "4907" "50484" "50940"
[73] "51082" "51251" "51292" "5136" "5137" "5138" "5139" "5140"
[81] "5141" "5142" "5143" "5144" "5145" "5146" "5147" "5148"
[89] "5149" "5150" "5151" "5152" "5153" "5158" "5167" "5169"
[97] "51728" "5198" "5236" "5313" "5315" "53343" "54107" "5422"
[105] "5424" "5425" "5426" "5427" "5430" "5431" "5432" "5433"
[113] "5434" "5435" "5436" "5437" "5438" "5439" "5440" "5441"
[121] "5471" "548644" "55276" "5557" "5558" "55703" "55811" "55821"
[129] "5631" "5634" "56655" "56953" "56985" "57804" "58497" "6240"
[137] "6241" "64425" "646625" "654364" "661" "7498" "8382" "84172"
[145] "84265" "84284" "84618" "8622" "8654" "87178" "8833" "9060"
[153] "9061" "93034" "953" "9533" "954" "955" "956" "957"
[161] "9583" "9615"
```

We have to get ready for the gage function format, which is what I'm doing here.

```
foldchanges = res$log2FoldChange
names(foldchanges) = res$entrez
head(foldchanges)
```

```
      1266      54855      1465      51232      2034      2317
-2.422719  3.201955 -2.313738 -2.059631 -1.888019 -1.649792
```

Next, I'm actually using the gage function, with `same.dir=TRUE` to help with order.

```
keggres = gage(foldchanges, gsets=kegg.sets.hs,same.dir=TRUE)
attributes(keggres)
```

```
$names
[1] "greater" "less"    "stats"
```

Checking out what is downregulated.

```
head(keggres$less)
```

	p.geomean	stat.mean	p.val
hsa04110 Cell cycle	8.995727e-06	-4.378644	8.995727e-06
hsa03030 DNA replication	9.424076e-05	-3.951803	9.424076e-05
hsa03013 RNA transport	1.375901e-03	-3.028500	1.375901e-03
hsa03440 Homologous recombination	3.066756e-03	-2.852899	3.066756e-03
hsa04114 Oocyte meiosis	3.784520e-03	-2.698128	3.784520e-03
hsa00010 Glycolysis / Gluconeogenesis	8.961413e-03	-2.405398	8.961413e-03

	q.val	set.size	exp1
hsa04110 Cell cycle	0.001448312	121	8.995727e-06
hsa03030 DNA replication	0.007586381	36	9.424076e-05
hsa03013 RNA transport	0.073840037	144	1.375901e-03
hsa03440 Homologous recombination	0.121861535	28	3.066756e-03
hsa04114 Oocyte meiosis	0.121861535	102	3.784520e-03
hsa00010 Glycolysis / Gluconeogenesis	0.212222694	53	8.961413e-03


And finally getting the pathview look at the pathway, inserted below.

```
pathview(gene.data=foldchanges, pathway.id="hsa04110")
```

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/savannahbogus/Documents/Classes/BGGN 213 Bioinformatics/202

Info: Writing image file hsa04110.pathview.png



```
[1] "hsa04640" "hsa04630" "hsa00140" "hsa04142" "hsa04330"
```

```
pathview(gene.data=foldchanges, pathway.id=keggresids, species="hsa")
```

```
'select()' returned 1:1 mapping between keys and columns
```

```
Info: Working in directory /Users/savannahbogus/Documents/Classes/BGGN 213 Bioinformatics/202
```

```
Info: Writing image file hsa04640.pathview.png
```

```
'select()' returned 1:1 mapping between keys and columns
```

```
Info: Working in directory /Users/savannahbogus/Documents/Classes/BGGN 213 Bioinformatics/202
```

```
Info: Writing image file hsa04630.pathview.png
```

```
'select()' returned 1:1 mapping between keys and columns
```

```
Info: Working in directory /Users/savannahbogus/Documents/Classes/BGGN 213 Bioinformatics/202
```

```
Info: Writing image file hsa00140.pathview.png
```

```
'select()' returned 1:1 mapping between keys and columns
```

```
Info: Working in directory /Users/savannahbogus/Documents/Classes/BGGN 213 Bioinformatics/202
```

```
Info: Writing image file hsa04142.pathview.png
```

```
Info: some node width is different from others, and hence adjusted!
```

```
'select()' returned 1:1 mapping between keys and columns
```

```
Info: Working in directory /Users/savannahbogus/Documents/Classes/BGGN 213 Bioinformatics/202
```



```
Info: Writing image file hsa04330.pathview.png
```











## Same procedure for the downregulated.

First, get the top 5 downregulated pathways.

```
keggrespathwaydown <- rownames(keggres$less)[1:5]
keggresidsdown = substr(keggrespathwaydown, start=1, stop=8)
keggresidsdown
```

```
[1] "hsa04110" "hsa03030" "hsa03013" "hsa03440" "hsa04114"
```

```
pathview(gene.data=foldchanges, pathway.id=keggresidsdown, species="hsa")
```

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/savannahbogus/Documents/Classes/BGGN 213 Bioinformatics/2020

Info: Writing image file hsa04110.pathview.png

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/savannahbogus/Documents/Classes/BGGN 213 Bioinformatics/2020

Info: Writing image file hsa03030.pathview.png

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/savannahbogus/Documents/Classes/BGGN 213 Bioinformatics/2020

Info: Writing image file hsa03013.pathview.png

'select()' returned 1:1 mapping between keys and columns


Info: Working in directory /Users/savannahbogus/Documents/Classes/BGGN 213 Bioinformatics/2020

Info: Writing image file hsa03440.pathview.png

'select()' returned 1:1 mapping between keys and columns



Info: Working in directory /Users/savannahbogus/Documents/Classes/BGGN 213 Bioinformatics/202

Info: Writing image file hsa04114.pathview.png





# DNA REPLICATION



## Replication complex (Bacteria)




## Replication complex (Archaea)



## Replication complex (Eukaryotes)



Data on KEGG graph  
Rendered by Pathview







G0:0002009	morphogenesis of an epithelium	1.396681e-04	3.653886	1.396681e-04
G0:0048729	tissue morphogenesis	1.432451e-04	3.643242	1.432451e-04
G0:0007610	behavior	1.925222e-04	3.565432	1.925222e-04
G0:0060562	epithelial tube morphogenesis	5.932837e-04	3.261376	5.932837e-04
G0:0035295	tube development	5.953254e-04	3.253665	5.953254e-04

		q.val	set.size	exp1
G0:0007156	homophilic cell adhesion	0.1952430	113	8.519724e-05
G0:0002009	morphogenesis of an epithelium	0.1952430	339	1.396681e-04
G0:0048729	tissue morphogenesis	0.1952430	424	1.432451e-04
G0:0007610	behavior	0.1968058	426	1.925222e-04
G0:0060562	epithelial tube morphogenesis	0.3566193	257	5.932837e-04
G0:0035295	tube development	0.3566193	391	5.953254e-04

\$less

		p.geomean	stat.mean	p.val
G0:0048285	organelle fission	1.536227e-15	-8.063910	1.536227e-15
G0:0000280	nuclear division	4.286961e-15	-7.939217	4.286961e-15
G0:0007067	mitosis	4.286961e-15	-7.939217	4.286961e-15
G0:0000087	M phase of mitotic cell cycle	1.169934e-14	-7.797496	1.169934e-14
G0:0007059	chromosome segregation	2.028624e-11	-6.878340	2.028624e-11
G0:0000236	mitotic prometaphase	1.729553e-10	-6.695966	1.729553e-10

		q.val	set.size	exp1
G0:0048285	organelle fission	5.843127e-12	376	1.536227e-15
G0:0000280	nuclear division	5.843127e-12	352	4.286961e-15
G0:0007067	mitosis	5.843127e-12	352	4.286961e-15
G0:0000087	M phase of mitotic cell cycle	1.195965e-11	362	1.169934e-14
G0:0007059	chromosome segregation	1.659009e-08	142	2.028624e-11
G0:0000236	mitotic prometaphase	1.178690e-07	84	1.729553e-10

\$stats

		stat.mean	exp1
G0:0007156	homophilic cell adhesion	3.824205	3.824205
G0:0002009	morphogenesis of an epithelium	3.653886	3.653886
G0:0048729	tissue morphogenesis	3.643242	3.643242
G0:0007610	behavior	3.565432	3.565432
G0:0060562	epithelial tube morphogenesis	3.261376	3.261376
G0:0035295	tube development	3.253665	3.253665



## Reactome Analysis

```
sig_genes <- res[res$padj <= 0.05 & !is.na(res$padj), "symbol"]  
print(paste("Total number of significant genes:", length(sig_genes)))
```

```
[1] "Total number of significant genes: 8147"
```

```
write.table(sig_genes, file="significant_genes.txt", row.names=FALSE, col.names=FALSE, quo
```

#Q What pathway has the most significant “Entities p-value”? Do the most significant pathways listed match your previous KEGG results? What factors could cause differences between the two methods?

Cell cycle, mitotic has the most significant entities p-value. It’s the same for the KEGG results, interestingly.