

Simulations of Neutron Multiplicity Experiments with Nuclear Data Perturbations

S.R. Bolding¹, C.J. Solomon²

¹ *Texas A&M University, College station, TX*

² *Los Alamos National Laboratory, Los Alamos, NM*

ANS National Meeting, 14 November 2013



Outline

1. Background
2. Correlated Sampling
3. Methodology
4. Results
5. Conclusions

Outline

1. Background
2. Correlated Sampling
3. Methodology
4. Results
5. Conclusions

Neutron Multiplicity Distributions

Neutron Multiplicity Distributions






Figure: Multiplicity distributions [PANDA Manual, 1991]

- Provide **passive** multiplication information about a subcritical, fissionable system

Constructing a Multiplicity Distribution (Ideal Case)


Constructing a Multiplicity Distribution (Ideal Case)

○ = Detected Neutron




● Normalize to form a PDF

Multiplicity Experiments



*Not to scale


Multiplicity Experiments



*Not to scale

- Performed at LANL for validating subcritical simulations
- Experimental Parameters
 - 94% ^{239}Pu sphere
 - 5 Different HDPE shells:
 - ▶ None, 0.5 cm, 1.0 cm, 1.5 cm, 3.0 cm

Multiplicity Experiments



*Not to scale


- Performed at LANL for validating subcritical simulations
- Experimental Parameters
 - 94% ^{239}Pu sphere
 - 5 Different HDPE shells:
 - ▶ None, 0.5 cm, 1.0 cm, 1.5 cm, 3.0 cm
- Measured multiplicity distributions are well verified
- Repeated with ^{252}Cf

MCNP5 multiplicity simulations

- LANL experiments modeled with a modified **MCNP5** [Solomon, 2011]

MCNP5 multiplicity simulations


- LANL experiments modeled with a modified **MCNP5** [Solomon, 2011]
- Discrepancy b/w **simulation** & **experiment** for Pu, but not ^{252}Cf



Pu with 3.0-cm HDPE reflector

MCNP5 multiplicity simulations


- LANL experiments modeled with a modified **MCNP5** [Solomon, 2011]
- Discrepancy b/w **simulation** & **experiment** for Pu, but not ^{252}Cf
- Previous work by Mattingly [2010]
 - Caused by **nuclear data** of ^{239}Pu
 - Adjusted **energy-integrated $\bar{\nu}$**



Pu with 3.0-cm HDPE reflector

MCNP5 multiplicity simulations

- LANL experiments modeled with a modified **MCNP5** [Solomon, 2011]
- Discrepancy b/w **simulation** & **experiment** for Pu, but not ^{252}Cf
- Previous work by Mattingly [2010]
 - Caused by **nuclear data** of ^{239}Pu
 - Adjusted **energy-integrated $\bar{\nu}$**
- **ENDF** ^{239}Pu $\bar{\nu}$ adjusted to match k_{eff}
 - $\bar{\nu}$ is $\sim 2\sigma$ **above** measured data for $E < 1.5$ MeV



Pu with 3.0-cm HDPE reflector

Objectives

Objectives

- Reduce discrepancy in multiplicity distributions w/o significantly altering k_{eff}

Objectives

- Reduce discrepancy in multiplicity distributions w/o significantly altering k_{eff}
- Perform energy-dependent perturbations to $\bar{\nu}(E)$ in ^{239}Pu
 - Random samples that preserve covariance data
 - Compare experimental and simulated multiplicity distributions
 - ▶ Include a k_{eff} benchmark (Jezebel)

Objectives

- Reduce discrepancy in multiplicity distributions w/o significantly altering k_{eff}
- Perform energy-dependent perturbations to $\bar{\nu}(E)$ in ^{239}Pu
 - Random samples that preserve covariance data
 - Compare experimental and simulated multiplicity distributions
 - ▶ Include a k_{eff} benchmark (Jezebel)
- Compare $\bar{\nu}(E)$ results to energy-independent shifts of microscopic cross sections σ_i
 - Are these experiments a validation tool for $\bar{\nu}$?

Outline

1. Background
2. Correlated Sampling
3. Methodology
4. Results
5. Conclusions

Covariance and Correlation Matrices

Consider N *dependent* random variables $X_i : i = 1, 2, \dots, N$

- Covariance Matrix Σ :

$$\Sigma_{ij} = \text{Cov}(X_i, X_j)$$

Covariance and Correlation Matrices

Consider N *dependent* random variables $X_i : i = 1, 2, \dots, N$

- Covariance Matrix Σ :

$$\Sigma_{ij} = \text{Cov}(X_i, X_j)$$

- Correlation Matrix C :

$$C_{ij} = \frac{\Sigma_{ij}}{\sqrt{\Sigma_{ii}\Sigma_{jj}}}, \quad C_{ij} \in [-1, 1]$$

Sampling Correlated Random Values

1. Decompose correlation matrix:

$$\mathbf{V}\mathbf{V}^T = \mathbf{C}$$

Sampling Correlated Random Values

1. Decompose correlation matrix:

$$\mathbf{V}\mathbf{V}^T = \mathbf{C}$$

2. Generate vector \mathbf{R} of independent random samples from the [standard normal distribution](#) ($\mu = 0, \sigma^2 = 1$).

Sampling Correlated Random Values

1. Decompose correlation matrix:

$$\mathbf{V}\mathbf{V}^T = \mathbf{C}$$

2. Generate vector \mathbf{R} of independent random samples from the [standard normal distribution](#) ($\mu = 0, \sigma^2 = 1$).
3. Transform \mathbf{R} into vector of correlated samples

$$\tilde{\mathbf{R}} = \mathbf{V}\mathbf{R}$$

Sampling Correlated Random Values

1. Decompose correlation matrix:

$$\mathbf{V}\mathbf{V}^T = \mathbf{C}$$

2. Generate vector \mathbf{R} of independent random samples from the **standard normal distribution** ($\mu = 0, \sigma^2 = 1$).
3. Transform \mathbf{R} into vector of correlated samples

$$\tilde{\mathbf{R}} = \mathbf{V}\mathbf{R}$$

- **Cholesky** decomposition for $\mathbf{V}\mathbf{V}^T$

Outline

1. Background
2. Correlated Sampling
3. Methodology
4. Results
5. Conclusions

Generating Energy-Dependent $\bar{\nu}$ Samples

Generating Energy-Dependent $\bar{\nu}$ Samples

1. Read original $\bar{\nu}(E)$ from **ACE** file and **ENDF/B-VII.1** covariance data
 - Using the LANL **NDVV** python modules

Generating Energy-Dependent $\bar{\nu}$ Samples

1. Read original $\bar{\nu}(E)$ from **ACE** file and **ENDF/B-VII.1** covariance data
 - Using the LANL **NDVV** python modules
2. Sample vector of correlated random numbers $\tilde{\mathbf{R}}$ from \mathbf{C}
 - Each element $\tilde{R}(E_g)$ is # of σ to shift $\bar{\nu}(E)$ in energy group E_g

Generating Energy-Dependent $\bar{\nu}$ Samples

1. Read original $\bar{\nu}(E)$ from **ACE** file and **ENDF/B-VII.1** covariance data
 - Using the LANL **NDVV** python modules
2. Sample vector of correlated random numbers $\tilde{\mathbf{R}}$ from **C**
 - Each element $\tilde{R}(E_g)$ is # of σ to shift $\bar{\nu}(E)$ in energy group E_g
3. Generate new set of $\bar{\nu}$:

Generating Energy-Dependent $\bar{\nu}$ Samples

1. Read original $\bar{\nu}(E)$ from **ACE** file and **ENDF/B-VII.1** covariance data
 - Using the LANL **NDVV** python modules
2. Sample vector of correlated random numbers $\tilde{\mathbf{R}}$ from \mathbf{C}
 - Each element $\tilde{R}(E_g)$ is # of σ to shift $\bar{\nu}(E)$ in energy group E_g
3. Generate new set of $\bar{\nu}$:
 - IF E in E_g :

$$\bar{\nu}'(E) = \sigma_{rel}(E_g)\nu(E)\tilde{\mathbf{R}}(E_g) + \bar{\nu}(E)$$

Fractional shifts to Cross Sections

- Adjust **energy-integrated** cross sections (or $\bar{\nu}$)

$$\sigma_i = \int_0^{E_{max}} \sigma_i(E) dE$$

Fractional shifts to Cross Sections

- Adjust **energy-integrated** cross sections (or $\bar{\nu}$)

$$\sigma_i = \int_0^{E_{max}} \sigma_i(E) dE$$

- Increase cross section by **same fraction** at each energy

$$\sigma_i'(E) = (1 + \alpha)\sigma_i(E) = \sigma_c(E) + \epsilon_i(E)$$

Fractional shifts to Cross Sections

- Adjust **energy-integrated** cross sections (or $\bar{\nu}$)

$$\sigma_i = \int_0^{E_{max}} \sigma_i(E) dE$$

- Increase cross section by **same fraction** at each energy

$$\sigma_i'(E) = (1 + \alpha)\sigma_i(E) = \sigma_c(E) + \epsilon_i(E)$$

- Adjust σ_f and σ_c **separately**

Fractional shifts to Cross Sections

- Adjust **energy-integrated** cross sections (or $\bar{\nu}$)

$$\sigma_i = \int_0^{E_{max}} \sigma_i(E) dE$$

- Increase cross section by **same fraction** at each energy

$$\sigma_i'(E) = (1 + \alpha)\sigma_i(E) = \sigma_c(E) + \epsilon_i(E)$$

- Adjust σ_f and σ_c **separately**
 - Need to compensate for ϵ_c and ϵ_f with σ_t or σ_s

Fractional shifts to Cross Sections

- Adjust **energy-integrated** cross sections (or $\bar{\nu}$)

$$\sigma_i = \int_0^{E_{max}} \sigma_i(E) dE$$

- Increase cross section by **same fraction** at each energy

$$\sigma_i'(E) = (1 + \alpha)\sigma_i(E) = \sigma_c(E) + \epsilon_i(E)$$

- Adjust σ_f and σ_c **separately**
 - Need to compensate for ϵ_c and ϵ_f with σ_t or σ_s
- Compute **average** number of standard deviations shifted $\# s(\sigma_i)$

Comparing Results of Simulations

Comparing Results of Simulations

- Reduced χ^2 values for the 5 multiplicity experiments and criticality benchmark

$$\chi_{red,mult,m}^2 = \frac{1}{N_{bins} - 1} \sum_{i=1}^{N_{bins}} \frac{(P_i^{exp} - P_i^{mcnp})^2}{\sigma^2(P_i^{exp}) + \sigma^2(P_i^{mcnp})}$$

Comparing Results of Simulations

- Reduced χ^2 values for the 5 multiplicity experiments and criticality benchmark

$$\chi_{red,mult,m}^2 = \frac{1}{N_{bins} - 1} \sum_{i=1}^{N_{bins}} \frac{(P_i^{exp} - P_i^{mcnp})^2}{\sigma^2(P_i^{exp}) + \sigma^2(P_i^{mcnp})}$$

- Compute a Cost Function:

$$\text{Cost} = \sum_{m=1}^5 \chi_{red,mult,m}^2 + \chi_{red,k_{eff}}^2$$

Summary of Procedure

- FOR each trial :
 1. Generate a unique set of perturbed nuclear data
 2. Run **MCNP5_mult** simulations (5 multiplicity, JEZEBEL)
 3. Produce multiplicity distributions
 4. Compute χ_{red}^2 values and cost
- The lowest cost is the most accurate trial

Outline

1. Background
2. Correlated Sampling
3. Methodology
4. Results
5. Conclusions

Energy-dependent $\bar{\nu}$ perturbations

- Cost results from 500 trials:

Trial	Cost	$\chi^2_{k_{\text{eff}}}$
$\bar{\nu}$ -1.14%	164.24	33.66
303	197.07	4.18
55	267.9	0.01
Original	426.86	0.27

Energy-dependent $\bar{\nu}$ perturbations

- Cost results from 500 trials:

Trial	Cost	$\chi^2_{k_{\text{eff}}}$
$\bar{\nu}$ -1.14%	164.24	33.66
303	197.07	4.18
55	267.9	0.01
Original	426.86	0.27

- MCNP criticality test suite for [trial 303](#):
 - 39 different ^{239}Pu criticality benchmarks

- $$RMSD = \sqrt{\frac{\sum_{i=1}^{N_{cases}} (k_{\text{eff},i} - k_{\text{eff},i}^{\text{ref}})^2}{N_{cases}}} \times 100\%.$$

Energy-dependent $\bar{\nu}$ perturbations

- Cost results from 500 trials:

Trial	Cost	$\chi^2_{k_{\text{eff}}}$
$\bar{\nu}$ -1.14%	164.24	33.66
303	197.07	4.18
55	267.9	0.01
Original	426.86	0.27

- MCNP criticality test suite for [trial 303](#):


- 39 different ^{239}Pu criticality benchmarks

- $$RMSD = \sqrt{\frac{\sum_{i=1}^{N_{\text{cases}}} (k_{\text{eff},i} - k_{\text{eff},i}^{\text{ref}})^2}{N_{\text{cases}}}} \times 100\%.$$


Trial	<i>RMSD</i>
$\bar{\nu}$ -1.14%	1.23%
303	0.51%
Original	0.49%


Energy-dependent $\bar{\nu}$ perturbations – 3.0 cm HDPE reflector

Original $\bar{\nu}$ Data




Trial 303: Lowest Cost





Energy-dependent $\bar{\nu}$ perturbations – 3.0 cm HDPE reflectorOriginal $\bar{\nu}$ Data

Trial 303: Lowest Cost



- Trial 303 data reduced **bias** in 1st and 2nd moments, averaged over all 5 simulations, by $\sim 35\%$

Simulations of Multiplicity Experiments with Nuclear Data Perturbations

Trial 303 $\bar{\nu}$ Data

Capture Cross Section – 3.0 cm HDPE reflector



Capture Cross Section – 3.0 cm HDPE reflector

- Adjust total cross section (σ_t) to compensate for change in σ_c

Simulations of Multiplicity Experiments with Nuclear Data Perturbations

Capture Cross Section – 3.0 cm HDPE reflector

- Adjust total cross section (σ_t) to compensate for change in σ_c

Original σ_c Data σ_c increased 16%

- Correction is less for other experiments, and $\#s(\sigma_c) = 7\sigma$


Fission Cross Section – 3.0 cm HDPE reflector

- Adjust elastic scattering (σ_s) to compensate change in σ_f , for $E > 1$ keV


Fission Cross Section – 3.0 cm HDPE reflector

- Adjust elastic scattering (σ_s) to compensate change in σ_f , for $E > 1$ keV

Original σ_f Data




σ_f decreased 1.5%




- High accuracy for all simulations: $\sum_{i=1}^5 \chi^2_{red,mult,m} = 14.6$

Energy-Integrated $\bar{\nu}$ Shift – 3.0 cm HDPE reflector

Original $\bar{\nu}$ Data



-1.14% $\bar{\nu}$ Data



Outline

1. Background
2. Correlated Sampling
3. Methodology
4. Results
5. Conclusions

Summary

- Energy-dependent $\bar{\nu}$ perturbations **reduced inaccuracies** in multiplicity while **preserving** k_{eff}
 - Majority of cross-correlation terms $\mathcal{O}(10^{-4})$ or less

Summary

- Energy-dependent $\bar{\nu}$ perturbations **reduced inaccuracies** in multiplicity while **preserving** k_{eff}
 - Majority of cross-correlation terms $\mathcal{O}(10^{-4})$ or less
- Cross Section Results

Summary

- Energy-dependent $\bar{\nu}$ perturbations **reduced inaccuracies** in multiplicity while **preserving** k_{eff}
 - Majority of cross-correlation terms $\mathcal{O}(10^{-4})$ or less
- Cross Section Results
 - Increasing σ_c not effective, relative to uncertainties
 - Increasing σ_f is very effective, as expected

Summary

- Energy-dependent $\bar{\nu}$ perturbations **reduced inaccuracies** in multiplicity while **preserving** k_{eff}
 - Majority of cross-correlation terms $\mathcal{O}(10^{-4})$ or less
- Cross Section Results
 - Increasing σ_c not effective, relative to uncertainties
 - Increasing σ_f is very effective, as expected
- Multiplicity simulations need to be considered in validation of nuclear data, particularly $\bar{\nu}$

Summary

- Energy-dependent $\bar{\nu}$ perturbations **reduced inaccuracies** in multiplicity while **preserving** k_{eff}
 - Majority of cross-correlation terms $\mathcal{O}(10^{-4})$ or less
- Cross Section Results
 - Increasing σ_c not effective, relative to uncertainties
 - Increasing σ_f is very effective, as expected
- Multiplicity simulations need to be considered in validation of nuclear data, particularly $\bar{\nu}$
- Covariance sampling methodology has been developed and demonstrated, with promising results

Summary

- Energy-dependent $\bar{\nu}$ perturbations **reduced inaccuracies** in multiplicity while **preserving** k_{eff}
 - Majority of cross-correlation terms $\mathcal{O}(10^{-4})$ or less
- Cross Section Results
 - Increasing σ_c not effective, relative to uncertainties
 - Increasing σ_f is very effective, as expected
- Multiplicity simulations need to be considered in validation of nuclear data, particularly $\bar{\nu}$
- Covariance sampling methodology has been developed and demonstrated, with promising results
 - Ideally sample all cross sections and $\bar{\nu}$ simultaneously

Questions?

Simulations of Neutron Multiplicity Experiments with
Nuclear Data PerturbationsS.R. Bolding¹, C.J. Solomon²¹ *Texas A&M University, College station, TX*² *Los Alamos National Laboratory, Los Alamos, NM*

ANS National Meeting, 14 November 2013

Backup Slides

¹ *Texas A&M University, College station, TX*

² *Los Alamos National Laboratory, Los Alamos, NM*

Summary of Procedure

Summary of Procedure

- **FOR** each "Trial":
 1. Generate a new set of perturbed nuclear data
 2. Run **MCNP5_mult** simulations (5 multiplicity, JEZEBEL)
 3. Produce multiplicity distributions
 4. Compute χ^2_{red} values and FOM

Summary of Procedure

- FOR each "Trial":
 1. Generate a new set of perturbed nuclear data
 2. Run **MCNP5_mult** simulations (5 multiplicity, JEZEBEL)
 3. Produce multiplicity distributions
 4. Compute χ^2_{red} values and FOM
- The lowest FOM is the most "accurate" trial

Sample Statistics

- Consider independent random samples $\{x_i : i = 1, 2, \dots, N\}$ of a variable X with some PDF $f(x)$
- **Statistics** are some $f(x_1, x_2, \dots, x_N)$
 - **Sample Mean**: $\bar{x} = \frac{1}{N} \sum_{i=1}^N x_i$
 - **Sample Variance**: $s^2 = \frac{1}{N-1} \sum_{i=1}^N (x_i - \bar{x})^2$
- As $N \rightarrow \infty$, approach **population** (true) mean and variance

Energy-dependent $\bar{\nu}$ perturbations

Comparison of Moments

Reflector	Moment	ENDF/B-VII.1 $\bar{\nu}$			Trial 303 $\bar{\nu}$			Experimental	
		Value	σ_{rel}	# σ away	Value	σ_{rel}	# σ away	Value	σ_{rel}
None	1	1.76E+001	2.68E-003	14.11	1.74E+001	2.68E-003	10.13	1.69E+001	1.38E-003
	2	3.31E+002	2.94E-003	24.43	3.24E+002	2.95E-003	17.59	3.08E+002	1.52E-003
0.5	1	2.40E+001	2.67E-003	16.72	2.37E+001	2.67E-003	11.75	2.29E+001	1.51E-003
	2	6.13E+002	2.90E-003	29.51	5.97E+002	2.90E-003	20.84	5.61E+002	1.65E-003
1.0	1	3.17E+001	2.66E-003	23.52	3.11E+001	2.66E-003	16.67	2.97E+001	1.77E-003
	2	1.07E+003	2.89E-003	41.52	1.03E+003	2.89E-003	29.59	9.38E+002	1.93E-003
1.5	1	3.80E+001	2.67E-003	28.61	3.70E+001	2.67E-003	19.27	3.51E+001	1.84E-003
	2	1.54E+003	2.92E-003	50.25	1.46E+003	2.91E-003	34.14	1.32E+003	2.01E-003
3.0	1	3.19E+001	2.70E-003	34.04	3.06E+001	2.70E-003	19.44	2.90E+001	1.75E-003
	2	1.11E+003	3.04E-003	58.05	1.02E+003	3.03E-003	33.72	9.17E+002	1.96E-003

Nuclear Data Formats

• ACE format

```
94239.70c 236.998600 2.53010E-08 08/25/07
94-Pu-239 at 293.6K from endf/b-vii.0 njoy99.248 mat9437
808738 94239 72098 48 45 14 0 6
0 0 0 0 0 0 0 0
1 360491 371402 371450 371498 371546 371594 598924
598970 658265 658310 733847 805945 805959 805973 806478
806492 806492 806506 808735 371781 808738 715724 724200
724211 724253 724259 0 0 0 0 0
1.00000000000E-11 1.03125000000E-11 1.06250000000E-11 1.09375000000E-11
```

• ENDF format

```
7.000000+6 4.519930-9 7.520000+6 0.000000+0 943715102 56
0.000000+0 0.000000+0 0 0 0 0943715 099999
0.000000+0 0.000000+0 0 0 0 09437 0 0 0
9.423900+4 2.369986+2 0 0 0 1943731452 1
0.000000+0 0.000000+0 0 452 0 1943731452 2
0.000000+0 0.000000+0 1 5 1326 51943731452 3
1.000000-5 8.000000-3 1.000000+2 2.000000+2 3.000000+2 5.000000+2 1943731452 4
```

Capture Cross Section - Case 4

- Adjust elastic scattering (σ_s) to compensate for change in σ_c

$$\boxed{\sigma'_s = \sigma_s - \epsilon_c} \quad \epsilon_c = \alpha \sigma_c \quad \sigma'_c = \sigma_c + \epsilon_c$$

- Only for $E > 1keV$

Trial	χ^2_{mult}	χ^2_{keff}	$\#s(\sigma_c)$
$\bar{\nu}$ -1.14%	130.58	33.7	n/a
$\alpha = 10.0\%$	345.1	0.22	4.04
$\alpha = 4.0\%$	390.9	0.01	1.62
$\alpha = 1.0\%$	417.9	0.01	0.40
Original	426.6	0.27	0

Fission Cross Section – Case 1

Trial	χ^2_{mult}	$\chi^2_{k_{eff}}$	$\#s(\sigma_t)$	$s(\#s(\sigma_t))$
-4.0%	1318.2	167.72	-1.16	0.82
-2.0%	101.0	48.31	-0.58	0.41
-1.6%	27.1	22.97	-0.47	0.33
-1.4%	17.4	22.79	-0.41	0.29
-1.2%	23.1	14.25	-0.35	0.25
-1.0%	47.7	9.37	-0.29	0.21
-0.5%	178.7	1.33	-0.14	0.10
$\bar{\nu}$ -1.14%	130.58	33.7	n/a	n/a
Original	426.6	0.27	0	0

Changing Fission and Capture Cross Sections

- Adjust **fission** to compensate for change in **capture**

$$\sigma'_f = \sigma_f - \epsilon_c$$

$$\epsilon_c = -\alpha \sigma_f$$

$$\sigma'_c = \sigma_c + \epsilon_c$$

Trial	χ^2_{mult}	$\#s(\sigma_c)$
$\alpha = 10\%$	90.66	4.31
$\alpha = 4\%$	222.47	1.72
$\alpha = 2\%$	314.82	0.86
$\bar{\nu} - 1.14\%$	130.58	n/a
Original%	426.6	0.0

Changing Fission and Capture Cross Sections

- Adjust **fission** to compensate for change in **capture**

$$\sigma'_f = \sigma_f - \epsilon_c$$

$$\epsilon_c = -\alpha \sigma_f$$

$$\sigma'_c = \sigma_c + \epsilon_c$$

Trial	χ^2_{mult}	$\#s(\sigma_c)$
$\alpha = 10\%$	90.66	4.31
$\alpha = 4\%$	222.47	1.72
$\alpha = 2\%$	314.82	0.86
$\bar{\nu} - 1.14\%$	130.58	n/a
Original%	426.6	0.0

- $\sigma_f \gg \sigma_c$

Effect of altering σ_t

Monoenergetic neutrons, **unit** atom density, **two** reaction types: $\sigma_t = \sigma_a + \sigma_b$

- Perturb σ_a : $\sigma'_a = \sigma_a + \epsilon_a$, $\sigma'_t = \sigma_t + \epsilon_t$ $\sigma'_b = \sigma_b$

- Interaction Probability:

$$\begin{aligned} P(\text{Interaction } i, x) &= P(\text{Interaction}, x) * P(\text{Interaction } i \mid \text{Interaction}, x) \\ &= [1 - e^{-\sigma_t x}] \frac{\sigma_i}{\sigma_t}, \end{aligned}$$

- Probability for σ'_b :

$$P'(\text{Interaction } b) = p'_b(x) = \left[1 - e^{-\sigma'_t x}\right] \frac{\sigma_b}{\sigma'_t}$$

Effect of altering σ_t

- Expand probability for σ'_b with Taylor series:

$$p'_b(x) = \left[1 - (1 - \sigma'_t x + \frac{(\sigma'_t x)^2}{2} + \mathcal{O}(\sigma_t'^3 x^3)) \right] \frac{\sigma_b}{\sigma'_t}$$

$$p'_b(x) = \sigma_b x - \frac{\sigma'_t x^2}{2} - \mathcal{O}(\sigma_t'^2 x^3).$$

- Change in probability:

$$\Delta p_b(x) = p'_b(x) - p_b(x) = -\frac{(\sigma'_t - \sigma_t)x^2}{2} + \mathcal{O}((\sigma_t'^2 - \sigma_t^2)x^3)$$

$$\boxed{\Delta p_b(x) = -\frac{\epsilon_a x^2}{2} + \mathcal{O}((\sigma_t'^2 - \sigma_t^2)x^3)}$$

Simulations

Simulations

- Perform **MCNP5_mult** simulations
 - Use modified nuclear data from created **ACE** files
 - The 5 different Pu sphere **multiplicity** experiments
 - **JEZEBEL** fast critical benchmark

Simulations

- Perform **MCNP5_mult** simulations
 - Use modified nuclear data from created **ACE** files
 - The 5 different Pu sphere **multiplicity** experiments
 - **JEZEBEL** fast critical benchmark
- Generate multiplicity distributions with **mtool.pl** script
 - **Non-paralyzable** dead time correction

Comparing Results of Simulations

Comparing Results of Simulations

- Reduced χ^2 values for the 5 multiplicity experiments and criticality benchmark

$$\chi_{red,mult,m}^2 = \frac{1}{N_{bins} - 1} \sum_{i=1}^{N_{bins}} \frac{(R_i - S_i)^2}{\sigma^2(R_i) + \sigma^2(S_i)}$$

$$\chi_{red,k_{eff}}^2 = \frac{(k_{eff}^{MCNP} - k_{eff}^{ref})^2}{\sigma^2(k_{eff}^{MCNP}) + \sigma^2(k_{eff}^{ref})}$$

Comparing Results of Simulations

- **Reduced** χ^2 values for the 5 multiplicity experiments and criticality benchmark

$$\chi_{red,mult,m}^2 = \frac{1}{N_{bins} - 1} \sum_{i=1}^{N_{bins}} \frac{(R_i - S_i)^2}{\sigma^2(R_i) + \sigma^2(S_i)}$$

$$\chi_{red,k_{eff}}^2 = \frac{(k_{eff}^{MCNP} - k_{eff}^{ref})^2}{\sigma^2(k_{eff}^{MCNP}) + \sigma^2(k_{eff}^{ref})}$$

- Compute a **FOM**:

$$FOM = \sum_{m=1}^5 \chi_{red,mult,m}^2 + \chi_{red,k_{eff}}^2$$

Modifying Nuclear Data

Modifying Nuclear Data

- Covariance data: **ENDF/B-VII.1** library
- MCNP input nuclear data: **ACE** format

Modifying Nuclear Data

- Covariance data: **ENDF/B-VII.1** library
 - **Python** modules at LANL allow reading of ENDF files

- MCNP input nuclear data: **ACE** format

Modifying Nuclear Data

- Covariance data: **ENDF/B-VII.1** library
 - **Python** modules at LANL allow reading of ENDF files
 - Integrated ability to handle *certain* covariance data

- MCNP input nuclear data: **ACE** format

Modifying Nuclear Data

- Covariance data: **ENDF/B-VII.1** library
 - **Python** modules at LANL allow reading of ENDF files
 - Integrated ability to handle *certain* covariance data
 - ▶ Parsing data
- MCNP input nuclear data: **ACE** format

Modifying Nuclear Data

- Covariance data: **ENDF/B-VII.1** library
 - **Python** modules at LANL allow reading of ENDF files
 - Integrated ability to handle *certain* covariance data
 - ▶ Parsing data
 - ▶ Constructing full matrices
- MCNP input nuclear data: **ACE** format

Modifying Nuclear Data

- Covariance data: **ENDF/B-VII.1** library
 - **Python** modules at LANL allow reading of ENDF files
 - Integrated ability to handle *certain* covariance data
 - ▶ Parsing data
 - ▶ Constructing full matrices
 - ▶ Sampling routines
- MCNP input nuclear data: **ACE** format

Modifying Nuclear Data

- Covariance data: **ENDF/B-VII.1** library
 - **Python** modules at LANL allow reading of ENDF files
 - Integrated ability to handle *certain* covariance data
 - ▶ Parsing data
 - ▶ Constructing full matrices
 - ▶ Sampling routines
- MCNP input nuclear data: **ACE** format
 - Built Python modules to read, modify, and rewrite nuclear data

Modifying Nuclear Data

- Covariance data: **ENDF/B-VII.1** library
 - **Python** modules at LANL allow reading of ENDF files
 - Integrated ability to handle *certain* covariance data
 - ▶ Parsing data
 - ▶ Constructing full matrices
 - ▶ Sampling routines
- MCNP input nuclear data: **ACE** format
 - Built Python modules to read, modify, and rewrite nuclear data
- Random samples of $\bar{\nu}(E)$, rather than **linear optimization** (LO)

Modifying Nuclear Data

- Covariance data: **ENDF/B-VII.1** library
 - **Python** modules at LANL allow reading of ENDF files
 - Integrated ability to handle *certain* covariance data
 - ▶ Parsing data
 - ▶ Constructing full matrices
 - ▶ Sampling routines
- MCNP input nuclear data: **ACE** format
 - Built Python modules to read, modify, and rewrite nuclear data
- Random samples of $\overline{\nu}(E)$, rather than **linear optimization** (LO)
 - LO would not preserve statistical accuracy
 - Under-constrained problem, **infinite solutions**

Capture Cross Section

- Adjust **total cross section** (σ_t) to compensate for change in σ_c

$$\sigma'_t = \sigma_t + \epsilon_c$$

$$\epsilon_c = \alpha \sigma_c$$

$$\sigma'_c = \sigma_c + \epsilon_c$$

Capture Cross Section

- Adjust **total cross section** (σ_t) to compensate for change in σ_c

$$\sigma'_t = \sigma_t + \epsilon_c$$

$$\epsilon_c = \alpha \sigma_c$$

$$\sigma'_c = \sigma_c + \epsilon_c$$

Capture Cross Section


- Adjust **total cross section** (σ_t) to compensate for change in σ_c

$$\sigma'_t = \sigma_t + \epsilon_c$$

$$\epsilon_c = \alpha \sigma_c$$

$$\sigma'_c = \sigma_c + \epsilon_c$$

Trial	χ^2_{mult}	χ^2_{keff}	$\#s(\sigma_t)$	$\#s(\sigma_c)$
$\bar{\nu}$ -1.14%	130.6	33.66	n/a	n/a
$\alpha = 16.0\%$	142.6	1.86	3.47	6.90
$\alpha = 8.0\%$	237.5	0.51	1.74	3.45
$\alpha = 2.0\%$	371.2	0.02	0.43	0.86
$\alpha = 1.0\%$	396.4	0.16	0.22	0.43
Original	426.6	0.27	0	0



Capture Cross Section

- Adjust **total cross section** (σ_t) to compensate for change in σ_c

$$\boxed{\sigma'_t = \sigma_t + \epsilon_c}$$

$$\epsilon_c = \alpha \sigma_c$$

$$\sigma'_c = \sigma_c + \epsilon_c$$

Capture Cross Section

- Adjust **total cross section** (σ_t) to compensate for change in σ_c

$$\boxed{\sigma'_t = \sigma_t + \epsilon_c}$$

$$\epsilon_c = \alpha \sigma_c$$

$$\sigma'_c = \sigma_c + \epsilon_c$$

Capture Cross Section


- Adjust **total cross section** (σ_t) to compensate for change in σ_c

$$\sigma'_t = \sigma_t + \epsilon_c$$

$$\epsilon_c = \alpha \sigma_c$$

$$\sigma'_c = \sigma_c + \epsilon_c$$

Trial	χ^2_{mult}	χ^2_{keff}	$\#s(\sigma_t)$	$\#s(\sigma_c)$
$\bar{\nu}$ -1.14%	130.6	33.66	n/a	n/a
$\alpha = 16.0\%$	142.6	1.86	3.47	6.90
$\alpha = 8.0\%$	237.5	0.51	1.74	3.45
$\alpha = 2.0\%$	371.2	0.02	0.43	0.86
$\alpha = 1.0\%$	396.4	0.16	0.22	0.43
Original	426.6	0.27	0	0



Fission Cross Section

- Adjust elastic scattering (σ_s) to compensate for change in σ_f

$$\sigma'_s = \sigma_s + \epsilon_f$$

$$\epsilon_f = -\alpha \sigma_f$$

$$\sigma'_f = \sigma_f - \epsilon_f$$

- Only for $E > 1keV$

Fission Cross Section

- Adjust elastic scattering (σ_s) to compensate for change in σ_f

$$\sigma'_s = \sigma_s + \epsilon_f$$

$$\epsilon_f = -\alpha \sigma_f$$

$$\sigma'_f = \sigma_f - \epsilon_f$$

- Only for $E > 1\text{keV}$

Trial	χ^2_{mult}	χ^2_{keff}
$\alpha = 2.0\%$	65.8	29.6
$\alpha = 1.5\%$	14.6	24.4
$\alpha = 1.0\%$	56.5	9.4
$\alpha = 0.5\%$	195.7	3.0
$\bar{\nu} - 1.14\%$	130.58	33.7
Original%	426.6	0.0

Future Work

- More energy-dependent $\bar{\nu}$ trials

Future Work

- More energy-dependent $\bar{\nu}$ trials
- Energy-dependent σ_f trials

Future Work

- More energy-dependent $\bar{\nu}$ trials
- Energy-dependent σ_f trials
 - σ_f has 400 covariance energy groups vs 50 for $\bar{\nu}$
 - Need constrained global optimization method

Future Work

- More energy-dependent $\bar{\nu}$ trials
- Energy-dependent σ_f trials
 - σ_f has 400 covariance energy groups vs 50 for $\bar{\nu}$
 - Need constrained global optimization method