

# Homework 1

Simon Bolding  
NUEN 629

September 14, 2015

# NUEN 629, Homework 1

Due Date Sept. 17

## 1 Ayzman

(10 points) Henyey and Greenstein (1941) introduced a function which, by the variation of one parameter,  $-1 \leq h \leq 1$ , ranges from backscattering through isotropic scattering to forward scattering. In our notation we can write this as

$$K(\mu_0, v' \rightarrow v) = \frac{1}{2} \frac{1 - h^2}{(1 + h^2 - 2h\mu_0)^{3/2}} \delta(v' - v).$$

Verify that this is a properly normalized  $f(\mu_0)$  and compute  $K_l(v' \rightarrow v)$  for  $l = 0, 1, 2$  as a function of  $h$ .

## 2 Bolding

(20 points) In an elastic scatter between a neutron and a nucleus, the scattering angle in the center of mass system is related to the energy change as

$$\frac{E}{E'} = \frac{1}{2} ((1 + \alpha) + (1 - \alpha) \cos \theta_c),$$

where  $E$  is the energy after scattering and  $E'$  is the initial energy of the neutron and

$$\alpha = \frac{(A - 1)^2}{(A + 1)^2}.$$

The scattered angle in the center-of-mass system is related the lab-frame scattered angle as

$$\tan \theta_L = \frac{\sin \theta_c}{A^{-1} + \cos \theta_c}.$$

Also, the distribution of scattered energy is given by

$$P(E' \rightarrow E) = \begin{cases} \frac{1}{(1 - \alpha)E'}, & \alpha E' \leq E \leq E' \\ 0 & \text{otherwise} \end{cases}.$$

Derive an expression for  $K(\mu_0, E' \rightarrow E)$ , where  $\mu_0$  is  $\cos \theta_L$ . What is the distribution in angle of neutrons of energy in the range  $[0.05 \text{ MeV}, 10 \text{ MeV}]$  to energies below 1 eV if the scatter is hydrogen?

## 3

(70 points) Consider an infinite square lattice of infinitely tall cylindrical UO<sub>2</sub> fuel pins in water. A quarter of a pin cell looks for a square lattice is shown in Fig. 1 and an infinite hex lattice in Fig. 2. The cross-section data for each is given in Table 1. The neutron transport equation for this problem is given simply by

$$\hat{\Omega} \cdot \nabla \psi(x, y, \hat{\Omega}) + \Sigma_t \psi(x, y, \hat{\Omega}) = \frac{1}{4\pi} \Sigma_s \phi(x, y) + \frac{Q}{4\pi}.$$

You may choose whichever lattice you wish – square or hex. For one or the other, perform the following:

Table 1: Data for Test Problem

	Fuel	Moderator
$\Sigma_t$ (cm <sup>-1</sup> )	0.1414	0.08
$\Sigma_s$ (cm <sup>-1</sup> )	0	0
$Q$ (n/cm <sup>3</sup> ·s)	1	0




Figure 1: Quarter of a pin cell of infinite square lattice problem. The azimuthal angle  $\varphi$  is written as  $\gamma$  in the figure.




Figure 2: Pin cell of infinite hex lattice problem.

1. Calculate the angular flux as a function of the azimuthal angle,  $\phi$ , at the spatial points indicated in the figure (two points for the hex lattice; three for the square). Use two different polar angles:  $\pi/2$ , which means the neutrons are traveling in the x-y plane, and  $\pi/8$ . Use the dimensions and cross sections from Table 1. Note that to simplify the problem we have abolished scattering. In the highest energy group of a fine-group set, there is very little within-group scattering, so it does not change the problem very much to ignore scattering. We have also assumed that the neutrons are born uniformly in the fuel – a flat radial profile. This isn't precisely true, but again, the simplification does not change the character of the solution that we wish to study. You will need to write a simple computer program for this in whatever language you'd like. You will need to trace rays and compute points of intersection. When you reach the boundary of a pin cell you use a periodic boundary condition to translate the ray across the cell, and then you continue. You will need a strategy to know how far a ray must be traced before you say "enough." Your code should accept as input:
  - (a) the number of values of the azimuthal angle,  $\phi$  at which to calculate the angular flux;
  - (b) the precision to which to calculate the angular flux at a given spatial point and given  $\phi$ . (This tells you when you can say "enough." You can say "enough" when you've traced through  $\tau$  mean free paths, where  $\exp(-\tau) =$  the requested precision.)
2. Convince me that your code calculates the angular flux correctly.
3. Plot the angular flux as a function of  $\phi$  for each of the two polar angles, for each of the three spatial points (two if hex lattice). Use enough  $\phi$  values to convince yourself that you have resolved all the significant bumps and wiggles in the angular flux. Discuss your plots, and in particular compare them against what was shown in the notes for square pins. Do the circles make things smoother? Be prepared to present your solutions to the class, and (see part above) be prepared to argue that they are correct.

**Problem 1:**

Henyey and Greenstein (1941) introduced a function which, by the variation of one parameter,  $1 \leq h \leq 1$ , ranges from backscattering through isotropic scattering to forward scattering. In our notation we can write this as

$$K(\mu_0, v' \rightarrow v) = \frac{1}{2} \frac{1 - h^2}{(1 + h^2 - 2h\mu_0)^{3/2}} \delta(v' - v). \quad (1)$$

Verify that this is a properly normalized  $f(\mu_0)$  and compute  $K_l(v' \rightarrow v)$  for  $l = 0, 1, 2$  as a function of  $h$ .

**Solution:**

# Solution:

- First, expand scattering Kernel in Legendre Polynomials.
- Begin w/ Legendre generating fn:

$$\frac{1}{\sqrt{1-2xh+h^2}} = \sum_{n=0}^{\infty} P_n(x) h^n$$

(1) [wiki]

- Take a derivative w.r.t.  $x$

$$\frac{h}{(1-2xh+h^2)^{3/2}} = \sum_{n=0}^{\infty} P'_n(x) h^n$$

(2)

- Multiply both sides by  $\frac{(1-h^2)}{2h}$ :

$$K(x) = \frac{1}{2} \frac{(1-h^2)}{(1-2xh+h^2)^{3/2}} = \frac{1}{2} \sum_{n=0}^{\infty} P'_n(x) h^{n-1} (1-h^2)$$

(3)

- (3) is our scattering kernel. Now we need to eliminate the derivative in terms of  $P_n(x)$ . Rewrite sum by letting  $n \rightarrow n+1$

$$\frac{1}{2} \sum_{n=0}^{\infty} P'_n(x) h^{n-1} (1-h^2) = \frac{1}{2} \sum_{n=0}^{\infty} P'_{n+1}(x) h^n (1-h^2)$$

(4)

- Note that the sum still starts at zero because  $P'_0(x) = 0$ , so we can trivially add this term. From Abramowitz:

$$P'_{n+1}(x) = \sum_{m \text{ even}}^n (2m+1) P_m(x)$$

(5)

- (5)  $\rightarrow$  (4) and write as two sums:

$$K(x) = \frac{1}{2} \sum_{n=0}^{\infty} \sum_{m \text{ even}}^n (2m+1) P_m(x) h^n - \frac{1}{2} \sum_{n=0}^{\infty} \sum_{m \text{ even}}^n (2m+1) P_m(x) h^{n+2}$$

(6)

\* Note: Here orthogonality is defined as  $\int P_m P_n = \frac{\delta_{mn} 2}{2n+1} *$

• Shift the second infinite sum by  $n' = n+2$

$$K = \frac{1}{2} \sum_{n=0}^{\infty} \sum_{\substack{m \\ \text{even}}}^n (2m+1) P_m(x) h^n - \frac{1}{2} \sum_{n=2}^{\infty} \sum_{\substack{m \\ \text{even}}}^{n-2} (2m+1) P_m(x) h^{n'}$$

• Combine the two series, writing out  $n=0,1$  & letting  $n' \rightarrow n$

$$K = \frac{1}{2} \sum_{n=0}^1 (2n+1) P_n(x) h^n + \frac{1}{2} \sum_{n=2}^{\infty} \left( \sum_{\substack{m \\ \text{even}}}^n (2m+1) P_m(x) - \sum_{\substack{m \\ \text{even}}}^{n-2} (2m+1) P_m(x) \right) h^n$$

• For the inner sums all terms but  $n$  cancel, adding back energy:

$$K(\mu_0, v' \rightarrow v) = \frac{1}{2} \sum_{n=0}^{\infty} (2n+1) P_n(\mu_0) h^n \delta(v'-v) \quad (7)$$

• Taking Legendre moments:

$$K_e(v' \rightarrow v) = \frac{\delta(v'-v)}{2} \sum_{n=0}^{\infty} (2n+1) h^n \int_{-1}^1 d\mu_0 P_n(\mu_0) P_e(\mu_0)$$

$$K_e(v' \rightarrow v) = \delta(v'-v) \sum_{n=0}^{\infty} \left( \frac{2n+1}{2} \right) h^n \frac{\delta_{ne} 2}{(2n+1)}$$

$$K_e(v' \rightarrow v) = h^e$$

• Noting  $\int_0^{\infty} dv \int_{-1}^1 d\mu_0 = \int_0^{\infty} dv \int_{-1}^1 P_0(\mu_0) d\mu_0 = \int_0^{\infty} dv \int_{-1}^1 d\mu_0 K(\mu_0, v' \rightarrow v) = h^0 = 1$ ,  
so normalization is correct.

$$\begin{aligned} K_0 &= 1 \\ K_1 &= h \\ K_2 &= h^2 \end{aligned}$$

**Problem 2:**

In an elastic scatter between a neutron and a nucleus, the scattering angle in the center of mass system is related to the energy change as

$$\frac{E}{E'} = \frac{1}{2} ((1 + \alpha) + (1 - \alpha) \cos \theta_c) \quad (2)$$

where  $E$  is the energy after scattering and  $E'$  is the initial energy of the neutron and

$$\alpha = \frac{(A - 1)^2}{(A + 1)^2}. \quad (3)$$

The scattered angle in the center-of-mass system is related the lab-frame scattered angle as

$$\tan \theta_L = \frac{A \sin \theta_c}{1 + A \cos \theta_c} \quad (4)$$

Also, the distribution of scattered energy is given by

$$P(E' \rightarrow E) = \begin{cases} \frac{1}{(1-\alpha)E'} & E'\alpha \leq E \leq E' \\ 0 & \text{otherwise} \end{cases}. \quad (5)$$

Derive an expression for  $K(\mu_0, E' \rightarrow E)$ , where  $\mu_0$  is  $\cos \theta_L$ . What is the distribution in angle of neutrons of energy in the range [0.05 MeV, 10 MeV] to energies below 1 eV if the scatter is with hydrogen?

**Solution:****Scattering Kernel Derivation**

Due to Eq. (2), for a fixed  $A$ , a given value of  $E$  and  $E'$  fully define  $\mu_c$ ; the lab frame cosine of the scattering angle  $\mu_0$  is also fully defined through Eq. (4). As a result, the shape of the doubly differential scattering cross section is fully defined by the probability density function (PDF)  $P(E' \rightarrow E)$ . Thus, it is possible to write the scattering cross section in the COM frame as [1]

$$\Sigma_s(\mu_0, E' \rightarrow E) = \Sigma_s(E') P(E' \rightarrow E) \delta(\mu_c - f_\mu(E, E')) \quad (6)$$

where  $f_\mu(E, E')$  is the value of  $\mu_c$  that satisfies Eq. (2) for a given  $E$ , i.e.,

$$f_\mu(E, E') = \frac{2(\frac{E}{E'}) - (1 + \alpha)}{(1 - \alpha)} \quad (7)$$

Because we are interested in the scattering kernel as a function of the lab frame cosine  $\mu_0$ , we define the scattering cross section in an equivalent form

$$\Sigma_s(\mu_0, E' \rightarrow E) = \Sigma_s(E') P(\mu_0) \delta(E - f_E(\mu_c(\mu_0), E')) \quad (8)$$



where  $P(\mu_0)$  is a PDF for  $\mu_0$  given a certain value of  $E'$ ,  $f_E$  is defined as

$$f(\mu_0, E') = \frac{E'}{2} ((1 + \alpha) + (1 - \alpha)\mu_c), \quad (9)$$

and  $\mu_c$  as a function of  $\mu_0$  will be derived later in Eq. (21). The scattering kernel is defined as

$$K(\mu_0, E' \rightarrow E) = \frac{\Sigma_s(E' \rightarrow E, \mu_0)}{\int_0^\infty dE \int_{-1}^1 d\mu_0 \Sigma_s(E' \rightarrow E, \mu_0)} \quad (10)$$

The denominator is evaluated as

$$\int_{-1}^1 d\mu_0 \int_0^\infty dE \Sigma_s(E') P(\mu_0) \delta(E - f_E(\mu_c(\mu_0), E)) = \Sigma_s(E') \int_{-1}^1 d\mu_0 P(\mu_0) = \Sigma_s(E') \quad (11)$$

where the first equality is true because the argument of the delta function is zero for the value of  $\mu_0$  and  $E'$  that satisfy  $f$ , which in this case gives the  $\mu_0$  that is the integration variable of the outer integral. The scattering Kernel is then just

$$K(\mu_0, E' \rightarrow E) = P(\mu_0) \delta(E - f_E(\mu_c(\mu_0), E)). \quad (12)$$

We now need to transform the PDF  $P(E' \rightarrow E)$  into a density function  $P(\mu_0)$ . From Eq. (2), there is a one-to-one relationship between  $E$  and  $\mu_c = \cos(\Theta_c)$  in the range of  $E \in [\alpha E', E']$ , thus

$$P(E' \rightarrow E) dE = P(\mu_c) d\mu_c \quad (13)$$

or

$$P(\mu_c) = P(E' \rightarrow E) \frac{dE}{d\mu_c}. \quad (14)$$

Multiplication of Eq. (2) by  $E'$ , followed by differentiation, yields

$$\frac{dE}{d\mu_c} = \frac{1}{2}(1 - \alpha)E' \quad (15)$$

Evaluating  $\mu_c$  for  $E$  at the limits  $\alpha E'$  and  $E'$  gives the support for  $P(\mu_c)$ , defined for  $\mu_c \in [-1, 1]$ . Substitution of the above equation and Eq. (5) into Eq. (14) gives the PDF in the COM frame

$$P(\mu_c) = \frac{1}{(1 - \alpha)E'} \left( \frac{1}{2}(1 - \alpha)E' \right) = \frac{1}{2}, \quad \mu_c \in [-1, 1] \quad (16)$$

We must now transform to the lab frame scattering cosine  $\mu_0$ . First, we solve Eq. (4) for  $\mu_0$  in terms of  $\mu_c$  as follows:

$$\tan^2 \theta_L = \left( \frac{A \sin \theta_c}{1 + A \cos \theta_c} \right)^2 \quad (17)$$

$$\sec^2 \theta_L - 1 = \left( \frac{A \sin \theta_c}{1 + A \cos \theta_c} \right)^2 \quad (18)$$

$$\mu_0^{-2} = \frac{A^2(\sin^2 \theta_c + \cos^2 \theta_c) + 1 + 2A\mu_c}{(1 + A\mu_c)^2} \quad (19)$$

$$\mu_0 = \frac{1 + A\mu_c}{\sqrt{1 + 2\mu_c A + A^2}}. \quad (20)$$

Solution of the above equation for  $\mu_c$  in terms of  $\mu_L$  gives

$$\mu_c = -\frac{1}{A}(1 - \mu_0^2) + \mu_0 \sqrt{1 - \frac{1}{A^2}(1 - \mu_0^2)}. \quad (21)$$

Eq. (20) demonstrates a one-to-one relationship between  $\mu_0$  and  $\mu_C$ . As before,

$$P(\mu_0) = P(\mu_C(\mu_0)) \frac{d\mu_c}{d\mu_0}. \quad (22)$$

Differentiation of Eq. (21) with respect to  $\mu_0$  and algebraic manipulation ultimately yields

$$\frac{d\mu_c}{d\mu_0} = \frac{2\mu_0}{A} + \frac{1 - \frac{1}{A^2}(1 - 2\mu_0^2)}{\sqrt{1 - \frac{1}{A^2}(1 - \mu_0^2)}}. \quad (23)$$

Substitution of the above equation and Eq. (16) into Eq. (22) gives an expression for  $P(\mu_0)$ . The final expression for the scattering kernel is, for  $A > 1$

$$K(\mu_0, E' \rightarrow E) = \begin{cases} \left[ \frac{\mu_0}{A} + \frac{1 - \frac{1}{A^2}(1 - 2\mu_0^2)}{2\sqrt{1 - \frac{1}{A^2}(1 - \mu_0^2)}} \right] \delta(E - f_E(\mu_c(\mu_0), E')), & \mu_0 \in [-1, 1] \\ 0, & \text{otherwise} \end{cases} \quad (24)$$

where the support is from evaluation of Eq. (20) at  $\mu_c = -1, 1$ . The case of  $A = 1$  must be treated separately. This can be seen, for instance, because evaluation of Eq. (20) at  $\mu_c = -1$  results in an indeterminate  $0/0$ . Evaluation of Eq. (21) for  $A = 1$  gives a non-indeterminate expression for  $\mu_0$  as

$$\mu_0 = \sqrt{\frac{1 + \mu_c}{2}} \quad (25)$$

Thus, the support becomes  $\mu_0 \in [0, 1]$ . The kernel also simplifies significantly at  $A = 1$ . The final scattering kernel, for the case of  $A = 1$ , is

$$K(\mu_0, E' \rightarrow E) = \begin{cases} 2\mu_0 \delta(E - f_E(\mu_c(\mu_0), E')), & \mu_0 \in [0, 1] \\ 0, & \text{otherwise} \end{cases} \quad (26)$$

which is PDF normalized over  $\mu_0$  and  $E$ .

## Plots for A=1


To plot in terms of energy, we evaluate Eq. (2) at  $A = 1$ , giving

$$\frac{E}{E'} = \frac{1 + \mu_c}{2}. \quad (27)$$

Then, using Eq. (25),  $\mu_0$  in terms of  $E$  and  $E'$  is

$$\mu_0 = \sqrt{\frac{E}{E'}}. \quad (28)$$

A plot of  $P(\mu_0)$  vs  $\mu_0$  and  $P(\mu(E', E))$  vs  $E'$  are given below. The values of  $E'$  range between 0.05 MeV to 10 MeV, with  $E$  fixed at 1 eV. As expected, lower energy neutrons are more likely to scatter to 1 eV.



## References

- [1] W.L. Dunn and J.K. Shultis, *Exploring Monte Carlo Methods*, 2012.