

# Which constitutive model do measured cavitation records prefer?

Spencer H. Bryngelson

August 8, 2026

## Abstract

We compare seventeen constitutive models against laser-induced cavitation records in gelatin at three temperatures. The method is Bayesian model selection, with a noise scale measured from trial-to-trial spread and a prior that prices a model down where it merely reproduces a simpler one it contains.

A quadratic Zener solid – strain-stiffening elasticity *and* stress relaxation – wins at every temperature. It reaches  $\chi^2/N = 1.23$  to  $1.35$  on the selection grid, and  $0.96$  refitted to its true optimum at  $15^\circ\text{C}$ . Neither ingredient suffices alone: at that temperature relaxation without stiffening reaches  $1.98$ , and both memoryless candidates land at  $20$ , where they fit each other to one part in a thousand. The stiffening term does no work at all without a relaxation mode to act on. Its advantage collapses as the gel warms, the clearest physical signal in the comparison.

We report the limitations plainly. The residuals are ceilings rather than estimates, because the winning model’s best-fitting region is partly unintegrable. More importantly, a good  $\chi^2/N$  overstates how well the model does. The residual has lag-one autocorrelation  $0.918$ , so the  $201$  samples carry the information of about ten independent ones, and what remains is structure the model lacks rather than noise it failed to average away. The ranking survives this – a  $3\%$  change under a correlated likelihood – but the absolute claim does not.

We then ask what experiment would do better. Scored by parameter precision and by model discrimination, the criteria disagree, and both prove delicate: a criterion defined by an inner minimum is multimodal here, and every improvement to the optimizer lowers it.

Reformulating over design *measures* rather than single designs makes the problem convex, and the Kiefer–Wolfowitz equivalence theorem then certifies the answer. Over  $480$  candidates the optimum splits the runs three ways and beats the best single design by a D-efficiency factor of  $5.74$  – certified globally optimal at a gap of  $9 \times 10^{-10}$ , not merely the best found.

**Keywords:** inertial microcavitation rheometry; model selection; Bayesian evidence; identifiability; optimal experimental design

## 1 Introduction

Inertial microcavitation rheometry infers the mechanical response of a soft material from the radius history of a single collapsing bubble [1]. A laser generates the bubble, high-speed imaging records  $R(t)$ , and a constitutive model is fitted to that trace. The strain rates reached, above  $10^3 \text{ s}^{-1}$ , are inaccessible to conventional rheometry, which is what makes the technique attractive.

The inference rests on choices that are rarely varied. A constitutive law is selected, a bubble-dynamics equation is assumed, thermal transport is included or neglected, and the resulting parameters are reported as material properties. Each choice conditions the answer, and the conditioning is seldom quantified.

We take the model comparison itself as the object of study. Candidate models are ranked by Bayesian evidence rather than by fit, since the candidates nest and best fit therefore selects the most flexible one [2]. We then ask what the records can settle, which turns out to be considerably less than the reported parameters suggest.

Three results follow. First, the likelihood depends on the product  $g\alpha$  alone. The shear modulus is then set by where the prior box cuts a ridge, not by the data, and moving the box swings the model ranking by fifty nats.

Second, the bubble-dynamics operator – held fixed at Keller–Miksis in every published comparison we are aware of – is beaten on all three records by Gilmore/Tait. Third, the correlated residual that motivates richer




Figure 1: Where this work sits in the measurement chain. A laser nucleates a bubble, imaging records how its radius changes, and a material model is fitted to that curve. The fitted stiffness and viscosity are then quoted as material properties. This paper concerns the two right-hand boxes: which model to believe, and which experiment to run so that the question can be answered. Ranking on goodness of fit alone does not work here, because a model with more parameters always fits at least as well as the simpler model inside it.

constitutive models survives everything we tried against it: a second relaxation time, a rate-dependent one, fitting individual trials, and all six operators.

We also ask what experiment would settle the question. Treat the choice of model as one more unknown to be fitted, and one calculation says both how well an experiment separates the models and how well it pins the material. The answer comes with a proof of optimality – unusual for a design calculation, and following from a classical result [3].

Judged that way, the experiments analyzed here are near the best possible for the hardest of the three choices, and 24 to 41 times worse than the best for the two easiest.

## 2 Records and noise model

Each dataset is a set of repeated laser-induced cavitation events at one temperature, recorded as  $R(t)/R_{\max}$ . Figure 2 shows the mean trace and the trial-to-trial spread. We use that spread as the measurement noise. It is measured rather than assumed, so  $\chi^2/N \approx 1$  means a model reproduces the record to within experimental repeatability.




Figure 2: Measured records. Line is the mean over trials, band is  $\pm$  one standard deviation across trials at each time. The spread is the noise model.

Two samples are excluded before fitting. A trial reading above its own maximum radius is a tracking failure. And the  $t = 0$  sample, where every trial agrees exactly, is a definition rather than a measurement: giving it an error bar would invent an uncertainty and let it constrain the fit.

Each candidate is evaluated on a log-spaced grid over its own parameters, at one resolution for every model, so grid luck cannot decide the winner. The evidence is a grid quadrature over that box. It carries three further pieces: the noise scale marginalized under a half-Cauchy prior, a redundancy factor down-weighting parameters where a model merely reproduces a simpler model it contains, and a BIC-style complexity penalty.

Trajectories are compressible [4]. The incompressible approximation changes the residuals by a factor of two to four and must not be used for these records.




Figure 3: The Laplace expansion, and why the evidence is summed over modes. The integrand  $e^{\ell}\pi$  is expanded about each maximum as a Gaussian, whose height and width give that mode’s contribution  $Z_k$  in closed form; the evidence is  $\sum_k Z_k$ , computed as a `logsumexp` over the distinct endpoints a multistart finds. Taking only the best mode discards the rest of the integral, which is the difference between an integral criterion and a minimum: a missed mode biases the total slightly, where a missed basin corrupts a minimum entirely. This is why `fit_candidate` returns every distinct endpoint rather than its winner.

### 3 Method

Four objects carry the analysis. Each is derived here rather than quoted, because the choices between them are not neutral and two of the errors this study corrected came from treating them as interchangeable.

#### 3.1 The likelihood, and what it assumes

Each record gives a mean radius  $y_i$  at time  $t_i$  over 18 trials, with the trial-to-trial standard deviation  $s_i$  as the noise estimate. Modeling the observations as independent and Gaussian about the prediction  $m_i(\theta)$ ,

$$p(y \mid \theta) = \prod_i \frac{1}{\sqrt{2\pi} \sigma_i} \exp \left[ -\frac{(y_i - m_i(\theta))^2}{2\sigma_i^2} \right], \quad -2 \log p = \chi^2(\theta) + \sum_i \log(2\pi\sigma_i^2). \quad (1)$$

Three assumptions are buried here and each is separately checkable. *Gaussianity* is the weakest claim, and is reasonable because  $y_i$  is a mean over 18 trials. *Known  $\sigma_i$*  is not innocuous: using the sample spread as the true deviation understates uncertainty. Section 3.2 handles it by giving the noise scale a prior and integrating it out. *Independence* is the one that fails: the diagonal covariance in (1) says each residual carries fresh information, and Section 13.1 measures a lag-one autocorrelation of 0.90.

The deviations are not constant across the record. Optical tracking degrades where the wall moves fastest, so the variance is inflated by a logistic weight in the strain rate,  $\sigma_i^2 = s_i^2/w_i$  with  $w = w_0 + (1 - w_0) \logit^{-1}(\kappa(\dot{\epsilon}_{\text{th}} - |\dot{\epsilon}_i|)/\dot{\epsilon}_{\text{th}})$ . This is a statement about the measurement, not the model, and it is why  $\chi^2/N$  is dominated by the late record where the weights are large.

#### 3.2 Integrating out the noise scale

Rather than trust  $s_i$ , introduce a single scale  $\beta$  multiplying every deviation,  $\sigma_i \rightarrow \beta\sigma_i$ . Substituting into (1),

$$\log p(y \mid \theta, \beta) = -\frac{\chi^2(\theta)}{2\beta^2} - N \log \beta + \text{const}, \quad (2)$$

which depends on the data only through  $\chi^2$  and  $N$ . That is what makes the marginalization cheap: the forward model never re-enters, so

$$\log p(y \mid \theta) = \log \int_0^\infty p(y \mid \theta, \beta) \pi(\beta) d\beta \quad (3)$$

costs one pass over a quadrature grid rather than one solve per node. With the half-Cauchy prior  $\pi(\beta) = (2/\pi)/(1 + \beta^2)$  the integral is done numerically on  $\beta \in [0.05, 10]$ .

Why marginalize rather than fit  $\beta$ ? Profiling at its best value gives  $-\frac{1}{2}N [1 + \log(\chi^2/N)]$ , and marginalizing is always smaller. The gap is an Occam penalty: a model that fits only by inflating its own error bars is charged for the inflation. Fitting  $\beta$  would hand that flexibility over for free.




Figure 4: Why this study ranks by evidence and not by fit. The constitutive candidates nest – NHKV is qKV at zero stiffening and qSLS at zero relaxation time – so a larger model can never fit worse, and the best-fitting candidate is simply the most flexible one. The evidence adds the Occam factor of Figure 25, which charges a model for prior volume the data does not use, and so turns a monotone quantity into one with an interior maximum. Everything downstream depends on that penalty being computed correctly, which is why the pathology in Figure 25 matters.

### 3.3 Why the evidence, and not the best fit

From Bayes’ theorem, the posterior odds between two models are the prior odds times the ratio of

$$Z_M = p(y \mid M) = \int p(y \mid \theta, M) \pi(\theta \mid M) d\theta. \quad (4)$$

This is the only quantity that answers “which model should I believe” without an arbitrary complexity penalty. Comparing best fits cannot: a nested model never fits worse than the model containing it, so  $\chi^2$  ranks by flexibility.

Information criteria add a penalty –  $2k$  for AIC,  $k \log N$  for BIC. But BIC is an asymptotic approximation to  $-2 \log Z$  assuming a regular, well-identified model with  $N$  large, and Section 3.5 shows this one is not identified in a direction. The integral itself assumes none of that.

The penalty appears explicitly on expanding (4) about the maximum. With  $\hat{\theta}$  the best fit,  $H = -\nabla^2 \log p(y \mid \theta)$  evaluated there, and  $\theta = \hat{\theta} + \delta$ ,

$$Z_M \simeq p(y \mid \hat{\theta}) \pi(\hat{\theta}) \int \exp\left[-\frac{1}{2} \delta^T H \delta\right] d\delta = \underbrace{p(y \mid \hat{\theta})}_{\text{best fit}} \underbrace{\pi(\hat{\theta}) (2\pi)^{p/2} |H|^{-1/2}}_{\text{Occam factor}}. \quad (5)$$

The Gaussian integral is exact for a quadratic  $-\log p$ , so (5) is the Laplace approximation and its error is governed by the third derivative. Reading the second factor as  $\pi(\hat{\theta}) \times (\text{posterior volume})$  makes its meaning plain: it is the fraction of the prior the posterior actually occupies, so a model needing fine tuning pays. This is the sense in which a good fit and a credible model differ, and it is what Freund and Ewoldt [2] argue should replace goodness-of-fit ranking in rheology.

One consequence matters later and is easy to miss. Because  $|H|^{-1/2}$  grows as the posterior broadens, a sloppy direction *raises* the evidence. Model comparison therefore cannot detect non-identifiability – it rewards it – and the two questions must be asked separately.

### 3.4 The prior this study uses

Equation (4) needs  $\pi(\theta \mid M)$ , and a uniform box would make the answer depend on where the box was drawn. Two factors are used instead, both computed on the same grid the quadrature runs over.

The *bottleneck* maps each parameter to  $u_j \in [0, 1]$  in logarithms, then takes the harmonic mean  $H(\theta) = p / \sum_j u_j^{-1}$ . Logarithms because the parameters span decades, where a linear map would encode the bounds rather than the physics.




Figure 5: What lag-one autocorrelation measures, and what it does not. Sampled at the  $N = 201$  points of these records, lag-one is close to a measure of how much of a residual is *smooth*: white noise gives 0.04, a single smooth excursion 1.00, a mixture that is three-quarters smooth 0.92. It does not say what made the residual smooth – model discrepancy, variation between bubbles and a slow calibration drift are alike in this statistic. The qSLS residual gives 0.918; the trial deviations from the mean, with no model fitted at all, give 0.859. Most of the smoothness the fit residual shows is therefore already in the data.

The harmonic mean beats the arithmetic one here precisely because it is dominated by its smallest argument. A parameter driven to the bottom of its range drags the whole prior down. The reasoning: a parameter the data pushed to a boundary is one the model did not need.

The *redundancy* weight asks whether a model merely emulates a simpler one it contains. For a candidate and each nested child, the relative stress mismatch  $F$  is mapped by  $F^n/(F^n + \tau^n)$  against the resolvable scale  $\tau$ . That gives a number near 0 when the parent imitates the child, approaching 1 when it does something genuinely different.

The weight is the *minimum* over children. A model that can imitate any one simpler model is redundant there, however different it looks from the rest.

### 3.5 The Fisher information, and what its eigenvectors are

Differentiate  $-\log p$  twice at fixed data and take the expectation over noise. The term containing  $\partial^2 m$  vanishes because  $\mathbb{E}[y_i - m_i] = 0$ , leaving

$$F_{jk} = \sum_i \frac{1}{\sigma_i^2} \frac{\partial m_i}{\partial \theta_j} \frac{\partial m_i}{\partial \theta_k} = (J^\top J)_{jk}, \quad J_{ij} = \frac{1}{\sigma_i} \frac{\partial m_i}{\partial \theta_j}, \quad (6)$$

so the Fisher information is exactly the Gram matrix of the whitened sensitivities – no approximation, given (1). Near the optimum  $\Delta\chi^2 \simeq \Delta\theta^\top F \Delta\theta$ , so the surface of constant misfit is the ellipsoid  $\Delta\theta^\top F \Delta\theta = c$ . Diagonalizing  $F = V\Lambda V^\top$ , its axes are the eigenvectors and their half-lengths are  $\sqrt{c/\lambda_k}$ : a small eigenvalue is a long axis, a direction along which the data barely object.

The Cramér–Rao bound gives this statistical force. For any unbiased estimator,  $\text{Cov}(\hat{\theta}) \succeq F^{-1}$ , so  $F^{-1}$  is the best achievable covariance and its eigen-decomposition is the confidence ellipsoid. The standard deviation along direction  $k$  is  $\lambda_k^{-1/2}$ , and the ratio  $\kappa = \lambda_{\max}/\lambda_{\min}$  measures how far the problem is from determining all directions equally.

Two conventions are load-bearing. Taking  $\theta = \log(\text{parameters})$  makes an eigenvector a power law: moving along  $v$  sends  $\Delta \log g = v_g t$  and  $\Delta \log \alpha = v_\alpha t$ , so  $g \propto \alpha^{v_g/v_\alpha}$  and the exponent is read directly off the components. Second, sensitivities taken with respect to a scaled coordinate  $u_j = (\log \theta_j - a_j)/w_j$  give  $\Delta \log \theta_j = w_j \tilde{v}_j$ , so the exponent is  $(w_g \tilde{v}_g)/(w_\alpha \tilde{v}_\alpha)$ . With box widths of three and four decades that is a 25% correction. Getting it wrong produced a spurious agreement earlier in this work.

Finally,  $F$  is the observability Gramian of the system augmented with  $\dot{\theta} = 0$ : identifiability and observability are the same statement. The distinction that matters is *structural* unidentifiability, where  $F$  is singular and no data can help, against *practical* sloppiness, where  $F$  is invertible but ill-conditioned. Here  $\lambda_{\min} = 5.2 \times 10^{-2} \neq 0$  and the profile likelihood has a genuine minimum, so the problem is the second – which is why a different experiment can fix it.

### 3.6 What correlated residuals cost

Equation (1) counts  $N$  independent pieces of information. If instead the residuals have autocorrelation  $\rho_k$  at lag  $k$ , then for a sum of  $N$  correlated terms  $\text{Var}(\sum r_i) = N\sigma^2(1 + 2\sum_k(1 - k/N)\rho_k)$ , so the information is

that of

$$N_{\text{eff}} = \frac{N}{1 + 2 \sum_{k \geq 1} \rho_k} \quad (7)$$

independent samples, and parameter uncertainties are understated by  $\sqrt{N/N_{\text{eff}}}$ . The sum is truncated at the first negative  $\rho_k$  – Geyer’s initial positive sequence – because the long tail is estimated from few pairs each and contributes sampling error rather than signal.

The correct remedy is a covariance with off-diagonal terms, not a larger  $\sigma$ . Taking  $C_{ij} = \sigma_i \sigma_j \exp(-|t_i - t_j|/\tau)$ , the Cholesky factor  $C = LL^T$  gives the generalized least squares form

$$-2 \log p = r^T C^{-1} r + \log |C| + \text{const} = \|L^{-1} r\|^2 + 2 \sum_k \log L_{kk} + \text{const}, \quad (8)$$

which is what `FieldObservation` computes. Inflating  $\sigma$  instead would treat structure as noise: it widens the error bars while leaving the parameter estimates biased, which is the failure Brynjarsdóttir and O’Hagan warn about.

### 3.7 Which design criterion, and why they disagree

An experiment is chosen by a scalar summary of  $F$ . The classical family is

$$\text{D: } \max \det F, \quad \text{A: } \min \text{tr } F^{-1}, \quad \text{E: } \max \lambda_{\min}, \quad \text{c: } \min c^T F^{-1} c. \quad (9)$$

D minimizes the volume of the confidence ellipsoid, A its mean squared axis length, E its longest axis, and c the variance of one nominated combination. They rank designs differently because they summarize the same spectrum differently. The choice states what one wants to learn.

For identifiability the relevant fact is a scaling argument. A design change that only amplifies signal – more samples, less noise, a larger bubble – sends  $F \rightarrow sF$ . Then  $\det F \rightarrow s^p \det F$  and  $\lambda_{\min} \rightarrow s \lambda_{\min}$ , so D and E both improve, while

$$\kappa = \frac{\lambda_{\max}}{\lambda_{\min}} \quad \text{is unchanged.} \quad (10)$$

Degeneracy is a property of the *eigenvectors*, not the eigenvalues. Only a design that rotates them can remove it. That is why the E-optimal search preferred a larger bubble while conditioning preferred a gentler collapse: more signal against better separation.

The c-optimal choice carries a small trap. Nominating  $c$  as the sloppy direction *of the present design* optimizes for a direction that moves once the design changes. In this study that criterion recommended the opposite of the right answer.

When the model itself is in doubt, none of these apply: all presuppose the model whose  $F$  is being computed. T-optimality asks instead for the design maximizing the discrepancy between rival predictions,  $\max_{\xi} \|m_1(\hat{\theta}_1; \xi) - m_2(\hat{\theta}_2; \xi)\|$ , and privileges neither.

## 4 Identifiability of the constitutive parameters

The winner fits, but not every parameter it contains is measured by these data.

### Why the trade exists

Write  $\hat{g} = G/p_{\infty} = 1/\text{Ca}$  for the nondimensional modulus and  $\lambda = R_{\text{eq}}/R$  for the stretch the stress law sees. The equilibrium stress the Maxwell arm relaxes toward is

$$Z_e = R^3 \left[ \frac{3\alpha-1}{2} \hat{g} P(\lambda) + 2\alpha \hat{g} Q(\lambda) \right], \quad \begin{aligned} P(\lambda) &= 5 - \lambda^4 - 4\lambda, \\ Q(\lambda) &= \frac{27}{40} + \frac{1}{8}\lambda^8 + \frac{1}{5}\lambda^5 + \lambda^2 - \frac{2}{\lambda}. \end{aligned} \quad (11)$$

Collecting powers of  $\alpha$  separates it into two terms with different parameter dependence,

$$Z_e = \hat{g} \alpha A(\lambda) + \hat{g} B(\lambda), \quad A = R^3 \left[ \frac{3}{2} P + 2Q \right], \quad B = -\frac{1}{2} R^3 P. \quad (12)$$

This is exact; it reproduces the implemented expression to  $2.5 \times 10^{-14}$  over random states. Only the first term carries  $\alpha$ , so  $\hat{g}\alpha$  and  $\hat{g}$  are separately determined only while *both* terms contribute.

They do not, at depth. As  $\lambda$  grows,  $Q \rightarrow \frac{1}{8}\lambda^8$  dominates every other term in either bracket, so  $A \rightarrow \frac{1}{4}R^3\lambda^8$  while  $B \sim \frac{1}{2}R^3\lambda^4$ , and

$$Z_e \rightarrow \frac{1}{4}\hat{g}\alpha R^3\lambda^8, \quad (13)$$

a function of the *product*  $\hat{g}\alpha$  alone. The likelihood cannot then distinguish a soft material that stiffens sharply from a stiff one that barely stiffens. Holding  $Z_e$  fixed in (12) gives the trade in closed form,

$$\frac{\partial \log \hat{g}}{\partial \log \alpha} = -\frac{\alpha A}{\alpha A + B} \rightarrow -1 \quad (\lambda \rightarrow \infty), \quad (14)$$

and the share of the signal that separates the two parameters is

$$\frac{|\hat{g}B|}{|\hat{g}\alpha A|} = \frac{|B|}{\alpha|A|} \simeq \frac{2}{\alpha\lambda^4}. \quad (15)$$

Equations (14) and (15) are what the measurements see. These records collapse to  $\lambda = 2.215$ , where (14) gives  $-0.981$  against  $-0.99$  from the Fisher curvature – a prediction from the constitutive law, not a fit. The profile likelihood returns  $-0.80$  because it averages over a range of  $\lambda$ , and the exponent is  $-0.861$  at  $\lambda = 1.5$ . Separability falls as  $\lambda^{-4}$ : it is 2% of the signal at  $\lambda = 2.2$  and 16% at  $\lambda = 1.5$ .




Figure 6: The two terms of (12) against collapse depth. Left:  $\alpha A$ , which depends on the product  $g\alpha$ , outruns  $B$ , which carries  $g$  alone. Right: their ratio, the share of the signal that can separate the two parameters, following  $2/(\alpha\lambda^4)$  once the collapse is deep. It is 2% at the depth these records reach and 16% at  $\lambda = 1.5$ .

## What the Fisher information measures

For Gaussian observations with known deviations the log-likelihood is  $-\frac{1}{2}\sum_i(y_i - m_i(\theta))^2/\sigma_i^2$  up to a constant, so with the whitened sensitivity  $J_{ij} = \sigma_i^{-1}\partial m_i/\partial\theta_j$  the Fisher information is exactly  $F = J^T J$ , and  $\Delta\chi^2 \simeq \Delta\theta^T F \Delta\theta$ . Taking  $\theta$  to be the logarithms of the parameters makes the eigenvectors read as power-law combinations: along an eigenvector  $v$  the ratio  $v_g/v_\alpha$  is the exponent in  $g \propto \alpha^{v_g/v_\alpha}$ . This is also the observability Gramian of the system augmented with  $\dot{\theta} = 0$ , which is why identifiability and observability are the same statement here.

One caution, since it caught us. Sensitivities taken with respect to a unit cube  $u_j = (\log\theta_j - a_j)/w_j$ , rather than  $\log\theta_j$  directly, give eigenvector components that carry the box widths. Then  $\Delta\log\theta_j = w_j\tilde{v}_j$ , so the exponent is  $(w_g\tilde{v}_g)/(w_\alpha\tilde{v}_\alpha)$ , not  $\tilde{v}_g/\tilde{v}_\alpha$ . With  $w$  of three and four decades the two differ by 25%.




Figure 7: What an eigenvector means, shown rather than defined. The parameters are moved the same distance in log-space along each eigenvector. Along the sloppy direction a 27% change in the modulus, compensated by 38% in the stiffening, leaves the trajectory inside the measurement noise; along the stiff direction a change of the same size is obvious. The data cannot see the first and cannot miss the second.




Figure 8: Left: the 95% region from the inverse Fisher information, 10.7 long and 0.015 wide in log-parameters – an axis ratio of 706, which is why it draws as a line, and the analytic slope of  $-0.981$  lies along it. Right: the misfit measured by re-solving at perturbed parameters, not the quadratic approximation. Moving along the stiff direction costs  $10^4$  in  $\Delta\chi^2$ ; moving the same distance along the sloppy one stays below the 95% threshold, so the parameters may shift by a factor of  $e \approx 2.7$  without the data objecting.



**Modulus and stiffening trade against each other.** Two independent analyses agree. The Fisher information at the fitted optimum has an eigenvalue spectrum spanning six decades. The eigenvector of its smallest eigenvalue is almost purely a shear-modulus-against-stiffening direction: weights  $-0.796$  and  $+0.604$ , against contributions of  $-0.003$  and  $-0.032$  from viscosity and relaxation time. Separately, profiling the likelihood in  $\alpha$  – fixing it and re-optimizing everything else – gives  $g \propto \alpha^{-0.80}$  across two decades. Sampling the posterior gives a third, independent measurement: its widest direction is  $(-0.682, +0.002, -0.082, +0.727)$  in the logarithms of  $(g, \mu, \lambda_1, \alpha)$ , and  $\log g$  is 98.7% anticorrelated with  $\log \alpha$ .

The three exponents are  $-0.99$  from local curvature,  $-0.94$  from the posterior covariance, and  $-0.80$  from the profile. They agree the trade is near-reciprocal and disagree in the third digit. That is expected rather than troubling: the valley is curved, so a curvature estimate at one point and a chord across two decades measure different slopes. The correlation is the more robust statement.

So viscosity and relaxation time are cleanly determined, while modulus and stiffening are constrained only in combination. The identifiable quantity is closer to  $g \alpha^{0.8}$  than to either alone, and a fitted  $\alpha$  should be reported with its correlation to  $g$  rather than as an independent material property.




Figure 9:  $\chi^2/N$  over the modulus–stiffening plane, at the fitted viscosity and relaxation time. The dashed line is  $g \propto \alpha^{-0.80}$ , predicted from the Fisher curvature at a single point; it lies along the floor of the valley. Four “best fits” from four methods are strung out along that floor rather than disagreeing with one another. The white region at upper right is where the model is not physical.

Figure 9 shows why the four estimates in this study disagree about  $g$  by an order of magnitude while agreeing about the fit. They are different points on the floor of one flat-bottomed trench.

**The stiffening is nonetheless identified, once the prior allows it.** Sampling with the ceiling on  $\alpha$  placed at 1, 10 and 100:

| ceiling | log evidence | median $\alpha$ | 97.5% |
|---------|--------------|-----------------|-------|
| 1       | 2162.9       | 0.85            | 0.97  |
| 10      | 2173.9       | 6.37            | 9.04  |
| 100     | 2172.4       | 5.27            | 8.73  |

Widening the prior tenfold moves the median only from 6.37 to 5.27, and the evidence by 1.5 out of 2173. So  $\alpha$  is identified near 5, with 97.5% of posterior mass below about 9. A prior-dominated parameter would

have tracked its bound. The ceiling of 1 shows what genuine truncation looks like: the median pins against it and 11 log units of evidence are lost.

This corrects a natural reading of the profile likelihood alone, which appears flat to the boundary only because the boundary sits where the posterior’s upper tail does.




Figure 10: Left: the Fisher spectrum at the optimum, six decades between the stiffest and sloppiest directions, with the sloppiest being modulus against stiffening. Right: the posterior for  $\alpha$  against the ceiling imposed on its prior. It tracks the ceiling while truncated and departs from it once the ceiling is loose, which is what distinguishes an identified parameter from a prior-driven one.

## 5 Results

Table 1 and Figure 11 give the outcome. The quadratic Zener model qSLS – strain-stiffening elasticity with stress relaxation – wins at every temperature. The ordering beneath it separates into three tiers: models with both ingredients, models with relaxation only, and memoryless models.

Table 1: Best  $\chi^2/N$  on the selection grid by model class – the best node, not the best fit; Section 5.2 refits and reaches 0.962 for qSLS at 15 °C. Values near 1 indicate a fit within experimental repeatability.

|                                     | 15 °C       | 23 °C       | 33 °C       |
|-------------------------------------|-------------|-------------|-------------|
| qSLS (stiffening + relaxation)      | <b>1.35</b> | <b>1.33</b> | <b>1.23</b> |
| SLS (relaxation, linear elasticity) | 2.94        | 1.85        | 1.41        |
| viscoelastic fluids                 | 5.3–5.7     | 4.5–4.7     | 1.9–2.1     |
| memoryless                          | 12.6–16.7   | –           | –           |

Figure 12 overlays the winning model on the data it was selected against.

### 5.1 Reading the posteriors

The model posteriors saturate. The winner takes essentially all the mass and the runners-up are reported as  $10^{-29}$  or smaller. Do not read these magnitudes literally.

Evidence differences scale with sample count. With roughly 3600 samples, a modest per-sample difference in fit compounds into an astronomical Bayes factor. The defensible reading is ordinal: the winner is decisive and the ordering means something, the magnitudes do not. Quote the residual  $\chi^2/N$ .




Figure 11: Best  $\chi^2/N$  on the selection grid for every candidate, all three temperatures, logarithmic scale. The dashed line is the measurement-noise floor.




Figure 12: The selected model against the measured band it was fitted to.

## 5.2 How good are the fits, really?

The table above reports the best  $\chi^2/N$  found on the evaluation grid. Two questions follow, with different answers. How well does the winning model fit? And how much does a good  $\chi^2/N$  entitle us to claim?

**The grid understates the fit.** The model-selection grid evaluates each candidate at log-spaced parameter nodes, so its best value is the best *node*, not the best fit. Refitting each model to the 15 °C record by multistart Gauss–Newton with analytic Jacobians – eight starts from a Latin hypercube over the wide prior box, taking the lowest – gives:

| model | parameters | $\chi^2/N$   | lag-one $\rho$ | $N_{\text{eff}}$ | $\chi^2/N_{\text{eff}}$ | inflation |
|-------|------------|--------------|----------------|------------------|-------------------------|-----------|
| qSLS  | 4          | <b>0.962</b> | 0.918          | 10.3             | 18.9                    | 4.43      |
| SLS   | 3          | 1.984        | 0.932          | 7.1              | 56.6                    | 5.34      |
| qKV   | 3          | 20.18        | 0.948          | 13.4             | 302                     | 3.87      |
| NHKV  | 2          | 20.16        | 0.948          | 13.4             | 302                     | 3.87      |

At its true optimum qSLS reaches  $\chi^2/N = 0.96$  rather than the grid’s 1.35, which confirms in a second way that the tabulated residuals are ceilings. The ordering is unchanged, and the gaps are large: relaxation without stiffening doubles the residual, and both memoryless models sit twenty times above it.

**Stiffening buys nothing without relaxation.** qKV and NHKV differ only by the stiffening exponent  $\alpha$ , and they fit identically: 20.18 against 20.16, one part in a thousand on a residual twenty times too large. The strain-stiffening term is decisive alongside a relaxation mode and worth nothing without one.

That is sharper than the evidence ranking alone. It is not that qKV is a worse model than qSLS; its extra parameter does no work at all. It also explains why the temperature trend is carried by the interaction of the two mechanisms rather than by either alone.

**But  $\chi^2/N \approx 1$  does not mean the fit is within noise.** The statistic  $\chi^2/N$  compares the residual to its expectation under the assumption that the  $N$  residuals are independent draws of known scale. They are not. The lag-one autocorrelation of the qSLS residual is 0.918, and by the Geyer initial-positive-sequence estimator [5] of (7) the 201 samples carry the information of about  $N_{\text{eff}} = 10$  independent ones. Dividing by that count instead gives 18.9.

That number is not a calibrated test. For correlated residuals the correct statistic is  $r^T \Sigma^{-1} r$  with the true covariance, not  $\chi^2$  rescaled by an effective count. But its message is robust, and it is the opposite of the headline’s.

A residual with  $\rho = 0.918$  is not noise the model failed to average away. It is *structure* the model does not contain. The fit is excellent in that no candidate does better, and not excellent in that it is inconsistent with the measurement error.

**The classical test agrees, and says more.** The argument above rests on lag-one autocorrelation, which Section 13.1 shows is a blunt instrument. There is a standard test for the same question and these records already carry what it needs: replicates. The trial spread at each time is *pure error*, costing no model, so the residual of the mean splits into that and what the model cannot follow,

$$F = \frac{J \sum_i (\bar{y}_i - \hat{y}_i)^2 / (k - p)}{(J - 1) \sum_i s_i^2 / k(J - 1)}, \quad (16)$$

with  $J = 18$  trials at  $k = 201$  times and  $p = 4$  parameters.

| record | $\chi^2/N$ | pure error            | lack of fit           | $F$         | corrected   |
|--------|------------|-----------------------|-----------------------|-------------|-------------|
| 15 °C  | 0.973      | $5.20 \times 10^{-4}$ | $7.36 \times 10^{-3}$ | <b>14.2</b> | <b>23.3</b> |
| 23 °C  | 1.017      | $1.86 \times 10^{-3}$ | $5.47 \times 10^{-3}$ | 2.9         | 4.8         |
| 33 °C  | 1.130      | $1.91 \times 10^{-3}$ | $6.23 \times 10^{-3}$ | 3.3         | 5.4         |

At 197 against 3417 degrees of freedom the five percent critical value is near 1.2, so every record shows lack of fit and 15 °C shows it by an order of magnitude. The conclusion is the one above, reached with none of the same machinery.

Two things make it worth the page. It gets the factor of  $J$  right by construction — the fit compares the *mean* of the trials against the spread *between* them, and the  $J$  in the numerator is that missing  $\sqrt{J}$  — and it is conservative in a direction we can quantify: 39.3% of the trial variance is bubble-to-bubble parameter spread rather than measurement, so pure error is inflated and the corrected column divides that share out.

It also separates the records, which lag-one does not. The lag-one values are 0.919, 0.968 and 0.911, effectively identical. But 15 °C repeats itself three times more tightly than the others, so its systematic deviation stands 14 times above its own scatter where theirs stand 3. The strongest evidence that the model is incomplete comes from the record with the best apparatus, not the worst fit.

The  $p$ -value is not usable:  $F$  assumes independent errors across time and these are correlated at 0.918. The ratio needs no such assumption, and is read here as an effect size.

Three things follow, and they should be read together.

1. *The ranking survives.* Refitting under a correlated likelihood with a Cholesky whitening of an estimated AR(1) covariance changes the qSLS–SLS evidence gap by 3%. What the correlation destroys is the absolute claim, not the comparison, and the comparison is what the study is for.
2. *The denominator is the wrong yardstick.* The scale  $\sigma_i$  is the trial-to-trial spread. Much of that is variation in the parameters between bubbles, not error in measuring any one of them. A  $\chi^2$  formed against it is not comparing the model to measurement precision at all.
3. *The misfit is in the early record*, not where the tracking is hardest. Restricting the fit to a later window makes  $\chi^2/N$  worse, 0.97 to 1.95, and does not whiten the residual. The structure cannot be excised by discarding the awkward part of the record.

The honest summary: qSLS is the best available description of these records by a wide margin, it is measurably incomplete, and the size of what is missing is better read from the residual’s correlation than from its magnitude.

### 5.3 What the trial spread actually is

The likelihood of Section 3.2 treats the residuals as independent draws whose scale is the trial-to-trial spread  $s_i$ . Two assumptions are buried in that, and both are wrong in ways that matter for how the numbers above should be read.

**Part of the spread is the parameters, not the measurement.** If the 18 trials differed only by measurement error, their deviations from the trial mean would be white and point nowhere in particular. They do neither.

The singular spectrum of the  $18 \times 201$  deviation matrix puts 53.9% of the variance in one component, where white noise would give 5.9%. Projecting each deviation onto the span of  $G = \partial(R/R_{\max})/\partial\theta$  for the six parameters  $(R_0, R_{\text{eq}}, g, \mu, \lambda_1, \alpha)$  captures 39.3% of the variance against a chance level of  $6/201 = 3.0\%$ . Noise does not align with a six-dimensional subspace of a 201-dimensional space by accident.

The physical reading is that each bubble has its own parameters, drawn from a population:  $\theta_j \sim \mathcal{N}(\theta_{\text{pop}}, \Sigma)$ . The mean trace then has covariance

$$C = \frac{1}{J} \left( G \Sigma G^\top + \sigma_{\text{meas}}^2 I \right), \quad (17)$$

low rank plus diagonal rather than diagonal. Note that  $\Sigma$  itself is not estimable here. Regressing the deviations onto  $G$  inherits the  $g$ – $\alpha$  degeneracy of Section 3.5 and returns  $\text{sd}(\log g) = 159$ . But only the prediction covariance  $G \Sigma G^\top$  enters (17), and that is the sample covariance of the projected deviations, which needs no inversion.

**It is not what makes the residuals correlated.** The natural hypothesis is that the lag-one correlation of the fit residual is this same effect. Whitening by (17) tests it directly:

| covariance             | $\chi^2/N$ | lag-one $\rho$ |
|------------------------|------------|----------------|
| diagonal (independent) | 17.54      | 0.758          |
| hierarchical, (17)     | 21.96      | <b>0.516</b>   |

The correlation falls by about a third, and most of it survives.  $\chi^2$  gets *worse*, so the residual sits in directions where  $C$  is small. The dominant term is model-form error, not variation between bubbles. Both effects are real and the model-error one is larger, which separates two things Section 13.1 had conflated.




Figure 13: What the trial spread is, and what it does not explain. Left: the singular spectrum of the eighteen trial deviations. Independent measurement error would put an equal share in every component (dashed); instead the leading one carries 54%, and 39% of the total lies in the six-dimensional span of  $\partial R/\partial \theta$  against a 3% chance level. Right: the autocorrelation of the qSLS fit residual before and after whitening by the hierarchical covariance (17), with the white-noise band shaded. If the correlation were caused by variation between bubbles, whitening would flatten it into that band. It removes about a third and leaves the rest, so the remainder is structure the model does not contain.

**Which precision  $\chi^2/N$  is measured against.** The same computation exposes a factor of  $J = 18$  in the meaning of the headline number. The data is the *mean* of 18 trials, but  $s_i$  is the spread *between* trials, and the standard error of the mean is smaller by  $\sqrt{18} = 4.24$ . From the identical residual,

$$\chi^2/N = 0.974 \text{ against the trial spread,} \quad \chi^2/N = 17.54 \text{ against the standard error of the mean.}$$

Both are meaningful, and they answer different questions. *The model reproduces the record to within the scatter between individual experiments* is true, and is the claim this document makes. *The model is statistically adequate* is false: against the precision the 18-trial mean actually carries, it is off by a factor of seventeen. The noise-scale marginalization of Section 3.2 absorbs the difference – the implied  $\beta \approx 0.24$  lies inside its prior – so the evidence ranking is undisturbed. What changes is what a residual near one entitles one to say.

**What lag-one does and does not measure.** The inference above – that whitening removes a third of the correlation and the remainder is structure the model lacks – rules out variation between bubbles, and nothing more. Lag-one autocorrelation at  $N = 201$  samples is close to a measure of how much of a residual is *smooth*, not of what made it smooth: sampled at this density, white noise gives 0.04, a single sine period gives 1.00, a smooth random curve gives 0.999, and a mixture that is 80% smooth gives 0.944. A value of 0.918 says the residual is roughly three-quarters smooth structure. It does not say whose.

That matters because the trial deviations from the mean – pure measurement variation, with no model fitted at all – carry lag-one 0.859 (median over the eighteen trials, range 0.725 to 0.890). Most of the smoothness the

fit residual shows is already present in the data before any model is involved. Whitening by the hierarchical covariance removes the part of it that lies along  $\partial R/\partial \theta$ , which is why it removes only a third; the rest is smooth in the data for reasons the parameter sensitivities cannot represent, and model-form error is one candidate among several rather than the established remainder.

The searches of Section 5.4 were run on the stronger reading, and their negative results are consistent with the weaker one.

## 5.4 What to enrich, and how it would have to be judged

The previous section leaves a specific defect: the residual carries structure the model does not contain, and it is not accounted for by variation between bubbles. The natural suspect is the assumption of a *single* relaxation time, which is strong for a crosslinked biopolymer whose relaxation is in reality distributed. Two enrichments follow from that suspicion, and they are not the same claim.

**A second timescale.** `TwoModeQuadraticZener` adds a second Maxwell arm sharing the elastic equilibrium, with a weight  $w$  splitting both the elastic target and the viscous forcing,

$$\dot{Z}_1 = -\frac{Z_1 - (1-w)Z_e}{De_1} + \frac{(1-w)\Phi}{De_1}, \quad \dot{Z}_2 = -\frac{Z_2 - wZ_e}{De_2} + \frac{w\Phi}{De_2}, \quad (18)$$

where  $\Phi$  is the forcing the one-mode law already carries. Splitting *both* terms is what makes  $w = 0$  exact: the second memory is then driven by nothing, decays from zero, and stays there.

**A timescale that moves.** `CarreauZener` keeps one arm and lets its relaxation time depend on the shear rate. A Maxwell arm is a spring and a dashpot in series, so  $\lambda = \eta/G$ ; if the dashpot obeys Carreau rather than Newton,

$$De_{\text{eff}} = De \left[ 1 + (\Lambda \dot{\gamma})^2 \right]^{(n-1)/2}, \quad (19)$$

and the arm relaxes faster the harder the medium is sheared. A collapse spans decades of shear rate, so this is a genuinely different claim from (18) about the same residual. Both terms of the memory equation carry  $De_{\text{eff}}$ , since an arm has one time constant; at  $n = 1$  the bracket is unity at every shear rate and the law is identically the one-mode model.

**Why the grid cannot judge either.** Both carry six free parameters, and the quadrature used throughout is a Cartesian product costing count<sup>p</sup>. At the 12 per axis used throughout, six axes is 5 971 968 solves against 168 072 for the entire standard set. The remedy is to expand about a fit rather than quadrature over a grid, at  $1 + 2p$  solves – but doing so exposes a defect in the expansion itself.

**The Occam factor must be capped at the prior.** Written as in (5), the Laplace evidence contains  $-\frac{1}{2} \log \det J^T J$ . A direction the data does not determine sends an eigenvalue of  $J^T J$  to zero and that term to  $+\infty$ : the expansion *rewards* a model for a parameter its design cannot see. This is not a small correction. On synthetic data generated from the one-mode model, the two-mode model scores

$$+29.7 \text{ nats above the one-mode model at } w = 10^{-3}, \quad -2.6 \text{ at } w = 0.3,$$

so it wins precisely where its second arm does nothing and loses only where the arm does real work. The cause is the fitted Gaussian extending far outside the bounded cube the prior occupies. Capping each eigendirection's contribution at unity – a posterior cannot be wider than the prior it came from, Section A.6 – returns  $-0.0000$ ,  $-0.0004$ ,  $-0.033$  and  $-6.81$  at  $w = 10^{-3}, 10^{-2}, 0.05, 0.3$ : monotone, and never paying for what the data cannot resolve. The correction is a strict generalization, agreeing with the plain form to round-off wherever every direction is already sharper than the prior.

**The answer, on the 15 °C record.** Both were fitted by multistart in the prior’s unit coordinates, scored by the capped expansion, and summed over the modes the multistart found.

| model     | $p$ | $\chi^2/N$ | $\log Z$ | vs. qSLS | lag-one $\rho$ | $N_{\text{eff}}$ |
|-----------|-----|------------|----------|----------|----------------|------------------|
| qSLS      | 4   | 0.962      | 529.26   | –        | 0.918          | 10.3             |
| qSLS2     | 6   | 0.950      | 531.03   | +1.78    | 0.918          | 7.2              |
| qSLStthin | 6   | 0.922      | 532.05   | +2.79    | 0.875          | 8.2              |

The one-mode  $\chi^2/N$  of 0.962 reproduces Section 5.2 exactly, by a different optimizer in different coordinates, which is the check that the fits are comparable at all.

Both enrichments are preferred, and neither margin means what it appears to. The residual autocorrelation is the defect they were reached for, and it does not move: 0.918 unchanged for the second relaxation time,  $0.918 \rightarrow 0.875$  for the rate-dependent one, against  $N_{\text{eff}}$  near 8 out of 201 either way. Both  $\log Z$  and  $\chi^2/N$  presume independent residuals, so the likelihood overstates its own information by roughly  $N/N_{\text{eff}} \approx 25$ ; deflating a 2.79-nat margin by anything of that order leaves nothing. The second arm is not idle – it lands at 16% of the memory with a timescale 15.7 times the first – it simply does not address what is wrong. And qSLStthin pins its crossover at the bottom of its range, so that margin is bound-limited and the crossover is not resolved by this record at all.

**What this says.** A richer relaxation spectrum is not the missing physics, or at least not the part that matters here. Two independent enrichments, each reducing exactly to the one-mode law and each free to switch itself off, both decline to and both leave the correlation in place. That is evidence about where to look next – the residual is concentrated in the first collapse, where sphericity and the far-field boundary are least secure – and it is more useful than a small positive margin would have been.

## 6 The bubble-dynamics operator

Every comparison above varies the material and holds the bubble-dynamics equation at Keller–Miksis. That is an assumption. With the material at its fit, changing only the operator moves the trace by four to fourteen times the median noise, which exceeds any constitutive difference measured here.

The operators are not candidate materials, so they are compared differently. We hold the candidate fixed and vary the forward solve. The parameter space is then identical across the set, the Occam terms cancel, and the difference in  $\log Z$  is a Bayes factor between operators.

**The prior box sets the answer.** Scored with the bounds of Section 3.4, Keller enthalpy/Mie–Grüneisen ranks first at +4.02 nats and Gilmore/Tait second to last at –27.66. Three of the six fits sit on  $g = 10^2$  or  $\lambda_1 = 10^{-7}$ , their own floors. Widening the box to  $g \in (10^0, 10^6)$  and  $\lambda_1 \in (10^{-9}, 10^{-2})$  reverses the ranking: Gilmore/Tait leads at +21.93 and Keller enthalpy/Mie–G falls to +7.32.

Widening moves the pinning rather than removing it. Five of six fits then sit on  $\alpha = 10^3$  with  $g$  collapsed to 1.1 Pa to 3.5 Pa. Figure 26 shows the mechanism: the optimizer slides along the  $g\alpha$  valley until it meets an edge.

**Identified coordinates.** Fixing  $g/\alpha$  and fitting the product  $g\alpha$  removes the pinning entirely, with none of 54 fits on a bound. We repeat at three ratios spanning two decades and on all three records. An ordering counts as settled when one record decides it by more than a nat and none contradicts it.

| ordering                               | 15 °C        | 23 °C       | 33 °C       |
|----------------------------------------|--------------|-------------|-------------|
| compressible > Rayleigh–Plesset        | +162 to +182 | +74 to +79  | +74 to +75  |
| <b>Gilmore/Tait &gt; Keller–Miksis</b> | <b>+17.1</b> | <b>+4.2</b> | <b>+1.2</b> |
| Keller enthalpy/Tait > Keller–Miksis   | +5.8         | +2.0        |             |
| Keller enthalpy/Mie–G > Keller–Miksis  | +4.5         | +1.5        |             |






Figure 14: The size of the operator result, against the size of an error bar this work set to zero. Both arrows are what survives refitting the material – the part of a perturbation that lies outside the shaded span, and so the only part any experiment can see. They are drawn at their measured length and their measured separation. A 1.68 % error in the inferred equilibrium radius leaves exactly as much residual as replacing the bubble-dynamics operator does. The  $58.4^\circ$  between them is the reason this is a call to fit  $R_{\text{eq}}$  rather than to abandon the ranking: at that angle the two are separable in principle, and Section 7 confirms the initial state absorbs under 5 % of the operator difference.

Twelve of the fifteen pairwise orderings are settled. Compressibility is required on every record. The pressure form assumed by every candidate in this package is beaten by three separate operators, and by Gilmore/Tait on all three records.

**A search budget produced an earlier answer.** At six multistart restarts the same comparison placed Gilmore/Tait fourth on two records and first on the third, and reported the operator as undetermined. It also settled no ordering at  $33^\circ\text{C}$ , which we read as a record without discriminating power. At ten starts the  $33^\circ\text{C}$  fits reach  $\chi^2/N = 0.42$  where six starts found 1.151.

Every operator had been held in the same poor basin. A record that cannot decide and a search that has not converged produce the same output. Only a better search separates them.

**The initial radius is not exact.** Every fit above holds  $R_{\text{eq}}$  at its inferred value, which asserts that it is known exactly. It is not:  $R_{\text{max}}$  is read off the images, but  $R_{\text{eq}}$  is inferred, and the margins in the table are small enough that this matters. Perturbing  $R_{\text{eq}}$  alone and projecting off the material span, as in Section 7, an error of 1.68 % leaves the same residual as replacing Keller–Miksis with its Mie–Grüneisen enthalpy form.

That is not a license to ignore the ranking. The two remainders lie at  $58.4^\circ$ , so an initial-radius error and an operator change are separable in principle – an experiment can tell them apart, and Section 7 shows the initial state absorbs under 5 % of the operator difference. What it means is narrower and worse: a quantity we set to zero is the size of the effect being reported. The remedy is to fit  $R_{\text{eq}}$  rather than assert it, which Section 8 does.

The perturbation is superlinear – the residual per unit fractional error runs 79, 93, 138 across half, one and two percent – so 1.68 % is solved for by re-solving rather than scaled from any one of those rows. Read linearly off the 1 % row it would come out a third larger.

## 7 Separability of the model axes

With the constitutive law, the operator and the thermal treatment all uncertain, the question is not which experiment discriminates but whether the axes are separable at all. They need not be, in two ways. A difference lying in the span of  $\partial R/\partial\theta$  is reproduced by refitting the material, and is then invisible at every design (Figure 21). And two axes whose remainders are nearly parallel move the evidence together, so no design can say which moved it.




Figure 15: Turning a model choice into a parameter. Writing  $R(\theta, \varepsilon) = (1 - \varepsilon)R_A(\theta) + \varepsilon R_B(\theta)$  gives  $\partial R / \partial \varepsilon = R_B - R_A$  at  $\varepsilon = 0$ , so the difference between two models becomes one more column of the Jacobian. Discrimination and estimation then live in a single information matrix, and  $D$ -optimality over the augmented set balances learning the material against learning which model produced the trace. Nothing about the criterion changes, so the certificate still applies. The variance of  $\varepsilon$  after the material is fitted out is what says whether a design can tell the models apart at all.




Figure 16: Three choices go into every prediction, and they are made independently: the constitutive law, the bubble-dynamics equation, and whether heat transfer is modeled. Published comparisons vary the first and leave the other two at the origin. Some of each change can be canceled by refitting stiffness and viscosity, which makes it invisible no matter how the experiment is run; what remains is 22% of an operator change, 57% of a constitutive change and 53% of a thermal one, and no two of those remainders point in similar directions. All three are therefore distinguishable in principle.

| axis                                              | size         | after refit | absorbed     | + initial state |
|---------------------------------------------------|--------------|-------------|--------------|-----------------|
| dynamics (KM $\rightarrow$ Keller enthalpy/Mie-G) | 8.89         | <b>1.98</b> | <b>77.7%</b> | 1.89            |
| constitutive (one $\rightarrow$ two modes)        | 14.42        | 8.14        | 43.5%        | 7.72            |
| thermal (cold $\rightarrow$ bubble + medium)      | <b>15.99</b> | 8.46        | 47.1%        | 7.80            |

in units of the record's noise. The last column widens the refit span to include  $R_0$  and  $R_{eq}$ , which are held at their measured values everywhere else in this work. They take a further 4.7, 5.2 and 7.8 % of what the material left, so none of these differences is initial-state freedom in disguise. A near-zero number here is also what a broken projection would report, so the operation is checked both ways on the same vectors: handed the difference itself the basis absorbs it to  $5 \times 10^{-15}$ , handed a random column it absorbs 0.02 of 1.98.

The remainders sit at 46, 71 and 38° to one another. No pair is confounded, so the design problem is well posed for all three. Two things follow that were not expected.

Thermal is the *largest* lever, not a negligible one. It moves the trace most and survives refitting most. So the earlier observation – that switching it on made  $\chi^2/N$  slightly worse – says the direction it moves in is not the one the record wants, not that it is small.

The operator is the most absorbed axis by a wide margin. That is the mechanism behind Section 6: most of an operator change can be mimicked by adjusting the material.

**A design against all three.** Make each axis a parameter through  $R(\theta, \varepsilon) = (1 - \varepsilon)R_A + \varepsilon R_B$ , so  $\partial R / \partial \varepsilon = R_B - R_A$  is one more column of  $J$ . This puts discrimination and estimation in one information matrix and leaves the criterion concave, so the certificate of Figure 17 applies unchanged. The information is averaged over a decade of  $g\alpha$ , because operator information is governed by collapse depth and a measure certified at one material misranks others. The measure certifies at a gap of  $10^{-9}$  on four support points.

| axis         | variance at the optimum         | at the 15 °C geometry | efficiency |
|--------------|---------------------------------|-----------------------|------------|
| dynamics     | $6.78 \times 10^{-3}$ (binding) | $9.91 \times 10^{-3}$ | 68.4%      |
| constitutive | $5.45 \times 10^{-4}$           | $2.24 \times 10^{-2}$ | 2.4%       |
| thermal      | $4.40 \times 10^{-4}$           | $1.05 \times 10^{-2}$ | 4.2%       |

Reported per axis rather than as one determinant. A design maximizing det could buy its whole score on the two easy axes and leave the operator undetermined without saying so. The experiments performed are near-best for the hardest axis and 24 to 41 times off for the two easiest. So a change of geometry buys a great deal of constitutive and thermal discrimination at little cost to the operator. The support wants  $R_{\max}$  near 60  $\mu\text{m}$  at high stretch and near 1200  $\mu\text{m}$  at low stretch, against the 277  $\mu\text{m}$  and stretch 7.09 performed.

## 8 Fitting the initial radius

Section 6 reports operator margins of a few nats while holding  $R_{\text{eq}}$  fixed, and a 1.68% error in  $R_{\text{eq}}$  is worth as much as the operator change those margins measure. A quantity that large cannot be asserted. So we fit it, as a nuisance parameter with a prior. The question is not whether the evidence moves – adding a parameter always moves it – but whether the *ordering* does. An operator preferred by 17 nats with  $R_{\text{eq}}$  pinned, and not preferred once it floats, was never an operator result.

The prior width is the whole argument, and nothing measures it. Too wide and  $R_{\text{eq}}$  absorbs the operator difference, which reads as “the operators are indistinguishable” while being a statement about the box. Too narrow and the pinned answer is reproduced by construction. We therefore sweep the half-width over 3% to 25% rather than choose one. A conclusion that holds across that range is a conclusion; one that flips inside it is a prior.

**A convergence gate with no tuning constant.** Every wider box contains every narrower one, and all of them contain  $R_{\text{eq}}$  at its inferred value, which is the pinned fit. The best attainable  $\chi^2/N$  can therefore only fall as the width grows. A cell that rises has not found its own optimum. This matters because it caught a real failure: at ten multistart restarts, five of 72 cells failed it, one reporting  $-117$  nats between a  $+13$  and a  $+7$  – which reads as a prior effect and is a search that stopped early. At 24 restarts two cells fail, and both are excluded below rather than read as physics.

| operator              | 15 °C  |                | 23 °C  |               | 33 °C  |               |
|-----------------------|--------|----------------|--------|---------------|--------|---------------|
|                       | pinned | $\pm 25\%$     | pinned | $\pm 25\%$    | pinned | $\pm 25\%$    |
| Rayleigh–Plesset      | −165.0 | − <b>178.6</b> | −76.9  | − <b>81.8</b> | −74.4  | − <b>74.2</b> |
| Keller–Miksis         | 0      | 0              | 0      | 0             | 0      | 0             |
| Keller enthalpy/Tait  | +6.1   | +0.8           | +2.1   | +0.1          | +0.8   | +0.8          |
| Gilmore/Tait          | +18.5  | + <b>5.0</b>   | +4.3   | +0.5          | +1.9   | −0.0          |
| Keller enthalpy/Mie–G | +4.7   | +0.3           | +1.6   | +0.1          | +0.6   | +0.9          |
| Gilmore/Mie–G         | +20.7  | + <b>11.2</b>  | +4.3   | —             | +1.1   | −0.3          |

nats against Keller–Miksis; “—” marks the one cell the gate excluded. Of 17 orderings the pinned fit called decisive, 15 stay decisive and keep their sign at every width. None reverses. Two fall below five nats at some width, both at 15 °C.

The conclusions of Section 6 therefore survive, but not equally. *Compressibility is required* at every width and on every record, by 74 to 179 nats – the most robust statement in this work. *Gilmore beats the pressure form* at 15 °C, still by 5 to 11 nats with  $R_{\text{eq}}$  free. But at 23 °C and 33 °C every margin above Keller–Miksis collapses to under a nat. Those two records never had operator discrimination; pinning  $R_{\text{eq}}$  is what made it look as though they did.

**The data want a larger equilibrium radius.** The fitted scale is the same story told forwards, and it is the most consistent number in this study. At  $\pm 3\%$  every one of the 18 cells pins against the upper edge. Given room, they agree closely: 1.104 to 1.115 at 15 °C, 1.074 to 1.086 at 23 °C, and 1.032 to 1.040 at 33 °C, across six operators and two prior widths.

So  $R_{\text{eq}}$  is inferred about 3 % to 11 % too small, and the fit says so whichever operator it is asked through. The residual improves accordingly —  $\chi^2/N$  falls from 0.97 to 0.62 at 15 °C, 1.03 to 0.90 at 23 °C, and 0.44 to 0.38 at 33 °C. That the offset shrinks as the gel warms is a pattern we can measure but not yet explain, and it is larger than the 1.68 % that makes  $R_{\text{eq}}$  worth as much as the operator choice.

## 9 Scoring the thermal treatment, and failing to

Section 7 finds the thermal treatment the largest of the three model axes: it moves the trace by 16 noise units and keeps 8.5 after the material absorbs what it can. It is also the only one never scored. The one number in circulation is a  $\chi^2/N$  that got slightly worse when heat transfer was switched on, which is a fit statistic and not evidence.

Scoring it means running the same comparison as Section 6 with the thermal treatment varied and the material axes held identical, so the Occam terms cancel and the difference is a Bayes factor between physics. We report the attempt because it did not work, and because how it failed is worth knowing.

| record | cold | bubble       | bubble + medium |
|--------|------|--------------|-----------------|
| 15 °C  | 0    | −6.4         | +21.6           |
| 23 °C  | 0    | + <b>2.6</b> | + <b>8.0</b>    |
| 33 °C  | 0    | −5.7         | −66.0           |

nats against the cold model, at 24 multistart restarts. Only the middle row may be read. Repeating the study at ten restarts and again at 24 – a  $2.4\times$  change of budget – moves the 23 °C rows by 0.00, 0.00 and 0.01 nats, and the others by 15, 153, 19 and 68. The cold baselines are stable everywhere, so the record and the material are not at fault; the thermal likelihood surface is.

The decisive detail is that *more restarts fitted worse*. At 15 °C the bubble treatment reached  $\chi^2/N = 0.895$  from ten starts and 1.039 from twenty-four; at 33 °C bubble + medium went from 0.428 to 1.073. Best-of-24 cannot be worse than best-of-10 at a converged optimum, so the search is landing in different basins rather than refining one. A third increase of the budget is therefore not the remedy, and this is the third budget tried.

What this does and does not say. It does not say the thermal treatment is unsupported: on the one record where the fit converges, both treatments are favored, bubble + medium by 8 nats. It says that scoring thermal physics by multistart least squares does not work on these records, and that the fix is a better search – warm-starting each thermal fit from the converged cold optimum, rather than sampling a four-dimensional box afresh. Until then the 15 °C and 33 °C rows above are search noise, and we decline to read the sign of either.

## 10 Experimental design

### Designing without assuming the model

Everything above optimizes the Fisher information of one model at its fitted parameters, which is circular if that model is wrong. Two checks that do not privilege it.

**The record is information-rich; the parameterization is not.** Draw 400 trajectories from the posterior and take the singular values of the centered ensemble. Eleven modes clear the trial noise floor of  $0.018 R_{\max}$ . That is not a count of identifiable combinations: four parameters span at most a four-dimensional manifold, and the extra modes are its curvature. But the record distinguishes far more shapes than the fit extracts. The limit is qSLS, not the measurement.

**Discrimination does not privilege a model.** T-optimality [6] asks a different question: which design makes the candidates disagree most? The criterion is

$$T(d) = \min_{\theta_2} \|m_1(\hat{\theta}_1, d) - m_2(\theta_2, d)\|/\sigma, \quad (20)$$

and the minimization is the whole content of it. A rival that retunes its parameters to imitate has not been discriminated against. So the separation is measured against the rival’s *best imitation*, never against its own fitted values.

Two errors here inflated  $T$ , both for one structural reason.  $T$  is a *minimum*, so anything stopping short of it reports discrimination that is not there. Evaluating each rival at its own fit gave 2.69 for qSLS–SLS at the present design; re-fitting with one Nelder–Mead pass gave 0.67.

Restarting and polishing changed other entries again. At  $1200\ \mu\text{m}$  the reported 4.92 fell to 2.38. A 0.1% change of design moved one loose evaluation from 7.25 to 1.36 – the optimizer’s signature, not the physics. The converged values are:

| design                                       | qSLS–SLS    | qSLS–qKV    | qSLS–NHKV   |
|----------------------------------------------|-------------|-------------|-------------|
| 277 $\mu\text{m}$ , stretch 7.09 (present)   | 0.67        | 4.53        | 4.37        |
| 277 $\mu\text{m}$ , stretch 5.00             | 0.72        | 4.61        | 4.61        |
| 277 $\mu\text{m}$ , stretch 3.00             | 0.74        | 1.72        | 1.72        |
| 199 $\mu\text{m}$ , stretch 3.55 (E-optimal) | 0.56        | 2.59        | 2.59        |
| 895 $\mu\text{m}$ , stretch 4.93             | <b>2.39</b> | 7.13        | <b>7.13</b> |
| 910 $\mu\text{m}$ , stretch 5.19             | 1.36        | 6.97        | 6.97        |
| 1200 $\mu\text{m}$ , stretch 7.09            | 2.38        | 5.03        | 5.63        |
| 100 $\mu\text{m}$ , stretch 20.0             | 1.26        | <b>8.04</b> | 4.17        |
| 57 $\mu\text{m}$ , stretch 19.5              | 1.64        | 5.53        | 5.00        |

At the present design SLS imitates qSLS to within 0.67 noise units, inside the measurement scatter. An unrelated calculation corroborates it. A systematic offset of  $0.668\sigma$  over 201 samples contributes  $201 \times 0.668^2 = 89.7$  to  $\chi^2$ , against a measured log-evidence gap of 101.6. Two computations sharing no machinery, agreeing to 12%.

**These values are upper bounds, and the criterion is the reason.**  $T$  is an inner *global* minimization, and that minimization is multimodal. Nelder–Mead and Gauss–Newton both converge, to different basins. At 895  $\mu\text{m}$  Nelder–Mead returns 2.39 against Gauss–Newton’s 1.52; at 1200  $\mu\text{m}$  the ordering reverses.

Multistart Gauss–Newton finds a lower value in 12 of 27 cells. Analytic Jacobians make it  $44\times$  cheaper than the derivative-free version – 34 s against 25 minutes for the table. A stationarity check from the same Jacobians reports 11 cells still not at a minimum.

Every improvement to the optimizer lowers the scores, because a better-optimized rival imitates better. So the tabulated values are upper bounds on discrimination. The true separations are smaller than any of them.

This is not a defect of the implementation but of the criterion. A minimum is attained at a point, so landing in the wrong basin gives a wrong answer with no indication that anything went wrong, and enumerating the outer design space exactly does not repair it: a dense grid over an unreliable integrand produces a front that is precisely wrong. Any criterion built on an argmin –  $T$ -optimality, profile likelihood, minimax design – inherits this.

The structural fix is to replace the minimum with an integral. Treating the model label as a discrete parameter makes discrimination an expected-information-gain problem over the augmented parameter  $(\theta, m)$ , whose inner quantity is the marginal likelihood  $\int p(Y \mid \theta, M, d) \pi(\theta \mid M) d\theta$ . An integral accumulates over all

modes instead of requiring the best one be found. Multimodality stops being a failure mode and becomes a property of the integrand: a missed mode biases the evidence slightly rather than corrupting the criterion.

It also puts discrimination and parameter precision in one framework. The cost is that the marginal likelihood is prior-sensitive where  $T$  is not. The prior becomes part of the criterion rather than a nuisance.

**The criteria do not agree.** Computed correctly, the criteria want different experiments. E-optimality peaks at  $199\text{ }\mu\text{m}$  and stretch 3.55. Separating qSLS from SLS or from NHKV is best at  $895\text{ }\mu\text{m}$  and stretch 4.93. Separating qSLS from qKV is best at  $100\text{ }\mu\text{m}$  and stretch 20, near the opposite corner of the design space. At the E-optimal design the discrimination scores are the worst in the table, so the conflict is not marginal.

The reason is physical, not numerical. Telling a material with a relaxation mode from one without needs a collapse fast enough to excite it. Separating  $g$  from  $\alpha$  needs a gentle collapse, because the term of  $Z_e$  carrying that distinction dies as  $\lambda^{-4}$ . No design is both fast and gentle.

There is accordingly no single best design, and a scalar design objective is answering a question that has no scalar answer. What replaces it is the *Pareto front*: the set of designs on which no other design is better on every criterion at once. Enumerating a  $14 \times 18$  grid over  $(R_{\max}, \lambda)$  and applying that test leaves 12 designs undominated, spanning  $50\text{--}1200\text{ }\mu\text{m}$  and stretch 4–18. Reporting that set rather than a single optimum is the honest summary of an experiment serving several questions at once; `pyimr.pareto.explore_tradeoff` computes it.

The  $T$  columns entering the domination test are upper bounds, so the front is provisional in its discrimination coordinates. The E-optimality coordinate is an eigenvalue of  $J^T J$  with no inner optimization, and carries no such reservation.

## Discrimination as an integral rather than a minimum

Everything above scores a design by  $T$ , defined through a *global minimum* over the rival’s parameters. That definition is the trouble in the previous subsection, and the trouble is structural rather than computational. Here we replace the minimum with an integral, say why that is more robust, and report what it gives.

**The model label is a parameter.** Write  $M \in \{M_1, M_2\}$  for the model index with prior  $p(M)$ , and let each model carry its own parameter  $\theta$  with prior  $\pi(\theta | M)$ . Nothing distinguishes  $M$  from any other unknown, so the information a design  $d$  yields about it is the mutual information

$$I(M; Y | d) = \mathbb{E}_M \mathbb{E}_{Y|M,d} \left[ \log \frac{p(Y | M, d)}{p(Y | d)} \right], \quad p(Y | d) = \sum_M p(M) p(Y | M, d), \quad (21)$$

the expected reduction in entropy of  $M$  – exactly as the expected information gain above is the expected reduction in entropy of  $\theta$ . Discrimination and estimation are one criterion applied to different components of the augmented unknown  $(\theta, M)$ . They are not competing philosophies: the earlier tension between  $T$ - and  $E$ -optimality is about *which component we care about*, not about two schools.

The quantity entering (21) is the marginal likelihood, or evidence,

$$p(Y | M, d) = \int p(Y | \theta, M, d) \pi(\theta | M) d\theta, \quad (22)$$

the same object as (4) but now at a hypothetical design and for hypothetical data. It is an integral over  $\theta$ , and that single fact is what the rest of this subsection turns on.

**The criterion actually computed.** Rather than (21) we score a design by the expected log Bayes factor in favor of the model that generated the data,

$$U(d) = \mathbb{E}_{Y \sim p(\cdot | M_1, d)} [\log p(Y | M_1, d) - \log p(Y | M_2, d)] = D_{\text{KL}}(p(\cdot | M_1, d) \| p(\cdot | M_2, d)). \quad (23)$$

Recognizing it as a Kullback–Leibler divergence between the two prior-predictive distributions settles its properties. By Gibbs’ inequality  $U(d) \geq 0$ , with equality exactly when the two predictives agree almost everywhere: *a model cannot discriminate against itself, at any design.*

That is sharp and falsifiable, and it is what the implementation is tested against. Two independently drawn banks from one family must score zero; any systematic excess is bias.

Equation (21) relates by  $I(M; Y | d) = \frac{1}{2}[D_{\text{KL}}(p_1 \| \bar{p}) + D_{\text{KL}}(p_2 \| \bar{p})]$  with  $\bar{p}$  the equal mixture. So (23) is the same comparison read against the rival rather than the mixture – the more direct statement of “how strongly will this experiment favor the truth”.

**Why an integral is robust where a minimum is not.** Both criteria require an inner computation that can go wrong. The asymmetry is in how the error propagates.

For  $T$ , any optimizer returns  $\hat{T} = \|m_1 - m_2(\hat{\theta}_2)\|/\sigma$  at whatever point it stopped, and since  $T$  is a minimum,  $\hat{T} \geq T$  always. If  $\hat{\theta}_2$  lies in the correct basin but is imprecise, stationarity gives  $\hat{T} - T = O(\|\hat{\theta}_2 - \theta_2^*\|^2)$  and the error is negligible. If it lies in the *wrong basin*, the error is  $O(1)$ : a fixed, unbounded overstatement with nothing in the returned value to indicate it. This is what was measured – Nelder–Mead and Gauss–Newton returning 2.39 and 1.52 at one design and reversing at another.

For the evidence, suppose the posterior has modes  $k = 1, \dots, K$  contributing  $Z = \sum_k Z_k$ , and a procedure finds only a subset  $\mathcal{S}$ . Then

$$\log \hat{Z} - \log Z = \log \left( 1 - \sum_{k \notin \mathcal{S}} \frac{Z_k}{Z} \right), \quad (24)$$

so the error is controlled by the *posterior weight* of what was missed. Missing a subdominant mode costs  $O(Z_k/Z)$  and missing a negligible one costs nothing. Because the contributions add, a multistart that finds several modes can sum them rather than choose between them, and each additional mode found monotonically improves the estimate. There is no analogue for a minimum, where finding more candidates only ever changes which single one is reported.

This is the whole argument for the change, and it is worth stating plainly: multimodality is a *failure mode* for a minimum and merely a *property of the integrand* for an integral.

**Estimating the evidence I: prior sampling, and why it fails here.** The direct estimator draws  $\theta_s \sim \pi(\theta | M)$  and averages the likelihood,

$$\hat{Z} = \frac{1}{S} \sum_{s=1}^S p(Y | \theta_s, M, d), \quad w_s = p(Y | \theta_s, M, d), \quad (25)$$

which is unbiased for  $Z$ , though  $\log \hat{Z}$  is biased low by Jensen’s inequality. Its reliability rides on the spread of the weights, summarized by the effective sample size

$$S_{\text{eff}} = \frac{(\sum_s w_s)^2}{\sum_s w_s^2} \in [1, S], \quad (26)$$

which equals  $S$  when all draws contribute equally and 1 when a single draw dominates. It costs nothing to compute. It is also the only thing standing between this estimator and a confidently reported number with no content.

That it must fail here can be seen in advance. With  $N$  samples the log-likelihood scales as  $\log p(Y | \theta) \simeq -\frac{1}{2}N\bar{d}(\theta)$  for a mean squared discrepancy  $\bar{d}$ , so weights are  $w_s \propto \exp[-\frac{1}{2}N\bar{d}(\theta_s)]$  and the ratio of the two largest is exponential in  $N$ . Useful draws are those landing inside the posterior. Its width is  $O(N^{-1/2})$  in each of  $p$  directions, so the expected number of useful draws is

$$\mathbb{E}[\#\text{useful}] \sim S \cdot \frac{\text{vol}(\text{posterior})}{\text{vol}(\text{prior})} \sim S N^{-p/2}. \quad (27)$$

For the present study  $N = 201$ ,  $p = 4$  and  $S = 220$ , giving  $220 \times 201^{-2} \approx 5 \times 10^{-3}$  – fewer than one useful draw.

The measurement agrees exactly.  $S_{\text{eff}} = 1.0$  at *every* design tried, with reported values between 793 and 10979 nats that are pure noise. The standard errors ran 56 to 130 throughout and looked entirely respectable. Without (26) the failure is invisible.



**Estimating the evidence II: Laplace.** Expand  $\log[p(Y | \theta)\pi(\theta)]$  to second order about its maximum  $\hat{\theta}$  and do the resulting Gaussian integral. As in (5),

$$\log Z \simeq \log p(Y | \hat{\theta}) + \log \pi(\hat{\theta}) + \frac{p}{2} \log 2\pi - \frac{1}{2} \log \det H, \quad H = -\nabla^2 \log p(Y | \theta)|_{\hat{\theta}}. \quad (28)$$

For Gaussian data  $H = J^T J + \sum_i r_i \nabla^2 r_i$ . Dropping the second term – the Gauss–Newton approximation, valid when residuals are small or the model nearly linear – leaves  $H \simeq J^T J$ , which the fit already computed. Parameterizing so the prior is uniform on the unit cube makes  $\pi(\hat{\theta}) = 1$ , and with whitened residuals  $r = (m - y)/\sigma$ ,

$$\log Z \simeq -\frac{1}{2} \|r\|^2 - \frac{N}{2} \log(2\pi\sigma^2) + \frac{p}{2} \log 2\pi - \frac{1}{2} \log \det(J^T J). \quad (29)$$

The first two terms are the best fit; the last two are the Occam factor, and their sign is what prevents a more flexible model from winning on fit alone. Crucially for the present purpose, (29) does not care how sharp the likelihood is, which is precisely the regime that destroyed (25). It is affordable only because the fit uses analytic Jacobians: derivative-free minimization of the same objective costs 55 s against 7.6 s, and the Jacobian needed for  $\det(J^T J)$  is a by-product of the solve rather than an extra cost.

**The two criteria are consistent where both can be computed.** The criteria are not independent, and the relation between them is a check on both. If the true model fits to within noise,  $\chi_1^2 \approx N$ , while a rival that cannot fit leaves a residual of root-mean-square  $T\sigma$ , so  $\chi_2^2 \approx NT^2$ . Substituting into the leading term of (29),

$$\log \text{BF} \approx \frac{1}{2} (\chi_2^2 - \chi_1^2) \approx \frac{N}{2} (T^2 - 1) \xrightarrow{T \gg 1} \frac{N}{2} T^2, \quad (30)$$

up to the Occam terms, which  $T$  has no way to express. Testing (30) against the measured values:

| design                           | $T$ (vs SLS) | $\frac{N}{2} T^2$ predicted | measured $U$ (nats) |
|----------------------------------|--------------|-----------------------------|---------------------|
| 199 $\mu\text{m}$ , stretch 3.55 | 0.56         | 31.5                        | $35.4 \pm 12.2$     |
| 895 $\mu\text{m}$ , stretch 4.93 | 1.52         | 232                         | $150 \pm 73$        |

Two computations sharing no machinery – one a constrained least-squares minimum, the other a marginal likelihood by Laplace – agreeing to within their stated uncertainty. The residual gap at 895  $\mu\text{m}$  is of the right sign and size for the Occam penalty that qSLS pays for its fourth parameter.

**What it costs, and where it is not yet trustworthy.** Two prices are paid for removing the minimum, and both are real.

*Prior sensitivity.*  $Z$  depends on  $\pi(\theta | M)$  through the normalization of (22). Widening a prior by a factor  $c$  in each of  $p$  directions divides the evidence by roughly  $c^p$  without changing the fit at all. This is the Lindley–Jeffreys effect [7, 8], and it is not a defect to engineer away: a model whose prior is diffuse really is less supported by data that pin it down.

But it makes the prior part of the criterion, where  $T$  needed none, so the prior must be stated. Centering each rival’s prior on its fit to the *existing* record produced a criterion measuring prior placement instead of discriminability, with every Laplace expansion invalid. The results here use the wide, physically motivated box of the model-selection grid, identical across models on every shared parameter.

*Boundary pinning.* Equation (28) presumes an interior maximum. In 50–90% of the fits at each design the rival’s best parameters lie on a face of the prior box, where the expansion does not apply and the value is not to be trusted.

That it happens is itself informative. A rival that can imitate qSLS only by driving a parameter to the edge of physical plausibility has, in a meaningful sense, been discriminated against. But saying so needs a criterion built for a constrained maximum, not a quadratic expansion about an interior one. Which parameter pins, and whether the pinning is physical or an artifact of the box, is the outstanding step.

The honest summary: the integral criterion is better founded than  $T$  and not yet more reliable on this problem. The multimodality that made  $T$  untrustworthy is genuinely gone. What replaces it is a dependence on the prior – now explicit rather than hidden – and a boundary condition the current estimator does not handle.






Figure 17: The certificate, which is what makes an optimal design a proof rather than a stopping rule. Kiefer and Wolfowitz: a measure  $\xi^*$  is  $D$ -optimal *if and only if* its sensitivity  $d(x, \xi^*)$  never exceeds the parameter count  $p$  anywhere in the design space, and attains  $p$  exactly on the points carrying weight. Both halves are visible at once – a curve capped by a level it touches and never crosses – so a search reports not merely that it stopped improving but that there is nowhere left for a better design to hide. The gap  $\max_x d(x, \xi^*) - p$  is the proof’s residue;  $10^{-9}$  in the designs here.




Figure 18: A design is a measure, not a point. Each candidate  $x_i$  carries an information matrix  $M(x_i)$ , and a design assigns weights  $\xi_i$  summing to one, read as the fraction of runs spent at that candidate. The averaged information  $M(\xi)$  is linear in  $\xi$ , so any concave function of it is concave in the design, and the maximization becomes convex. That convexity is what makes Figure 17 possible: a local optimum is global, and a certificate can therefore exist. Weights of zero are the common case, so the support of the optimal measure is usually a handful of designs out of hundreds.

## Which criterion the conflict is between

The scaling argument of Section 3.7 says a design change that merely amplifies signal leaves the condition number  $\kappa$  untouched. Degeneracy is a property of the eigenvectors, and only a design that rotates them can remove it.

For these records, E-optimality preferred a larger bubble ( $895 \mu\text{m}$ ,  $\lambda_{\min} = 12.8$ ). Conditioning and eigenvector composition preferred a gentler collapse at the present size ( $277 \mu\text{m}$  at stretch 5,  $\kappa = 4.2 \times 10^3$ , the modulus-stiffening share of the worst direction falling from 87% to 66%). Both are correct answers to different questions, and (15) says which one matters here.

So there are two distinct disagreements in play, and they have different resolutions. The classical criteria disagree among themselves because they summarize one spectrum differently; that is a choice about what one wants to learn. Estimation and discrimination disagree because they are about different quantities altogether. The next subsection resolves the second, and in doing so dissolves the first.

## Designing over measures, and certifying the answer

Every design result above shares a defect that no amount of care in computing them repairs: each is the output of a search over a single design point, that search is not convex, and nothing in its output distinguishes a good answer from a bad one. The evidence that this is not hypothetical is in the preceding subsections. A surrogate search recovered 16% of the best conditioning actually available and reported the result without qualification; a dense grid over a criterion whose inner minimization was unreliable produced a front that was precise and wrong. Both look exactly like success.

The classical formulation removes the defect rather than mitigating it.




Figure 19: How the optimal design is found. Every candidate starts with equal weight, and each step multiplies a candidate’s weight by how informative it is relative to the current average, then renormalizes. Weights therefore stay positive and keep summing to one by construction, and the criterion increases at every step. Candidates that are not worth running decay to zero, so the answer is a handful of settings out of hundreds. The final weights are the fraction of bubbles to run at each setting, and Figure 17 is what certifies that no other allocation does better.

**Designs as measures.** Instead of choosing one design  $x$  from a candidate set  $\mathcal{X}$ , choose a probability measure  $\xi$  on  $\mathcal{X}$ : run a fraction  $\xi_i$  of the bubbles at setting  $x_i$ . This is not an abstraction imposed for mathematical convenience – it is what an experimenter with a run budget does anyway, and it is strictly more expressive, since a single design is the special case of a measure concentrated on one point. The information available from a run allocated by  $\xi$  is the average

$$M(\xi) = \sum_i \xi_i M(x_i), \quad M(x) = J(x)^\top J(x), \quad (31)$$

because information from independent runs adds and  $\xi$  says how many of each to take.

The consequence is the whole point:  $M(\xi)$  is *linear* in  $\xi$ .

**Concavity, and why the first-order condition is enough.**  $\log \det$  is concave on the positive-definite cone, and the composition of a concave function with a linear map is concave, so  $\Phi(\xi) = \log \det M(\xi)$  is concave on the simplex – a convex set. A concave function on a convex set has no local maximum that is not global, and its first-order condition is not merely necessary but sufficient. Every pathology this study ran into – wrong basins, surrogate blind spots, answers that improve whenever the optimizer does – is a symptom of non-convexity, and none of them can occur here.

The same holds for  $\Phi_A = -\text{tr } M(\xi)^{-1}$  and  $\Phi_E = \lambda_{\min}(M(\xi))$ , and for any non-negative combination of concave criteria.

**The directional derivative.** Move mass  $\alpha$  from  $\xi$  toward the single design  $x$ , giving  $\xi_\alpha = (1 - \alpha)\xi + \alpha \delta_x$ . By linearity  $M(\xi_\alpha) = (1 - \alpha)M(\xi) + \alpha M(x)$ , and using  $\partial \log \det A = \text{tr}(A^{-1} \partial A)$ ,

$$\phi(x, \xi) \equiv \left. \frac{\partial}{\partial \alpha} \log \det M(\xi_\alpha) \right|_{\alpha=0} = \text{tr}[M(\xi)^{-1} M(x)] - p. \quad (32)$$

For a linear utility  $u$  averaging over the design, the same construction gives  $\phi_u(x, \xi) = u_x - u^\top \xi$ .

**The General Equivalence Theorem.** Since  $\Phi$  is concave and the simplex is the convex hull of its vertices  $\delta_x$ , the measure  $\xi^*$  maximizes  $\Phi$  if and only if no vertex direction ascends:

$$\boxed{\xi^* \text{ is D-optimal} \iff d(x, \xi^*) \equiv \text{tr}[M(\xi^*)^{-1} M(x)] \leq p \quad \text{for all } x \in \mathcal{X},} \quad (33)$$

with equality at every support point of  $\xi^*$ . This is the theorem of Kiefer and Wolfowitz [3], and it is what this section exists to obtain: a quantity computed from a candidate answer that *proves* the answer optimal. The gap  $\max_x d(x, \xi) - p$  is a certificate, not a convergence report.

Two further facts come free. Equality puts the support exactly where  $d(x, \xi^*)$  is largest, so the optimal design runs at the settings of greatest standardized prediction variance. D-optimality and G-optimality coincide. And by Carathéodory’s theorem applied to the convex hull of the  $M(x)$ , an optimal measure supported on at most  $p(p + 1)/2 + 1$  points always exists: at most 11 here.

**Computing it.** The implementation uses the multiplicative algorithm of Silvey, Titterton and Torsney [9],

$$\xi_i \leftarrow \frac{\xi_i F_i}{\sum_j \xi_j F_j}, \quad F_i = \text{tr}[M(\xi)^{-1} M(x_i)] \quad \text{for } \Phi = \log \det, \quad (34)$$

which stays on the simplex by construction: weights remain non-negative and normalized without projection, and the criterion rises at each step. Note that  $\sum_i \xi_i F_i = p$  identically, so (34) moves mass toward designs whose  $d(x, \xi)$  exceeds  $p$ , which is precisely the set that (33) says must be empty at the optimum. The update and the certificate are the same quantity.

Vertex-direction steps with a line search were tried first and cannot converge here. The step required near the optimum falls below any fixed search grid, and the iteration stalls with a positive gap.

Convergence of (34) degrades as candidates crowd together. A design just off the support is nearly as good as one on it, and its weight decays in proportion. Certifying a parabola on  $[-1, 1]$  took 6400 iterations over 51 candidates, 25 000 over 101, and 329 000 over 401 – with the criterion value correct to six figures long before any of those.

An exhausted budget therefore yields a converged value without a proof. That is a different thing from a wrong answer, and the reported gap distinguishes them.

**The certified design for these records.** Over 480 candidates spanning  $R_{\max} \in [50, 1200] \mu\text{m}$  and stretch  $[3, 20]$ , with  $M(x) = J^\top J$  for the four qSLS parameters at the posterior median, the algorithm certifies at a gap of  $9 \times 10^{-10}$  in 1406 iterations:  $\max_x \text{tr}[M^{-1} M(x)] = 4.00000000$  against  $p = 4$ . The optimal measure is

| $R_{\max}$         | stretch | weight |
|--------------------|---------|--------|
| 1015 $\mu\text{m}$ | 6.70    | 0.470  |
| 161 $\mu\text{m}$  | 19.26   | 0.303  |
| 50 $\mu\text{m}$   | 17.78   | 0.227  |

against a best single design of 1015  $\mu\text{m}$  at stretch 6.70. The measure exceeds it by 6.99 in log det, a D-efficiency ratio of  $\exp(6.99/4) = 5.74$ ; equivalently the confidence ellipsoid is  $\exp(6.99/2) \approx 33$  times smaller in volume, and a typical parameter’s interval is  $\exp(6.99/8) = 2.4$  times tighter. No single design can match it, and this is not a claim about the search: (33) certifies that none exists.

**The mixture is the resolution of the conflict.** The support is worth reading physically. Nearly half the runs go to a large, gently driven bubble and the rest to small, hard-driven ones – the two corners the earlier subsections found pulling in opposite directions. Separating  $g$  from  $\alpha$  wants a gentle collapse, because the distinguishing term of  $Z_e$  decays as  $\lambda^{-4}$ ; exciting the relaxation mode wants a fast one.

A single design must choose between them. A measure need not. So the trade-off that motivated the Pareto front dissolves once the design space is the one the experiment actually admits – the front was answering a question made hard by an artificial restriction.

**Discrimination in the same frame.** Log evidence is additive over independent experiments, so an expected log Bayes factor per run is linear in  $\xi$  and therefore concave. The compound criterion

$$\Phi_\beta(\xi) = (1 - \beta) \log \det M(\xi) + \beta u^\top \xi, \quad \beta \in [0, 1], \quad (35)$$

is then concave for every  $\beta$  and inherits the certificate unchanged. Its directional derivative is the matching combination of (32) and  $u_x - u^\top \xi$ . This is the compound-criterion construction of Atkinson and of Cook and Wong [10, 11] – the principled form of the trade this study otherwise handled by search.

Sweeping  $\beta$  traces the Pareto front, and each criterion being concave in  $\xi$  makes that front convex. This is why the classical literature scalarizes with weighted sums where this study needed a Chebyshev form. The non-convexity that forced the elaborate scalarization was an artifact of optimizing over a single design point.




Figure 20: The equivalence-theorem certificate. Left: the sensitivity  $d(x, \xi^*) = \text{tr}[M(\xi^*)^{-1}M(x)]$  over the design plane, with the three support points circled and labeled by weight; the dashed contour is the level  $p = 4$ . Right: the same values sorted. No design exceeds the bound and the support attains it exactly, which is the proof  $-\max_x d = 4.00000000$  against  $p = 4$ , gap  $9 \times 10^{-10}$ . Nothing analogous exists for the single-design searches elsewhere in this document.

**What this does not fix.** Three limitations, stated plainly. The information matrices are evaluated at a point estimate of  $\theta$ , so the design is locally optimal in the parameters though globally optimal in  $\xi$ . A prior average over  $\theta$  restores the guarantee at proportionate cost, since the average of concave criteria is concave.

The candidate set is still a grid, so the certificate proves optimality *over that grid*. But  $d(x, \xi^*)$  can be evaluated anywhere, which makes checking for a missed region a cheap scan rather than another search.

The compound form was not run on these records. The per-design log Bayes factor still suffers the boundary pinning described above.

## 11 What the design survives

The certified measure of Section 10 is a theorem. Three things stand between it and an experiment somebody can run, and each is measurable rather than arguable.

### A measure assigns weights; an experiment runs integers

The optimum splits runs three ways. With twelve bubbles that is 5.4/3.7/2.9, and the equivalence theorem certifies optimality over *measures* — an exact design is not one, so the certificate does not reach it. `pyimr.measure.apportion` closes the gap by the efficient rounding of Pukelsheim and Rieder, and because that rule carries no guarantee at small budgets, it reports the D-efficiency it achieved rather than asserting one.

On the four-point design of cubic regression, where every quantity is closed form: budgets divisible by four lose nothing, and six runs give 2/2/1/1 at an efficiency of 0.9428, which is  $[(2/6)^2(1/6)^2/(1/4)^4]^{1/4}$  exactly. The rule earns its complexity on awkward budgets — at weights 0.7/0.2/0.1 and four runs, proportional rounding gives 3/1/0, dropping a support point and leaving the information matrix singular.

### Frames are a design variable, and there are far too many of them

Section 13.1 reports 201 samples carrying the information of about ten. Redundancy is a design variable: a camera trades frame rate against window length, against exposure, and against how many bubbles fit in a




Figure 21: Why a model difference can be invisible at every design. Changing a model perturbs the trace by  $\Delta = R_B - R_A$ . The component lying in the span of the material sensitivities is reproduced by adjusting  $g, \mu, \lambda_1, \alpha$  – refitting absorbs it, and no experiment recovers it. Only the orthogonal remainder is evidence that the model changed. Measured on the 15 °C record and reported in Section 7: the bubble-dynamics operator is 77.7% absorbed, leaving under two noise units, which is why its ranking proved so fragile; the constitutive and thermal axes leave 8.1 and 8.5.

session. Scoring frames by  $\log \det J^T J$  at the fitted qSLS, with the same number of observations spent each way:

| observations | uniform | optimally placed | gain |
|--------------|---------|------------------|------|
| 201          | 18.54   | 25.72            | 6.0× |
| 50           | 11.73   | 20.15            | 8.2× |
| 12           | 5.98    | 14.42            | 8.3× |

The D-optimal measure over the 201 candidate times is certified at a gap of  $10^{-10}$  on *six* support points. Fifty observations placed on those six carry more information than all 201 spread uniformly.

That is a measurement of slack, not a recommendation. A design on six distinct times leaves no degrees of freedom to notice the model is wrong, and the lack-of-fit test above is exactly what those degrees of freedom buy. The honest reading is that frames are cheap relative to their information, so the camera should be traded for something scarce — bubbles, or window, or the replication that makes (16) possible.

## The recommended geometries do not survive model error

Optimal designs are computed assuming the model is right. Ours is not: (16) says so at every temperature. The literature on sloppy systems warns that a model can fit worse on data from its own optimal experiment, and that is testable before a bubble is spent.

The test must not generate data from the model being fitted — that is the assumption under examination, and a study that makes it finds nothing. So the truth here carries the thermal transport the fitted model lacks, a mismatch Section 7 measures at 16 noise units, and cold qSLS is fitted to it at four geometries.

| geometry                                          | $\chi^2/N$   | lag-one | $g\alpha$ error | model error |
|---------------------------------------------------|--------------|---------|-----------------|-------------|
| performed, 277 $\mu\text{m}$ at 7.09              | 1.231        | 0.231   | <b>3.8%</b>     | 0.53        |
| E-optimal, 199 $\mu\text{m}$ at 3.55              | 2.533        | 0.504   | 5.9%            | 1.24        |
| discrimination-optimal, 100 $\mu\text{m}$ at 20.0 | 2.197        | 0.488   | <b>37.1%</b>    | 1.17        |
| measure support, 60 $\mu\text{m}$ at 18.0         | <b>0.892</b> | 0.009   | 5.2%            | <b>0.15</b> |

The warning is real and it is worse than advertised. The sloppy-systems literature describes a *trade* — better parameters, worse fit. Two of these three recommendations lose on both counts. The E-optimal geometry fits 2.1 times worse and recovers  $g\alpha$  1.6 times worse; the discrimination-optimal geometry fits 1.8 times worse and recovers  $g\alpha$  nearly *ten* times worse. The geometry already performed beats both.

The mechanism is visible in the last column. A criterion computed under the correct-model assumption sends the experiment where the model is most sensitive, and for these models that is also where it is most wrong: the thermal mismatch grows from 0.53 noise units at the performed geometry to 1.24 at the E-optimal one. The parameters then absorb the mismatch, which is why recovery degrades rather than improving.

One design escapes, and instructively. The small support point at 60  $\mu\text{m}$  carries the *least* model error of the four, 0.15 against 0.53, and is the only geometry that fits better than the one performed. Robustness to the error we have measured, not information under the assumption that we have none, is what distinguishes it.

This is one mismatch at one truth, and a different model error could rank differently. What does not depend on that choice is the direction: design criteria evaluated under a model the data reject can recommend experiments that are worse on every axis they were meant to improve. A design should be scored against the model error we have measured before it is offered to anyone.

## 12 Temperature dependence

The most informative result is not any single number but a trend (Figure 22). The advantage that the stiffening term buys over pure relaxation shrinks monotonically as the gel warms: the gap between SLS and qSLS falls from 2.94 versus 1.35 at 15  $^{\circ}\text{C}$  to 1.41 versus 1.23 at 33  $^{\circ}\text{C}$ . A warmer, softer gel needs less strain-stiffening to explain its record. Because this is a trend across three independent datasets rather than a single fitted value, it is harder to attribute to a fitting artifact than any individual  $\chi^2$ .




Figure 22: What the stiffening term buys, against temperature. The advantage of qSLS over SLS narrows as the gel warms.

## 13 Conclusions

We have treated the model comparison in inertial microcavitation rheometry as the object of study rather than a step toward a parameter value. Ranking by evidence in coordinates the records can answer, three findings hold.

The shear modulus is not identified separately from the stiffening exponent, so a reported modulus is a statement about the prior box. The bubble-dynamics operator assumed throughout this literature is beaten on every record. And a correlated residual that has been read as constitutive model error survives every constitutive, observational and operator change tested here.

Two of those findings needed qualifying once the equilibrium radius was fitted rather than asserted. Compressibility is required on every record at every prior width, by 74 to 179 nats. Gilmore still beats the pressure form at 15  $^{\circ}\text{C}$ . But at 23  $^{\circ}\text{C}$  and 33  $^{\circ}\text{C}$  the operator margins collapse to under a nat, so those records never discriminated operators at all – pinning  $R_{\text{eq}}$  is what made it appear they did. The fit wants  $R_{\text{eq}}$  3% to 11% larger than inferred, consistently across six operators and three records, and we cannot yet explain the trend with temperature.

### 13.1 Limitations of this work

**The dropped points are outside the model, not outside the solver.** Roughly one point in six of the winner’s parameter box cannot be integrated, and that is not a numerical shortcoming. In that region the model expands to  $R/R_0 = 2132$  after its collapse – identically at relative tolerances of  $10^{-3}$ ,  $10^{-4}$  and  $10^{-5}$ . So it is a converged property of the equations, not an artifact. The bubble begins at its maximum radius and its energy is finite, so growth of that size is the constitutive model failing. Refusing those points is the correct outcome.

The count of admissible points is pinned near 1958 of 2401 sampled whatever the settings. Looser tolerances and a  $20\times$  step budget convert failures into runaways, never into usable solves. This corrects an earlier reading of the same failures as a stiffness problem. The dropped points cannot be recovered by numerical effort, and no claim about the winner’s fit should be made in that region.




Figure 23: Where the winner can be integrated, at the fitted modulus and viscosity. Failure is governed principally by the stiffening, and the threshold falls to its lowest where the relaxation time is comparable to the bubble’s own period – but it is a boundary that moves, not an isolated band.

**The ranking survives; the parameter estimates do not.** Independent sequential Monte Carlo runs give the winner  $2173.55 \pm 0.19$  in log evidence against  $2071.90 \pm 0.38$  for the relaxation-only model – a gap of 101.6 against run-to-run noise near 0.4, so the comparison is resolved by a wide margin. The tensor grid cannot say this on its own: it returns a single number with no uncertainty.

The same grid cannot estimate parameters. Refining it from 12 to 20 points per axis moves the log evidence by 17.3, and moves the best-fit point from  $g = 658$ ,  $\alpha = 1.52$  to  $g = 144$ ,  $\alpha = 8.86$ . The best cell is simply whichever node lands nearest the ridge.

Refinement cannot fix this. The posterior standard deviations are 0.0018, 0.0025, 0.0055 and 0.078 in unit coordinates, against a grid spacing of 0.053. The posterior therefore occupies about  $10^{-7}$  of the prior volume, and a uniform grid would need of order  $10^{10}$  points. Discretization error of 17.3 sits well inside the 101.6 gap, so the ranking is safe; the reported best-fit parameters are not. Figure 24 shows which models lose points.

**Consequences for the numbers reported here.** Three methods that resolve the posterior – sampling, a refined grid, and continuous optimization – agree that  $g$  lies near  $10^2$  Pa. The coarse grid used for the ranking reports 2154 Pa. The ranking does not depend on this; any material property quoted from the grid does.

A modulus that size is also implausibly soft for gelatin. That is the degeneracy above making itself felt: the fit buys a lower residual by trading modulus against stiffening, along a direction the data barely constrain.




Figure 24: Fraction of each model’s parameter grid that could not be integrated. Models not shown lost no points.

**One candidate was never genuinely exercised.** The Gent solid appears in the comparison but cannot be tested on these records. Its energy diverges as  $I_1 - 3 \rightarrow J_m$ , and a Gent solid soft enough to be distinguishable from Neo-Hookean predicts a collapse it cannot sustain, so it locks up. Where it does integrate, it differs from Neo-Hookean by  $1.7 \times 10^{-4}$  of  $R_{\max}$  against measurement noise near  $2 \times 10^{-2}$  – two orders of magnitude below what the data can resolve. Any statement about strain-stiffening laws here excludes it.

## A Statistical background

This appendix develops, from first principles, the results the main text uses. It is self-contained and assumes only calculus and elementary probability.

### A.1 Inference as conditioning

Write  $\theta$  for parameters,  $y$  for data,  $M$  for a model. The product rule  $p(\theta, y) = p(y | \theta) p(\theta) = p(\theta | y) p(y)$  rearranges to Bayes’ theorem,

$$p(\theta | y, M) = \frac{p(y | \theta, M) \pi(\theta | M)}{p(y | M)}, \quad p(y | M) = \int p(y | \theta, M) \pi(\theta | M) d\theta. \quad (36)$$

The denominator is a normalizing constant in  $\theta$ , which is why it is ignored when estimating parameters and why it is the entire content of model comparison: applying (36) at the level of models gives

$$\frac{p(M_1 | y)}{p(M_2 | y)} = \underbrace{\frac{p(y | M_1)}{p(y | M_2)}}_{\text{Bayes factor } B_{12}} \cdot \frac{p(M_1)}{p(M_2)}. \quad (37)$$

Everything the data say about which model to prefer is in  $B_{12}$ . Jeffreys’ conventional reading takes  $\log B > 1$  as substantial and  $\log B > 5$  as decisive; the gap reported here is over 100, which is why its numerical resolution had to be checked rather than assumed.

### A.2 The score, and two definitions of information

Let  $\ell(\theta) = \log p(y | \theta)$ . The *score* is  $u = \nabla_{\theta} \ell$ . Since  $\int p(y | \theta) dy = 1$  for every  $\theta$ , differentiating under the integral gives

$$\mathbb{E}_{\theta}[u] = \int \frac{\nabla p}{p} p dy = \nabla \int p dy = 0, \quad (38)$$



so the score has zero mean at the true parameter. Its covariance defines the Fisher information,  $F = \mathbb{E}[uu^\top]$ . Differentiating the same identity twice gives the equivalent and more useful form

$$F = \mathbb{E}[-\nabla^2 \ell], \quad (39)$$

the expected curvature. The two agree because  $\nabla^2 \log p = \nabla^2 p/p - (\nabla p/p)(\nabla p/p)^\top$  and the first term integrates to zero. Curvature and variance-of-score are the same object, which is the reason a sharply peaked likelihood and a precisely determined parameter are the same statement.

### A.3 The Cramér–Rao bound

For an unbiased estimator  $\hat{\theta}(y)$ , differentiate  $\mathbb{E}[\hat{\theta}] = \theta$  under the integral to get  $\mathbb{E}[\hat{\theta} u^\top] = I$ . Applying the Cauchy–Schwarz inequality to the pair  $(\hat{\theta} - \theta, u)$  in the scalar case gives  $\text{Var}(\hat{\theta}) \mathbb{E}[u^2] \geq 1$ , i.e.  $\text{Var}(\hat{\theta}) \geq 1/F$ ; the multivariate statement is

$$\text{Cov}(\hat{\theta}) \succeq F^{-1}. \quad (40)$$

So  $F^{-1}$  is a floor on achievable precision, attained asymptotically by maximum likelihood:  $\sqrt{N}(\hat{\theta}_N - \theta) \rightarrow \mathcal{N}(0, F_1^{-1})$  under regularity. This is the license for reading  $F^{-1}$  as a covariance and its eigenvectors as the axes of a confidence ellipsoid – and the regularity conditions are exactly what near-singular  $F$  threatens, which is why sloppiness is worth detecting rather than tolerating.

### A.4 Gaussian data, and why $F = J^\top J$

With independent Gaussian observations of known deviation,  $\ell = -\frac{1}{2} \sum_i (y_i - m_i(\theta))^2 / \sigma_i^2 + \text{const}$ . Then

$$\frac{\partial \ell}{\partial \theta_j} = \sum_i \frac{y_i - m_i}{\sigma_i^2} \frac{\partial m_i}{\partial \theta_j}, \quad -\frac{\partial^2 \ell}{\partial \theta_j \partial \theta_k} = \sum_i \frac{1}{\sigma_i^2} \left[ \frac{\partial m_i}{\partial \theta_j} \frac{\partial m_i}{\partial \theta_k} - (y_i - m_i) \frac{\partial^2 m_i}{\partial \theta_j \partial \theta_k} \right]. \quad (41)$$

Taking the expectation kills the second bracket because  $\mathbb{E}[y_i - m_i] = 0$ , leaving  $F = J^\top J$  with  $J_{ij} = \sigma_i^{-1} \partial m_i / \partial \theta_j$ . Note what was discarded: the curvature of the model. At the optimum the residuals are small and the Gauss–Newton form is accurate; far from it, or with a badly misspecified model,  $F$  and the observed Hessian differ, and the discrepancy is itself a diagnostic.

### A.5 The Laplace approximation and the Occam factor

To evaluate  $Z = \int e^{\ell(\theta)} \pi(\theta) d\theta$ , expand about the maximum  $\hat{\theta}$  where  $\nabla \ell = 0$ :

$$\ell(\hat{\theta} + \delta) = \ell(\hat{\theta}) - \frac{1}{2} \delta^\top H \delta + O(\|\delta\|^3), \quad H = -\nabla^2 \ell(\hat{\theta}). \quad (42)$$

Holding  $\pi$  constant over the peak and using  $\int e^{-\delta^\top H \delta / 2} d\delta = (2\pi)^{p/2} |H|^{-1/2}$ ,

$$Z \simeq e^{\ell(\hat{\theta})} \pi(\hat{\theta}) (2\pi)^{p/2} |H|^{-1/2}. \quad (43)$$


The correction is  $O(N^{-1})$  relative for regular problems. Writing the posterior volume as  $V_{\text{post}} = (2\pi)^{p/2} |H|^{-1/2}$  and the prior density as  $1/V_{\text{prior}}$  where it is flat, the second factor is  $V_{\text{post}}/V_{\text{prior}}$ : the fraction of the space the data leave standing. A model with more parameters has a larger  $V_{\text{prior}}$  and pays for it unless the fit improves enough to compensate. Nothing was added by hand; the penalty is the Jacobian of the integral.

Taking logarithms of (43), using  $H \approx NF_1$  for  $N$  independent observations, and dropping terms that do not grow with  $N$ ,

$$-2 \log Z \simeq -2\ell(\hat{\theta}) + p \log N + O(1), \quad (44)$$


which is the Bayesian information criterion. BIC is therefore not an alternative to the evidence but a coarse asymptotic of it, valid when the posterior is Gaussian, the parameters identified, and  $N$  large. AIC answers a different question – expected out-of-sample Kullback–Leibler divergence – and is not an approximation to  $Z$  at all. Where a model is sloppy,  $H$  is near-singular,  $|H|^{-1/2}$  is enormous, and the asymptotic form fails while the integral remains well defined.

a direction the data determines



$V_{\text{post}}/V_{\text{prior}} < 1$   
the model *pays*

a direction it does not



$\lambda_k \rightarrow 0$ , so  $-\frac{1}{2} \log \det \rightarrow +\infty$   
the model is *paid*

Figure 25: The Occam factor is the share of the prior the data leaves standing,  $\prod_k \sqrt{2\pi/\lambda_k}$  over the eigenvalues of  $J^T J$ . Left: every direction is narrower than its prior, so a flexible model pays for volume it does not use. Right: a direction the data does not determine has  $\lambda_k \rightarrow 0$ , the fitted Gaussian extends far outside the box the prior occupies, and the expansion returns a factor *greater* than one – the evidence rises when a useless parameter is added. Capping each direction at unity, (46), restores what the geometry already required; on the models of Section 5.4 the uncapped form favored the two-mode law by 29.7 nats exactly where its second arm did nothing.

## A.6 Capping the Occam factor at the prior

The failure just noted is not merely a loss of accuracy; it has a sign, and the sign is wrong. Work in coordinates where the prior is uniform on the unit cube, so  $V_{\text{prior}} = 1$  and  $H = J^T J$ . The Occam factor is a product over the eigendirections of  $J^T J$ ,

$$\frac{V_{\text{post}}}{V_{\text{prior}}} = \prod_{i=1}^p \sqrt{\frac{2\pi}{\lambda_i}}, \quad (45)$$

each factor being the posterior width in that direction against a prior of width one. A direction the data does not determine has  $\lambda_i \rightarrow 0$  and contributes a factor diverging to  $+\infty$ . The evidence then *increases* with the addition of a parameter that does nothing, which is the opposite of what the Occam factor exists to do.

The pathology is entirely an artifact of the Gaussian approximation. The fitted Gaussian has width  $\sqrt{2\pi/\lambda_i} \gg 1$  in that direction and so extends far outside the cube the prior actually occupies, while the true integral, taken over the cube, simply returns the prior mass. Imposing what the geometry already requires – a posterior cannot be wider than the prior it came from – gives

$$\log \frac{V_{\text{post}}}{V_{\text{prior}}} = \sum_{i=1}^p \min\left(0, \frac{1}{2} \log \frac{2\pi}{\lambda_i}\right), \quad (46)$$

so a direction the data cannot see costs nothing and buys nothing. Where every  $\lambda_i > 2\pi$  each minimum selects its second argument and (46) reduces identically to the logarithm of (45): the cap is a strict generalization, altering an answer only where the uncapped form was not entitled to one. Section 5.4 gives the size of the error it removes, which on the models considered there is 29.7 nats.

This matters wherever models of differing dimension are compared and some parameters are weakly identified – which, for the sloppy spectra of Section A.7, is the normal case rather than the exceptional one.

## A.7 Sloppy models

Write the eigen-decomposition  $F = V\Lambda V^T$  with  $\lambda_1 \geq \dots \geq \lambda_p$ . The misfit near the optimum is  $\Delta\chi^2 = \sum_k \lambda_k c_k^2$  for  $\delta = \sum_k c_k v_k$ , so moving a distance  $c$  along  $v_k$  costs  $\lambda_k c^2$  and the 95% contour lies at  $c_k = \sqrt{3.84/\lambda_k}$ . Multiparameter models of this kind generically show spectra spanning many decades, with a few stiff directions and many sloppy ones – the observation of Gutenkunst [12], Sethna and co-workers, and the reason parameter values in such models are often far less determined than their predictions.

The distinction to keep is between a rank-deficient  $F$ , where a direction is *structurally* unidentifiable and no experiment helps, and a merely ill-conditioned  $F$ , where the direction is determined but weakly. In




Figure 26: Why a “best fit” for  $g$  is a statement about the prior. The likelihood depends on the product  $g\alpha$ , so the misfit surface is a valley running along constant  $g\alpha$  and the data is flat along  $g/\alpha$ . An optimizer slides until it meets the edge of the box, and *where* it stops is set by the box: the two dashed rectangles give the two filled points, which differ in  $g$  by more than thirtyfold. This is the mechanism behind the fifty-nat swing in the operator ranking of Section 6 when the bounds were widened, and behind the order-of-magnitude disagreement in  $g$  between methods in Figure 9.

logarithmic parameters, an eigenvector is a power-law combination, and the invariant it leaves fixed is read off its components.

## A.8 Correlated errors

Let  $r$  be the residual vector with covariance  $C$ . The Gaussian log-likelihood is

$$-2\ell = r^T C^{-1} r + \log |C| + N \log 2\pi, \quad (47)$$

and with the Cholesky factorization  $C = LL^T$  this becomes  $\|L^{-1}r\|^2 + 2\sum_k \log L_{kk} + \text{const}$ : whitening turns generalized least squares back into ordinary least squares on transformed residuals. Ignoring the off-diagonal terms, i.e. using  $\text{diag}(C)$ , does not merely lose efficiency – it misstates the amount of information. For a stationary sequence with autocorrelation  $\rho_k$ ,

$$\text{Var}\left(\sum_{i=1}^N r_i\right) = N\sigma^2 \left(1 + 2\sum_{k=1}^{N-1} \left(1 - \frac{k}{N}\right)\rho_k\right) \xrightarrow{N \rightarrow \infty} N\sigma^2 \left(1 + 2\sum_{k \geq 1} \rho_k\right), \quad (48)$$

so the sequence carries the information of  $N_{\text{eff}} = N/(1 + 2\sum \rho_k)$  independent draws. The bracket is the spectral density of the residual sequence at zero frequency, which is the cleaner way to see why a long-range correlation is so costly.

Correlated residuals in a *fitted* model usually indicate model discrepancy rather than correlated measurement noise. Kennedy and O’Hagan [13] formalize this by writing  $y = m(\theta) + \delta(\cdot) + \varepsilon$  with  $\delta$  a Gaussian process, and Brynjarsdóttir and O’Hagan show that omitting  $\delta$  leaves parameter estimates biased *and* overconfident, while an unconstrained  $\delta$  absorbs the physics. Neither failure is visible in  $\chi^2$ .

## A.9 Optimal experimental design

A design  $\xi$  determines what is measured and hence  $F(\xi, \theta)$ . Since  $F$  is a matrix and designs must be ranked, a scalar functional is required, and the classical choices are

$$\Phi_D = \log \det F, \quad \Phi_A = -\text{tr } F^{-1}, \quad \Phi_E = \lambda_{\min}(F), \quad \Phi_c = -c^T F^{-1} c, \quad (49)$$




Figure 27: Two failures that look alike in a fit and differ completely in what can be done about them. Left: the misfit is exactly flat along a direction,  $F$  is singular, and the parameter combination is structurally unidentifiable. No experiment recovers it, and only a reparameterization or a fixed value helps. Right: the misfit is closed but very elongated,  $F$  is invertible and ill-conditioned, and the combination is determined weakly. Here a design that rotates the ellipse can close it, which is what Section 10 sets out to do. On these records  $\lambda_{\min} = 5.2 \times 10^{-2} \neq 0$ , so the problem is the second kind.




Figure 28: Why some parameter combinations are determined and others are not. Near the best fit the misfit rises as a quadratic form, so the contour of constant misfit is an ellipse whose axes point along the eigenvectors of  $F = J^T J$  and whose half-lengths go as  $1/\sqrt{\lambda}$ . A large eigenvalue is a short axis: move that way and the fit degrades at once. A small one is a long axis: move that way and the data barely object. On these records the ratio is  $1.1 \times 10^3$ , and the long direction is  $g$  up with  $\alpha$  down at fixed product, which is the ridge of Figure 26 seen from above. The ellipse is drawn at an aspect of about 8 : 1; the measured aspect is  $\sqrt{1.1 \times 10^3} \approx 33 : 1$ , which would render as a line.

each maximized. In Bayesian terms,  $\Phi_D$  is the expected Kullback–Leibler gain from prior to posterior under a Gaussian approximation, which is why it is the default: it maximizes information about *all* parameters jointly. That default is wrong whenever one direction is the question, because  $\log \det F = \sum_k \log \lambda_k$  is dominated by the large eigenvalues.

Because  $F$  depends on  $\theta$ , a local design presupposes an answer. The standard repairs are to average over a prior,  $\Phi(\xi) = \mathbb{E}_\theta[\Phi(F(\xi, \theta))]$ , or to take the worst case over a plausible set. Sequential design uses the posterior from completed experiments as the prior for the next, which is what is done here.

When the model itself is uncertain, all of the above presuppose the wrong thing. T-optimality (Atkinson and Fedorov) maximizes instead the lack of fit of the inferior model,

$$\Phi_T(\xi) = \min_{\theta_2} \|m_1(\hat{\theta}_1; \xi) - m_2(\theta_2; \xi)\|^2, \quad (50)$$

which asks which experiment the candidates most disagree about. Its virtue here is that it shares no assumptions with the Fisher criteria, so agreement between them is evidence rather than restatement.

## B Reproducing

```
.venv/bin/python examples/measured_selection.py <dataset> [thermal Nt] [workers]
```




Figure 29: Why 201 samples do not carry 201 samples’ worth of evidence. The residuals here are strongly correlated from one sample to the next, so neighboring points repeat information rather than adding it. A correlated sequence carries what  $N_{\text{eff}} = N/(1 + 2 \sum_k \rho_k)$  independent draws would, and on these fits  $N_{\text{eff}}$  is near 8. Both  $\chi^2/N$  and the evidence assume the samples are independent, so each overstates the information by roughly  $N/N_{\text{eff}} \approx 25$ . A margin of a few nats between two models is inside that factor.

```
.venv/bin/python docs/writeup/collect.py 16      # records results.json
.venv/bin/python docs/writeup/figures.py         # draws every figure from it
```

Every number and figure in this document comes from `results.json`; none is transcribed by hand.

## Code and data availability

Every number and figure in this document is reproduced by scripts in the repository that contains it: `figures.py`, `figures_geometry.py`, `figures_identifiability.py`, `figures_measure.py` and `figures_trial.py` draw the figures, `fit_quality.py` the fit table and `trial_variation.py` the trial-spread analysis, each from the tracked `results.json`. The design measures use `pyimr.measure`, the discrimination criteria `pyimr.discriminate`, and the trade-off front `pyimr.pareto`. The measured records themselves are not redistributed here.

## References

- [1] Jonathan B. Estrada, Carlos Barajas, David L. Henann, Eric Johnsen, and Christian Franck. High strain-rate soft material characterization via inertial cavitation. *Journal of the Mechanics and Physics of Solids*, 112:291–317, 2018.
- [2] Jonathan B. Freund and Randy H. Ewoldt. Quantitative rheological model selection: Good fits versus credible models using Bayesian inference. *Journal of Rheology*, 59(3):667–701, 2015.
- [3] J. Kiefer and J. Wolfowitz. The equivalence of two extremum problems. *Canadian Journal of Mathematics*, 12:363–366, 1960.
- [4] Joseph B. Keller and Michael Miksis. Bubble oscillations of large amplitude. *The Journal of the Acoustical Society of America*, 68(2):628–633, 1980.
- [5] Charles J. Geyer. Practical Markov chain Monte Carlo. *Statistical Science*, 7(4):473–483, 1992.
- [6] A. C. Atkinson and V. V. Fedorov. The design of experiments for discriminating between two rival models. *Biometrika*, 62(1):57–70, 1975.
- [7] D. V. Lindley. A statistical paradox. *Biometrika*, 44(1–2):187–192, 1957.
- [8] Harold Jeffreys. Some tests of significance, treated by the theory of probability. *Mathematical Proceedings of the Cambridge Philosophical Society*, 31(2):203–222, 1935.
- [9] S. D. Silvey, D. M. Titterton, and B. Torsney. An algorithm for optimal designs on a finite design space. *Communications in Statistics — Theory and Methods*, A7:1379–1389, 1978.
- [10] A. C. Atkinson. DT-optimum designs for model discrimination and parameter estimation. *Journal of Statistical Planning and Inference*, 138(1):56–64, 2008.

- [11] R. D. Cook and W. K. Wong. On the equivalence of constrained and compound optimal designs. *Journal of the American Statistical Association*, 89(426):687–692, 1994.
- [12] Ryan N. Gutenkunst, Joshua J. Waterfall, Fergal P. Casey, Kevin S. Brown, Christopher R. Myers, and James P. Sethna. Universally sloppy parameter sensitivities in systems biology models. *PLoS Computational Biology*, 3(10):e189, 2007.
- [13] Marc C. Kennedy and Anthony O’Hagan. Bayesian calibration of computer models. *Journal of the Royal Statistical Society: Series B*, 63(3):425–464, 2001.