

goal: sufficient conditions for local optimality in NLP subject to constraints

refs:

OPTIMAL CONTROL AND ESTIMATION

Robert F. Stengel

Department of Mechanical and Aerospace Engineering
Princeton University, Princeton, New Jersey

Section 2.1

Nonlinear Programming

SECOND EDITION

Dimitri P. Bertsekas

Massachusetts Institute of Technology

Chapter 3

o consider NLP with constraints: $\min_{w \in \mathbb{R}^l} c(w)$ $c: \mathbb{R}^l \rightarrow \mathbb{R}$
s.t. $f(w) = 0$ $f: \mathbb{R}^l \rightarrow \mathbb{R}^n$

→ assuming you can explicitly solve the constraint equation
show how you can reduce to an equivalent unconstrained NLP

- suppose $w \in \mathbb{R}^l$ can be "split" into two pieces, $w = (x, u) \in \mathbb{R}^n \times \mathbb{R}^m$
and explicitly solve $f(x, u) = 0$ for x in terms of u ,

i.e. $x: \mathbb{R}^m \rightarrow \mathbb{R}^n \rightsquigarrow$ then $\min_{u \in \mathbb{R}^m} c(x(u), u)$ is equivalent
to original problem
: $u \mapsto x(u)$

i.e. $X: \mathbb{R}^n \rightarrow \mathbb{R}^m$ \mapsto then $\min_{u \in \mathbb{R}^m} C(X(u), u)$ is equivalent to original problem
 $: u \mapsto X(u)$

• can't solve $f(u)=0$ in general, so we'll consider another approach

idea: consider augmented cost function

$$\tilde{C}(x, u, \lambda) = C(x, u) + \lambda \cdot f(x, u)$$

$\lambda \in \mathbb{R}^{1 \times n}$ termed "Lagrange multipliers" or "dual variables"

and find stationary points of \tilde{C}

i.e. want $\underbrace{(x_0, u_0, \lambda_0)}_{\in \mathbb{R}^{(n+m+n)}} \text{ s.t. } \underbrace{D\tilde{C}(x_0, u_0, \lambda_0)}_{\in \mathbb{R}^{1 \times (n+m+n)}} = 0$

→ determine necessary condition for λ_0 to be stationary
 (assume $D_x f(x_0, u_0) \in \mathbb{R}^{n \times n}$ invertible / non singular)

- $D_x \tilde{C} = D_x C + \lambda \cdot D_x f$ so if $D_x f(x_0, u_0)$ invertible

then it's necessary that $\lambda_0 = -D_x C(x_0, u_0) \cdot [D_x f(x_0, u_0)]^{-1}$

def: $(x_0, u_0) \in \mathbb{R}^{n+m}$ is a stationary point for $\min_{(x,u) \in \mathbb{R}^{n+m}} C(x, u)$
 s.t. $f(x, u) = 0 \in \mathbb{R}^n$

if $D_u C(x_0, u_0) + \lambda_0 \cdot D_u f(x_0, u_0) = 0$ and $f(x_0, u_0) = 0$

where $\lambda_0 = -D_x C(x_0, u_0) \cdot [D_x f(x_0, u_0)]^{-1}$

→ why is it reasonable to assume $D_x f(x_0, u_0)$ invertible?

- otherwise constraints are redundant!

- otherwise constraints are redundant!

consider the linear case: $f(x,u) = L \cdot \begin{bmatrix} x \\ u \end{bmatrix}$

→ why are stationary points of constrained NLP the same as those of \tilde{c} ?

* full answer requires some work — cf wikipedia article "Lagrange multipliers"

→ when $n=1$ the geometry is easier to understand — cf pg 103 of
Folland's Advanced Calculus

ex: (Lewis Sympos 1.2-3) quadratic cost, linear constraint:

$$\min c(x,u) = \frac{1}{2} x^T Q x + \frac{1}{2} u^T R u \quad \text{— assume } Q^T = Q > 0$$

$$\text{s.t. } f(x,u) = x + B u + b = 0$$

$$R^T = R > 0$$

→ write augmented cost & stationarity conditions

$$- \tilde{c}(x,u,\lambda) = c(x,u) + \lambda \cdot f(x,u)$$

$$= \frac{1}{2} x^T Q x + \frac{1}{2} u^T R u + \lambda \cdot (x + B u + b)$$

$$- D_x \tilde{c} = D_x c + \lambda \cdot D_x f = x^T Q + \lambda = 0$$

$$- D_u \tilde{c} = D_u c + \lambda \cdot D_u f = u^T R + \lambda \cdot B = 0$$

$$- D_x \tilde{c} = f(x,u) = x + B u + b = 0$$

→ solve for stationary u (& λ)

$$- \text{from } D_x \tilde{c} = 0 \text{ we find } \lambda = -x^T Q \Leftrightarrow \lambda^T = -Q x$$

$$n \quad n \quad n \times 1 \quad n \times n \quad n \times 1 \quad n \times 1 \quad n \times 1 \quad n \times 1 \quad n \times 1 \quad n \times 1$$

- from $D_x C = 0$ we find $\lambda = -R^{-1}B^T Q x$
- from $D_u \tilde{C} = 0$ we find $u = -R^{-1}B^T \lambda$
- combining these we find $u^* = -R^{-1}B^T Q x$