

# The place of the local group in the cosmic web

Sebastian Bustamante <sup>★1</sup> Jaime E. Forero-Romero<sup>2</sup>

<sup>1</sup>*Instituto de Física - FCEN, Universidad de Antioquia, Calle 67 No. 53-108, Medellín, Colombia*

<sup>2</sup>*Departamento de Física, Universidad de los Andes, Cra. 1 No. 18A-10, Edificio Ip, Bogotá, Colombia*

17 September 2013

## ABSTRACT

We present a theoretical study Local Group in a cosmological context. The principal objective is to find a coherent sample of LG systems in the unconstrained simulation Bolshoi based upon the properties of hand construted LG systems in the constrained simulations CLUES. We obtain a privileged environment for these systems and significant bias in some of their properties. (**draft!**)

**Key words:** galaxies: high-redshift - galaxies: star formation - line: formation

## 1 INTRODUCTION

The spatial distribution of galaxies describes a web-like pattern, the so-called cosmic web. Today it is understood that such configuration is driven by gravitational instabilities. ...

The study of the influence of the cosmic web on galaxy properties start with the seminal work of Dressler [[reference here](#)] and extends to recent works using large observational surveys that look for signatures of the web into the evolution of galaxy populations. With the advent of more detailed observations and sophisticated computational models it is now within our reach to understand what physical processes dominate.

This makes that the mass assembly history of a galaxy is deeply connected with its position in the cosmic web. There is an extensive body of literature on the effects of the web environment on the observable properties of galaxies.

This environmental study is also of paramount importance to understand the formation of our Galaxy. In our local neighborhood, the observations of dwarf galaxies around the Milky Way (MW) and the Andromeda galaxy (M31) show filamentary and disk-like patterns that can be linked to a preferential infall direction, very likely connected with the cosmic web where the Local Group (LG) of galaxies is embedded.

In this paper we quantify the velocity shear environment of DM halo pairs representative of the principal members of the Local Group (LG), the Milky Way (MW) and Andromeda galaxy (M31). We perform this study in an unconstrained cosmological simulation from random phases in the initial conditions, and unlike previous works, where were used constrained cosmological simulations which have been setup as to reproduce the large scale structure of the local

universe, we use directly observational measurements of the kinematics properties of the local group [[Reference here](#)] in order to build faithful samples of LG-like systems.

We pay special attention to the correlation of the present velocity shear environment with the assembly and the kinematics properties of the pairs. The motivation to have that focus is that it has been previously shown that the LG present in three different realizations of the constrained simulations have assembly histories biased towards early formation times and absence of major mergers (ratio 1:10) in the last 10 Gyr. In the case of the kinematic properties, recent observational constrains to the galactocentric tangential velocity of M31 has enabled to establish how typical is the LG in a cosmological context [[reference to Forero-Romero 2013-1](#)], that is why we focus here how a specific kind of host environment biases these kinematics properties.

## 2 THE SIMULATION

As it was previously mentioned, we use an unconstrained cosmological simulation, the Bolshoi simulation, to identify the possible large scale environment of the Local Group. This is a similar approach to the one already used by [[reference here](#)].

The Bolshoi simulations follows the non-linear evolution of a dark matter density field on a cubic volume of size  $250h^{-1}\text{Mpc}$  sampled with  $2048^3$  particles. The cosmological parameters in the simulation are  $\Omega_m = 0.27$ ,  $\Omega_\Lambda = 0.73$ ,  $h = 0.70$ ,  $n = 0.95$  and  $\sigma_8 = 0.82$  for the matter density, cosmological constant, dimensionless Hubble parameter, spectral index of primordial density perturbations and normalization for the power spectrum. The mass of each particle in the simulation is  $m_p = 1.4 \times 10^8 h^{-1} M_\odot$ .

★ sbustama@pegasus.udea.edu.co

### 2.1 Halos and Merger Trees

We identify halos with two algorithms, the Friends-of-Friends [reference here] algorithm and the Bound Density Maximum algorithm. The constructed catalogues also provide the basis for the mass aggregation history studies. We also include in the catalogues information about the substructure.

All the results presented here must be interpreted in term of host halos, without any information of the substructure. In particular the merger of two FOF halos corresponds to the epoch of first overlap, and not to the fusion and/or disruption of an accreted sub-halo with a dominant halo.

The linking length is  $b = 0.17$  times the mean inter-particle separation. All objects with 20 particles or more are considered a bona fide halo and are included in the construction of the merger tree, this corresponds to a minimum halo mass of  $M_{\min} = 3.78 \times 10^8 h^{-1} M_{\odot}$  in the CLUES volumes and  $M_{\min} = 2.70 \times 10^9 h^{-1} M_{\odot}$  in the Bolshoi simulation.

The halo identification for the CLUES simulation was done for 80 snapshots in the redshift range  $0 < z < 7$  more or less evenly spaced in look-back time. For Bolshoi we use the information of XX snapshots in the same redshift range.

## 3 ALGORITHMS TO QUANTIFY THE COSMIC ENVIRONMENT

### 3.1 The tidal web - T-web

The first algorithm we use to identify the cosmic web is based on the diagonalization of the tidal tensor, defined as the Hessian of a normalized gravitational potential

$$T_{\alpha\beta} = \frac{\partial^2 \phi}{\partial x_{\alpha} \partial x_{\beta}} \quad (1)$$

where the physical gravitational potential has been rescaled by a factor  $4\pi G \bar{\rho}$  in such a way that  $\phi$  satisfies the following equation

$$\nabla^2 \phi = \delta, \quad (2)$$

where  $\bar{\rho}$  is the average density in the Universe,  $G$  is the gravitational constant and  $\delta$  is the dimensionless matter overdensity.

### 3.2 The velocity web - V-web

We use the a kinematical method to define the cosmic-web environment in the simulation. The method has been thoroughly described in XXX and applied to study the shape and spin alignment in the Bolshoi simulation here XX. We refer the reader to these papers to find a detailed description of the algorithm, its limitations and capabilities. Here we summarize the most relevant points for the discussion.

The V-web method for environment finding is based on the local shear tensor calculated from the smoothed DM velocity field in the simulation. The central quantity is the following dimensionless quantity

$$\Sigma_{\alpha\beta} = -\frac{1}{2H_0} \left( \frac{\partial v_{\alpha}}{\partial x_{\beta}} + \frac{\partial v_{\beta}}{\partial x_{\alpha}} \right) \quad (3)$$

where  $v_{\alpha}$  and  $x_{\alpha}$  represent the  $\alpha$  component of the comoving velocity and position, respectively.  $\Sigma_{\alpha\beta}$  can be represented by a  $3 \times 3$  symmetric matrix with real values, that ensures that is possible to diagonalize and obtain three real eigenvalues  $\lambda_1 > \lambda_2 > \lambda_3$  whose sum (the trace of  $\Sigma_{\alpha\beta}$ ) is proportional to the divergence of the local velocity field smoothed on the physical scale  $\mathcal{R}$ .

The relative strength of the three eigenvalues with respect to a threshold values  $\lambda_{th}$  allows for the local classification of the matter distribution into four web types. Voids, sheets, filaments and peaks, which correspond to regions with 3, 2, 1 or 0 eigenvalues with values larger than  $\lambda_{th}$ . In this paper we do not perform an definitive classification into web types, instead we express the results respect to the relative strength of the eigenvalues and in a range of  $\lambda_{th}$ . We consider *a posteriori* the possible web type interpretation that could be feasible within a range of thresholds for the eigenvalues.

### 3.3 The Web in Bolshoi

... Resolution and Smoothing  
... Database

### 3.4 Method to find void regions

Following the recent work of Courtois et al. 2013, we use a method based on a FOF algorithm to find extended regions of voids and then, we select halo systems according to a proximity criteria to those regions. In order to perform this, we build the input FOF catalogue with the positions of the center coordinate of every cell marked as void according to the web scheme adopted; furthermore we set an adequate linking length for connecting diagonal neighbour cells.

Following the work of Forero-Romero et al. 2008, we also perform a percolation analysis in order to select the best threshold parameter that reduce percolation in cells, thereby accounting for physical voids regions. In the next figure we show the obtained result of our percolation analysis for both web schemes. In both cases it can be noted that percolation (the largest void region volume) is completely reduced for  $\lambda_{th} = 0.0$ , so despite of the previously established  $\lambda_{th}$  values for each scheme, just for the detection of void regions we shall use  $\lambda_{th} = 0.0$

## 4 LOCAL GROUP SAMPLE DEFINITION

In order to establish an adequate set of criteria to define LG samples in unconstrained simulations, we proceed from the the general dark halo catalogues, constructed using the FOF scheme with a linking length of  $b = 0.17$ . These samples are defined for all simulation and will be referred as General Halos (GH) samples. To be consistent with the observationally determined mass range of halos that host disk-form galaxies, we select the halos in the mass range  $5 \times 10^{11} < M_h / h^{-1} M_{\odot} < 5 \times 10^{12}$ , referred here as Individual Halos (IH) samples.

As a primal approach to define gravitational bounded halo pairs we select all halo pairs in IH samples that satisfied to be the closest to each other, they constitute the Pairs (P) samples. To keep the concordance with previous

works in algorithms for LG selection ([[references here](#)]), we define here the next list of conditions that have to fulfill a pair system in order to construct the Isolated Pairs (IP) samples. All these considerations are based upon the relative dynamics of the Milky Way and M31, and its isolation from massive structures:

- (i) The distance between the center of the two halos should be less than  $0.7h^{-1}\text{Mpc}$ .
- (ii) The relative physical velocity between the two halos has to be negative.
- (iii) The distance to any halo more massive than any of the pair members must be less than  $2h^{-1}\text{Mpc}$ .
- (iv) The distance to cluster-like halos with masses larger than  $1 \times 10^{13} h^{-1}M_{\odot}$  must be larger than  $5h^{-1}\text{Mpc}$ .

In the case of a constrained simulation one can define samples of LG systems based upon the next condition

- (v) The Local Group pair must be located in the right environment with respect to the XX Cluster.

Due to the intrinsic nature of constrained simulations, we expect that these LG samples be the most faithful systems that resemble the properties and the environment of our LG. In fact, the three CLUES simulations appear to present a common V-web/T-web environment for LG systems, such as shown in the figure ???. Based in this we propose a new criteria of selection to define more realistic LG systems in the unconstrained simulation, this consists in tanking the extreme values of each V-web/T-web eigenvalue associated to the host environment of the three LG systems in CLUES. With this range established (see table 1) we filter the IP sample in Bolshoi simulation in order to build the samples Constructed Local Groups V-web based (CLGV) and T-web based (CLGT). Finally in the table ?? we show the sizes of all defined samples for each simulation.

## 5 FINDING A LOCAL GROUP ENVIRONMENT

In this section we prepare some numerical experiments with the halos samples and their environment, with special emphasis on the isolated pairs and LG samples. All this in order to find for environment correlations and common properties between LG systems.

### 5.1 Comparison of the two simulations

Prior to study of isolated and LG samples and to determine possible correlations between their properties, it is necessary to establish the equivalence between all simulations that we will use, with the aim to eliminate effects due to construction process of each one.

In first place, we analyse the mass distribution of individual halos and in next figure we show the integrated mass function (IMF) for halos sample with  $M \geq 1 \times 10^{11} M_{\odot}$ .

Although in high masses the IMF of each simulation are slightly different, in low mass region, where the most halos of our interest are, the IMFs are quite similar indicating that the sample of halos in each simulation have the same mass distribution, while the small differences are due to finite size of samples. Another aspect in the figure ?? is the position of

LG halos, they are distributed across the mass range of halos that we have set (see in section 4), indicating that there is not an apparent condition with their individual masses.

The situation is quite different respect to mass ratio index of isolated pairs sample (MRI). In the figure ?? (a) we show the integrated distribution of MRI, where the approximately linear behaviour indicates an uniformly distribution. The interesting aspect here is the closeness between the LG sample values, indicating a possible common property in these systems, even though this could be established a priori by construction. In the ?? (b) is plotted the integrated distribution of total pair mass, and again like the IMF, there is not a preferential value.

Once established the concordance between the defined samples, we proceed to analyse the distribution of the eigenvalue of shear velocity tensor 3, for this we assume that the halos are good tracers of the environment properties and therefore we evaluate each eigenvalue in the center mass of each halo, mapping of this way the complete distribution (figure ??). The black curves correspond to the unconstrained simulation (Bolshoi) and the color curves to the constrained simulations (CLUES), additionally we calculate the cosmic variance, showed in purple curves, dividing the Bolshoi volume in smaller parts with a size comparable to the CLUES volume ( $64h^{-1}\text{Mpc}$  of side). What is interesting here is the clear difference between the eigenvalues distributions of the both types of simulations, inclusively the constrained distributions does not match between the cosmic variance, indicating that although both simulations were made with the same cosmology (see subsections ?? and ??), the constrained simulations have a significantly different environment properties (spatial matter distribution) compared to the average expected from a random volume with the same comoving size.

### 5.2 Defining the LG environment

With the aim of constructing a sample of LG systems in Bolshoi simulations, we use the eigenvalues of the three LG systems in each CLUES simulation and choose a range for each one based in the extreme values of the six halos, this with the expectation of reproducing the LG specific conditions in constrained simulations. This criteria can be thought as a first approximation to establish a more faithful sub-sample into the isolated pairs. In table 1 we show the used extreme values for each eigenvalue of local shear tensor.

As a self-consistency test, we apply this criteria to CLUES simulations and find, on average, three LG systems in each one. But to avoid confusion, we keep defining the LG sample as the three initial pairs. Finally, we construct the LG sample of Bolshoi, where the sample size is illustrated in Table ??.

Once we have established the equivalence of samples in each simulation and have defined the LG samples, we proceed to calculate correlations between halo samples and their environment. As was mentioned in section 3.2, we do not use an specific value of eigenvalue threshold  $\lambda_{th}$ , instead of this, we explore a relatively wide range of this parameter (i.e.  $0 \leq \lambda_{th} \leq 1$ ) and calculate distributions respect to each eigenvalue individually.

At first place, we calculate the environment of each LG system in CLUES simulations, for this we use the  $\lambda_{th}$  scheme

to classify it in void, sheet, filament or knot (see 3.2), with  $\lambda_{th}$  into the threshold range. Figure ?? is obtained.

At first place, we calculate the integrated distribution of each eigenvalue for individual halos, isolated pairs and LG samples. For this, we associate to each halo a value of environment (set of eigenvalues) according to its center of mass position in a smoothed grid ( $256^3$  cells for Bolshoi and  $64^3$  for CLUES, or equivalently a resolution of  $1.0\text{Mpc h}^{-1}$  per cell, according to the physical size of the pairs.)

## 6 RESULTS

### 6.1 Bias induced on kinematic and dynamics properties

- ... Total Mass
  - ... Radial vs. tangential velocities
  - ... Angular Momentum
  - ... Total Mechanical energy.
  - ... Reduced Spin

### 6.2 Bias induced on the Mass Assembly Histories


- ... Last major merger. Formation Time. Assembly Time.

### 6.3 Pair Alignment with the Cosmic Web

- ... Separation.
  - ... Relative velocity.
  - ... Angular Momentum.

## 7 CONCLUSIONS


## ACKNOWLEDGMENTS



**Figure 1.** Integrated mass function of individual halos of each simulation used. In the center-left, with vertical dot lines, the individual masses of each LG system in CLUES simulations are represented.

	$\lambda_1 [10^{-1}]$	$\lambda_2 [10^{-1}]$	$\lambda_3 [10^{-1}]$
Minim value	1.78	$-6.29 \times 10^{-2}$	-1.98
Maxim value	3.49	1.21	$-8.85 \times 10^{-1}$

**Table 1.** Extreme values for each V-web eigenvalues to construct LG samples.



**Figure 2.** Integrated mass function of individual halos of Bolshoi.