


# Research Proposal for a Master Thesis in Physics

## Verifying the VPH scheme in Galaxy Formation

---

Sebastian Bustamante Jaramillo



Time evolution of a gas cloud in a supersonic wind using a VPH scheme. Taken from (Heß & Springel, 2010)

## Contents

<b>1</b>	<b>General Information</b>	<b>2</b>
<b>2</b>	<b>Abstract</b>	<b>2</b>
<b>3</b>	<b>Introduction</b>	<b>3</b>
<b>4</b>	<b>Objectives</b>	<b>4</b>
<b>5</b>	<b>Theoretical Framework</b>	<b>4</b>
<b>6</b>	<b>Methodology</b>	<b>4</b>
<b>7</b>	<b>Expected Results</b>	<b>5</b>
<b>8</b>	<b>Scientific Impact</b>	<b>6</b>
<b>9</b>	<b>Schedule</b>	<b>6</b>
<b>8</b>	<b>Bibliography</b>	<b>7</b>

# 1 General Information

## Information of the Student

<b>Name</b>	Sebastian Bustamante Jaramillo
<b>Degree</b>	B.Sc. in Physics, Universidad de Antioquia (2013)
<b>E-mail 1</b>	macsebas33 <i>at</i> gmail.com (personal)
<b>E-mail 2</b>	sebastian.bustamante <i>at</i> udea.edu.co (academic)

More detailed information of the applicant can be found here <http://goo.gl/BPZGzK>

## Information of the Project

<b>Title</b>	<b>Verifying the VPH scheme in Galaxy Formation</b>
<b>Field</b>	Cosmology, Astrophysics, Physical Sciences
<b>Advisor 1</b>	Professor Juan Carlos Munoz-Cuartas. Universidad de Antioquia, Colombia.
<b>University</b>	Universidad de Antioquia, Master of Physics program
<b>Time Frame</b>	2 years

# 2 Abstract

### 3 Introduction

As we understand more deeply the physical processes involved in astrophysical phenomena, it becomes necessary to compute complex interactions of a ever increasing number of single components. Some prominent examples include the large-scale Universe, galaxy evolution, stellar interior, star formation and protoplanetary disk dynamics. A common aspect of these examples is that all of them can be regarded basically as a fluid mechanic problem.

Although the development of analytical approaches has demonstrated to be a valuable resource for studying these processes, their increasing complexity makes necessary to invoke numerical solutions as a more viable alternative. For this purpose, two different families of hydrodynamics solvers has been explored and widely used by the astrophysical community. First, a family of moving-mesh-based techniques (e.g. *Smoothed Particle Hydrodynamics* SPH (Monaghan, 1992), *Voronoi Particle Hydrodynamics* VPH (Heß & Springel, 2010)), and a second family of fixed-mesh-based techniques (e.g. *Adaptive Mesh Refinement* AMR (Berger & Colella, 1989)).

Due to the Lagrangian character of moving-mesh methods, techniques like SPH are easily implemented on a computer. Furthermore, as the physical system evolves, the mass particles naturally move into higher density regions, providing a self-adjusting spatial resolution. Nevertheless, SPH has been shown to produce spurious suppression of fluid instabilities due to its kernel-based density estimator, making it unsuitable to model some of the dynamics accurately. On the other hand, fixed-mesh methods like AMR are more efficient for capturing shock dynamics. However, due to the conservative nature of the hydrodynamical equations, a fixed mesh causes a lack of Galilean invariance. Furthermore, the sampling of physical properties over the grid introduces spurious vorticity to the fluid, making this technique poor suitable for studying turbulent flows.

A completely new approach to solve hydrodynamical problems was introduced by Springel (2010) and implemented into the AREPO code. It combines the strengths of AMR and SPH but overcomes many of their weaknesses, hence it can be thought as a mixed technique. AREPO uses a moving mesh based on a Voronoi tessellation defined over a set of particles that represents the fluid. The geometry of the mesh resembles very closely that of the point distribution, retaining the self-adaptivity inherent of SPH and also keeping a grid to capture shocks like AMR does. These features make AREPO highly accurate for simulating a wide range of hydrodynamical problems. Nevertheless, there is a price to pay for this accuracy, AREPO demands a huge computing time as compared with SPH and even AMR.

A very interesting alternative was introduced by Heß & Springel (2010), i.e. the *Voronoi Particle Hydrodynamics* VPH technique. This approach consists of an implementation of SPH with a modified density estimator based on the *Voronoi Tessellation Field Estimator* VTFE. The new estimator has demonstrated to improve substantially the spurious suppression of fluid instabilities as well as retaining the computational efficiency of the original formulation.

Finally, galaxy evolution and large-scale structure formation are very rich astrophysical phenomena where a plethora of hydrodynamical processes can be found and studied. In this fashion, cosmological simulations are a quite suitable scenario for performing detailed physical and computational comparisons of all above-mentioned techniques. It is especially interesting to quantify the computational performance of the VPH technique in terms of its physical accuracy as compared with the classic approaches and AREPO.

## 4 Objectives

### General Objective

Quantifying the computational performance of the VPH technique in terms of its physical accuracy for a cosmological setup.

### Specific Objectives

- Evaluating the physical accuracy provided by VPH for a cosmological setup as compared with AMR, SPH and AREPO.
- Exploring and quantifying the differences between VPH and AMR for describing shock dynamics in specific hydrodynamical instabilities.
- Exploring and quantifying the differences of VPH and SPH for describing turbulent flows.
- Measuring the computational performance of VPH as compared with AREPO.

## 5 Theoretical Framework

## 6 Methodology

The proposed project is subject to a M.Sc. study and will cover the following steps:

- ✓ *First, a bibliographic review of the original papers where were formulated each of the discussed methods will be done. Also a review of previous comparison projects.*

Before carrying out our enterprise in quantifying the performance of VPH over cosmological setups, it is necessary to understand deeply the foundations of the classic approaches. At this point, a detailed bibliographic review of the original papers (for SPH, AMR, VPH and AREPO) should be done. Although no previous works have been done in comparing thoroughly the performance of VPH with other approaches over cosmological setups, there are a plenty of

comparison projects for the classic approaches and even AREPO over galaxy simulations and commonly used benchmark problems. This literature will have to be reviewed as well.

- ✓ *Second, a design of the numerical experiments should be done at this point. This includes making cosmological simulations using different techniques and if necessary, constructing and simulating specific benchmark problems.*

As this project will be entirely based on numerical results, computing a set of cosmological simulations as well as some benchmark problems is one of the key steps. For this purpose, we will use some packages like GADGET (Springel, 2005) for SPH simulations, RAMSES Teyssier (2002) for AMR and a modified version of GADGET for VPH. Other standard benchmark problems will be also simulated, e.g. the sod shock tube, Kelvin-Helmholtz instabilities, a gas cloud in a supersonic wind.

- ✓ *Third, a thorough analysis of the numerical results will be done.*

Once obtained the numerical results from the performed simulations, a thorough analysis of the physical accuracy of VPH as compared with the other techniques will be done for each situation. A computational performance analysis of the VPH technique will be also carried out, i.e. computing time, memory and processor usage.

- ✓ *Fourth, a first-author paper with the main result will be prepared.*

The more relevant results of our project will be prepared as a paper and submitted to some high impact international journal. If possible, a participation in some international event is also included in this step.

- ✓ *Fifth, a thesis will be written.*

A dissertation for obtaining a M.Sc. in Physics degree will be prepared. A streamlined description of each technique will be included as well as the presentation of the performed simulations and a discussion of all our results and conclusions.

## 7 Expected Results

At the end of the stipulated development time for this project, we hope to have obtained the following results:

- A toolbox of codes to study the performance of hydro-solvers over cosmological setups and over standard benchmark problems in fluid mechanics.

- A set of cosmological simulations computed by using each of the studied techniques.
- A M.Sc. thesis.
- Submitting a first-author paper to an international journal.
- Participating with a poster or an oral presentation in an international event.

## 8 Scientific Impact

## 9 Schedule

Goals	Semester I	Semester II	Semester III	Semester IV
Bibliographic review	X			
Numerical experiments	X	X		
Analysis of results		X	X	
International journal paper			X	
Dissertation				X

## 8 Bibliography

Berger M. J., Colella P., 1989, *Journal of Computational Physics*, 82, 64

Heß S., Springel V., 2010, *MNRAS*, 406, 2289

Monaghan J. J., 1992, *ARA&A*, 30, 543

Springel V., 2005, *MNRAS*, 364, 1105

Springel V., 2010, *MNRAS*, 401, 791

Teyssier R., 2002, *A&A*, 385, 337