


Verifying the Voronoi-Particle Hydrodynamics (VPH) scheme in Galaxy Formation





Sebastian Bustamante
Universidad de Antioquia
Medellín, Colombia

Motivation


- Understanding and simulating hydrodynamics processes with good accuracy in astrophysics is a very important task as many phenomena are modeled as fluid systems.
- Provided the complexity of the hydrodynamic equations, numerical solutions are usually the only feasible approach.

Motivation



Motivation

The large-scale Universe is a very rich scenario where a plethora of hydrodynamical processes occur. In this fashion, cosmological setups are excellent systems for comparing hydrodynamic solvers.




The cosmological setup

The Universe at very large scales can be modeled as a homogeneous and isotropic fluid, following the next equation

$$H^2(t) = H_0^2 \left[(1 - \Omega_0) \frac{1}{a^2} + \Omega_m \frac{1}{a^3} + \Omega_r \frac{1}{a^4} + \Omega_\Lambda \right]$$

where $a(t)$ is the scale factor, $H = a/\dot{a}$, H_0 is the current Hubble constant.

The cosmological setup



The cosmological setup

However, a homogeneous and isotropic Universe can not account for complex structures like galaxies, stars, planets and us!

Continuity equation

$$\frac{\partial \delta}{\partial t} = -\frac{1}{a} \nabla_r \cdot [(1 + \delta) \mathbf{v}]$$

Euler's equation

$$\frac{\partial \mathbf{v}}{\partial t} + \frac{\dot{a}}{a} \mathbf{v} + \frac{1}{a} (\mathbf{v} \cdot \nabla_r) \mathbf{v} = -\frac{\nabla_r P}{a \bar{\rho}(1 + \delta)} - \frac{1}{a} \nabla_r \Phi$$

Poisson's equation

$$\nabla_r^2 \Phi = 4\pi G \bar{\rho} a^2 \delta$$

$$\rho = \bar{\rho} + \delta \rho = \bar{\rho}(1 + \delta)$$

The cosmological setup

When modes of the density field are small enough ($\delta \ll 1$), it is possible to decouple them, obtaining the next equation:

$$\frac{d^2\delta_{\mathbf{k}}}{dt^2} + 2\frac{\dot{a}}{a}\frac{d\delta_{\mathbf{k}}}{dt} = \left[4\pi G\bar{\rho} - \frac{c_s^2}{a^2}k^2\right]\delta_{\mathbf{k}}$$

Obtaining the analytical solution for the lineal regime:

$$\delta_{\mathbf{k}}(t) = \delta_{\mathbf{k}}(0)D(t)$$


$$D(a) = \frac{5}{2}\Omega_m \left[\Omega_m^{4/7} - \Omega_\Lambda + \left(1 + \frac{\Omega_m}{2}\right) \left(1 + \frac{\Omega_\Lambda}{70}\right) \right]^{-1}$$

The cosmological setup

Even for the quasi-linear regime ($\delta \sim 1$), it is possible to obtain analytical solutions through the Zel'dovich approximation:


$$\mathbf{r}(t, \mathbf{q}) = a(t) [\mathbf{q} + \Psi(\mathbf{q}, t)]$$

$$\Psi(\mathbf{q}) = \int \frac{d^3 k}{(2\pi)^3} e^{i\mathbf{k}\cdot\mathbf{q}} \frac{i\mathbf{k}}{k^2} \delta_{\mathbf{k}}(0)$$



The cosmological setup

For the non-linear regime, the Fourier modes of the density are not longer coupled. Numerical evolution is necessary henceforth. However, quasi-linear solutions provide initial conditions for numerical schemes.



Hydrodynamic solvers

Once clarified the importance of the hydrodynamics in Galaxy formation (for evolving the non-linear regime), two families of numerical approaches has been adopted throughout the literature:

- **Lagrangian schemes:** the fluid is evolved following the dynamics of fluid parcels.
- **Eulerian schemes:** the evolution performed over a fixed coordinate system, namely a mesh.


Lagrangian schemes

- The most representative Lagrangian scheme is **SPH** (Smoothed-Particle Hydrodynamic).
- Other schemes, normally modifications of the original **SPH**, have been also implemented.
- Among the most interesting proposals is the **VPH** (Voronoi-Particle Hydrodynamic) scheme. This scheme greatly improves the accuracy of **SPH**.

Smoothed-Particle Hydrodynamic

- Although a fluid is a continuous medium, numerical solutions always requires some kind of discretization.
- SPH reaches this by sampling the fluid through particles that represent a parcel.
- The equations of motions are then integrated over each of moving particles.
- Continuous properties are then recovered through interpolation techniques.

Smoothed-Particle Hydrodynamic



Smoothed-Particle Hydrodynamic

The heart of **SPH** is the kernel interpolant formalism:

$$F_s(\mathbf{r}) = \int_V \mathbf{F}(\mathbf{r}') W(\mathbf{r} - \mathbf{r}'; h) d\mathbf{r}'$$

Discretization can be reached approximating
 $\Delta\mathbf{r}_i = m_i/\rho_i$, yielding:

$$F'_i \equiv F_s(\mathbf{r}_i) \approx \sum_j^{N_{tot}} \frac{m_j}{\rho_j} F_j W(\mathbf{r}_i - \mathbf{r}_j; h_i)$$

Where h_j is the smoothing length and is associated to the volume of the fluid parcel.

Smoothed-Particle Hydrodynamic

The equations of motions may be derived from the Lagrangian

$$\mathcal{L} = \int \rho \left(\frac{v^2}{2} - u \right) dV$$


Using the equation of state of inviscid gas and discretizing, we obtain

$$\frac{d\mathbf{v}_i}{dt} = - \sum_{j=1}^N m_j \left[f_i \frac{P_i}{\rho_i^2} \nabla_i W_{ij}(h_i) + f_j \frac{P_j}{\rho_j^2} \nabla_i W_{ij}(h_j) \right]$$

Smoothed-Particle Hydrodynamic

Computational performance is greatly improved if we use a finite-range kernel:

$$W\left(\frac{r}{2h}\right) = W(q) = \frac{8}{\pi} \begin{cases} 1 - 6q^2 + 6q^3, & 0 \leq q \leq 1/2, \\ 2(1 - q)^3, & 1/2 < q \leq 1, \\ 0, & q > 1 \end{cases}$$



Wabeeja
Medawigie
Meri
unichesh
Tingki
Komapsuminda
Shukuria
Paldies
Hater
hui
Sanco
bolzin
Maake
Denkaaja
Agoy
Spasibo
gozaimashita
Fakause
Spasibo
Mehrbani
Nurachallya
Ekhmet
Arigato
Dankscheen

Tashakkur
Maketal
hui
Sanco
bolzin
Maake
Denkaaja
Agoy
Spasibo
gozaimashita
Fakause
Spasibo
Mehrbani
Nurachallya
Ekhmet
Arigato
Dankscheen

Balika Yaspagaraktam
Minmonchar Atto
Garjho
Sakoms
Malteika
ekojia
Tariqach
lah
Merastawly
Dhanybaad
Chilla

Merci
suksama
ekojia
Tariqach
lah
Merastawly
Dhanybaad
Chilla

Shukria
Gracias
Tank
Biiyan
Grazie
Juspaxar