

Neural Network solutions to Witsenhausen problem

Jiaojiao Fan

GTID:903565753

Email: jiaojiaofan@gatech.edu

Abstract—In this report, several neural networks with different structures are implemented to solve Witsenhausen problem. Other improving strategies include optimizers, initializations and forced function fixing. Finally, the result are compared with former people and a better result is obtained. Also, the shortcoming of the neural network also shows in this project. The neural network may be stuck into a near local minima.

1. Introduction

In this report, we proposed several solutions to the well-known and still unsolved Witsenhausen counterexample. [1] There have been some meaningful tries to detect the global minima of the min problem, such as Lee [2] and M. Baglietto [3]. Some of their manipulations are also referred in this project. Other than that, thanks to the development of the neural networks, many other meaningful attempts are also taken such as input convex neural network (ICNN) structure [4]. Different results would be listed to show the effect.

2. The Witsenhausen Counterexample

The Witsenhausen counterexample has been outstanding for more than 50 years. It is formulated by Hans Witsenhausen in 1968. [1] It is a counterexample to a natural conjecture that in a system with linear dynamics, Gaussian disturbance, and quadratic cost, affine control laws are optimal to minimize the cost. However, Witsenhausen counterexample, shown in figure below, has nonlinear control laws




Figure 1. Witsenhausen counterexample

that outperform all linear laws.

$$f(x) = \gamma_1(x) + x \quad g(x) = \gamma_2(x) \quad (1)$$

As a result, $f(x_0) = x_1$ and $g(x_1 + z) = u_2$. Our goal is to minimize the quadratic cost: $k^2 \mathbb{E}[U_1^2] + \mathbb{E}[X_2^2]$, which can also be written as

$$\min J(f, g) := k^2 \mathbb{E}[f(X_0) - X_0]^2 + \mathbb{E}[(f(X_0) - g(f(X_0) + N))^2] \quad (2)$$

In equation (2), there is a parameter k^2 , which in fact determines the cost gap between the linear controller and nonlinear controller [3]. If k^2 is smaller, the gap is bigger. For better comparison with the results got by previous researchers, k^2 is set as 0.04 in this report.

In addition, it is already known that $f(x)$ must have some strict property to be optimal [5] :

- Any optimal controller f is a strictly increasing unbounded piecewise real analytic function with a real analytic inverse

Acknowledgments

Thanks to my advisor: Dr. Yongxin Chen and the other PhD students: Rahul Singh and Qinseng Zhang for their kind help. They give me a lot of ideas about how to improve the performance during the process of project.

References

- [1] H. S. Witsenhausen, "A counterexample in stochastic optimum control," *SIAM Journal on Control*, vol. 6, no. 1, pp. 131–147, 1968.
- [2] J. T. Lee, E. Lau, and Y.-C. Ho, "The witsenhausen counterexample: A hierarchical search approach for nonconvex optimization problems," *IEEE Transactions on Automatic Control*, vol. 46, no. 3, pp. 382–397, 2001.
- [3] M. Baglietto, T. Parisini, and R. Zoppoli, "Numerical solutions to the witsenhausen counterexample by approximating networks," *IEEE Transactions on Automatic Control*, vol. 46, no. 9, pp. 1471–1477, 2001.
- [4] B. Amos, L. Xu, and J. Z. Kolter, "Input convex neural networks," in *Proceedings of the 34th International Conference on Machine Learning-Volume 70*. JMLR. org, 2017, pp. 146–155.
- [5] Y. Wu and S. Verdú, "Witsenhausen's counterexample: A view from optimal transport theory," in *2011 50th IEEE Conference on Decision and Control and European Control Conference*. IEEE, 2011, pp. 5732–5737.