
Change with RET: CRE vs PLA (mixed-model ANOVA)
Model-estimated POST − PRE ± 95% CI; points = observed change (paired participants)
Change (POST − PRE)
Weight (kg)
Group (main) p = 0.128
Pooled change p = 0.471
CRE vs PLA p = 0.634
BMI
Group (main) p = 0.159
Pooled change p = 0.597
CRE vs PLA p = 0.619
Lean Body Mass (kg)
Group (main) p = 0.080
Pooled change p = 0.047
CRE vs PLA p = 0.414
Appendicular Lean Mass (kg)
Group (main) p = 0.096
Pooled change p < 0.001
CRE vs PLA p = 0.050
ALM Index
Group (main) p = 0.107
Pooled change p = 0.001
CRE vs PLA p = 0.097
Fat Mass (kg)
Group (main) p = 0.533
Pooled change p = 0.028
CRE vs PLA p = 0.941
Fat (%)
Group (main) p = 0.473
Pooled change p = 0.015
CRE vs PLA p = 0.913
Sit-to-Stand Max Power
Group (main) p = 0.076
Pooled change p = 0.543
CRE vs PLA p = 0.574
Grip Strength (lbs)
Group (main) p = 0.078
Pooled change p = 0.601
CRE vs PLA p = 0.968
Chest Press (lbs)
Group (main) p = 0.067
Pooled change p < 0.001
CRE vs PLA p = 0.299
Leg Extension (lbs)
Group (main) p = 0.153
Pooled change p < 0.001
CRE vs PLA p = 0.762
PLA
CRE