

This holds for $p = 7$ and $g = 3$. The remaining two cases $-g^2 \pm g + 1 = 0$ are equivalent because $-g^2 + g + 1 = 0$ if and only if $-g^{-2} - g^{-1} + 1 = 0$. In other words, we only need to consider the equality $g^2 = g + 1$ over \mathbb{F}_p . Such a primitive root g is called a *Fibonacci primitive root* modulo p , as the *golden ratio* ϕ_{gr} satisfying $\phi_{gr}^2 = \phi_{gr} + 1$. The primes p such that \mathbb{F}_p has a Fibonacci primitive root is the sequence [A003147](#): 5, 11, 19, 31, 41, 59, 61, 71, 79, 109, ... on [OEIS](#) [\[Slo\]](#). Consequently, we have $N(p, (p-1)/2) > 0$ if and only if \mathbb{F}_p has a Fibonacci primitive root. On the other hand, the order $|H|$ of the subgroup $H = \langle -1, 2 \rangle$ is an even integer. It follows that $\ell_0 = \frac{p-1}{|H|}$ must divide $\frac{p-1}{2}$. By Proposition [5.3](#), we see that Equation [\(2\)](#) is solvable over \mathbb{F}_p if \mathbb{F}_p has a Fibonacci primitive root. In this case, there exists a solution $(x, y) \in \mathbb{F}_p^2$ such that $xy \neq 0$ and thus Conjecture [C](#) holds.

Proposition 5.4. *If \mathbb{F}_p has a Fibonacci primitive root, then Conjecture [C](#) holds for this p .*

On a related issue, for the valid cases in Conjecture [C](#) established above we would like to know how many prime numbers p are there such that the subgroup H generated by -1 and 2 has the given index ℓ_0 in \mathbb{F}_p^\times . This question can be viewed as a generalization of the Artin's primitive root conjecture. It is shown in [\[Mur91, Theorem 1\]](#) that there are infinitely many primes p such that the index $[\mathbb{F}_p^\times : H] = \ell_0$ under the *Generalized Riemann Hypothesis* (GRH). Assuming GRH, we conclude from Corollary [B](#) that there are infinitely many prime numbers p with ℓ_0 satisfying conditions in Theorem [5.2](#) and therefore, the size of optimal CACs of prime lengths is equal to $M(p, \ell_0)$ for *infinitely many* primes p .

ACKNOWLEDGMENT

We would like to thank Professor Yuan-Hsun Lo for bringing [\[FLS14\]](#) to our attention. The first named author is partially supported by MOST grant 110-2115-M-003-007-MY2. The second named author is partially supported by MOST grant 111-2115-M-003-005. The third named author is supported by MOST grant 110-2811-M-003-530.

REFERENCES

- [AA53] D.R. Anderson and T.M. Apostol. The evaluation of Ramanujan's sum and generalizations. *Duke Math. J.*, 20(2):211–216, 1953. [doi:10.1215/S0012-7094-53-02021-3](#)
- [Bur11] D.M. Burton. *Elementary Number Theory*. McGraw-Hill, New York, 2011.
- [FLM10] H.-L. Fu, Y.-H. Lin, and M. Mishima. Optimal conflict-avoiding codes of even length and weight 3. *IEEE Trans. Inform. Theory*, 56(11):5747–5756, 2010. [doi:10.1109/TIT.2010.2069270](#)
- [FLS14] H.-L. Fu, Y.-H. Lo, and S.W. Shum. Optimal conflict-avoiding codes of odd length and weight three. *Des. Codes Cryptogr.*, 72(2):289–309, 2014. [doi:10.1007/s10623-012-9764-5](#)
- [GV93] L. Györfi and I. Vajda. Constructions of protocol sequences for multiple access collision channel without feedback. *IEEE Trans. Inform. Theory*, 39(5):1762–1765, 1993. [doi:10.1109/18.259673](#)
- [Har77] R. Hartshore. *Algebraic Geometry*, volume 52 of *Graduate Texts in Mathematics*. Springer, New York, 1977. [doi:10.1007/978-1-4757-3849-0](#)