

# Intermixing-driven surface and bulk ferromagnetism in the quantum anomalous Hall candidate $\text{MnBi}_6\text{Te}_{10}$

Abdul V. Tcakaev,<sup>1,2</sup> Bastian Rubrecht,<sup>3</sup> Jorge I. Facio,<sup>3,4</sup> Volodymyr B. Zabolotnyy,<sup>1,2</sup> Laura T. Corredor,<sup>3</sup> Laura C. Folkers,<sup>5,2</sup> Ekaterina Kochetkova,<sup>3</sup> Thiago R. F. Peixoto,<sup>6,2</sup> Philipp Kagerer,<sup>6,2</sup> Simon Heinze,<sup>7</sup> Hendrik Bentmann,<sup>6,2</sup> Robert J. Green,<sup>8,9</sup> Pierluigi Gargiani,<sup>10</sup> Manuel Valvidares,<sup>10</sup> Eugen Weschke,<sup>11</sup> Maurits W. Haverkort,<sup>7</sup> Friedrich Reinert,<sup>6,2</sup> Jeroen van den Brink,<sup>3,12,2</sup> Bernd Büchner,<sup>3,5,2</sup> Anja U. B. Wolter,<sup>3,2</sup> Anna Isaeva,<sup>13,3,\*</sup> and Vladimir Hinkov<sup>1,2,†</sup>

<sup>1</sup>*Physikalisches Institut (EP-IV), Fakultät für Physik und Astronomie, Universität Würzburg, Am Hubland, D-97074 Würzburg, Germany*

<sup>2</sup>*Würzburg-Dresden Cluster of Excellence ct.qmat, Germany*

<sup>3</sup>*Leibniz Institut für Festkörper- und Werkstoffforschung (IFW) Dresden, Helmholtzstraße 20, D-01069 Dresden, Germany*

<sup>4</sup>*Centro Atómico Bariloche, Instituto de Nanociencia y Nanotecnología (CNEA-CONICET) and Instituto Balseiro. Av. Bustillo 9500, Bariloche (8400), Argentina.*

<sup>5</sup>*Institut für Festkörper- und Materialphysik, Technische Universität Dresden, 01062 Dresden, Germany*

<sup>6</sup>*Physikalisches Institut (EP-VII), Fakultät für Physik und Astronomie, Universität Würzburg, Am Hubland, D-97074 Würzburg, Germany*

<sup>7</sup>*Institute for Theoretical Physics, Heidelberg University, Philosophenweg 19, 69120 Heidelberg, Germany*

<sup>8</sup>*Department of Physics and Astronomy and Stewart Blusson Quantum Matter Institute, University of British Columbia, Vancouver, BC V6T 1Z4, Canada*

<sup>9</sup>*Department of Physics and Engineering Physics,*

*University of Saskatchewan, SK S7N 5E2 Saskatoon, Canada*

<sup>10</sup>*ALBA Synchrotron Light Source, E-08290 Cerdanyola del Vallès, Barcelona, Spain*

<sup>11</sup>*Helmholtz-Zentrum Berlin für Materialien und Energie, Albert-Einstein-Straße 15, D-12489 Berlin, Germany*

<sup>12</sup>*Institut für Theoretische Physik, Technische Universität Dresden, D-01062 Dresden, Germany*

<sup>13</sup>*Van der Waals-Zeeman Institute, Department of Physics and Astronomy,*

*University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands*

(Dated: February 3, 2023)

The recent realizations of the quantum anomalous Hall effect (QAHE) in  $\text{MnBi}_2\text{Te}_4$  and  $\text{MnBi}_4\text{Te}_7$  benchmark the  $(\text{MnBi}_2\text{Te}_4)(\text{Bi}_2\text{Te}_3)_n$  family as a promising hotbed for further QAHE improvements. The family owes its potential to its ferromagnetically (FM) ordered  $\text{MnBi}_2\text{Te}_4$  septuple layers (SL). However, the QAHE realization is complicated in  $\text{MnBi}_2\text{Te}_4$  and  $\text{MnBi}_4\text{Te}_7$  due to the substantial antiferromagnetic (AFM) coupling between the SL. An FM state, advantageous for the QAHE, can be stabilized by interlacing the SL with an increasing number  $n$  of  $\text{Bi}_2\text{Te}_3$  layers. However, the mechanisms driving the FM state and the number of necessary QLs are not understood, and the surface magnetism remains obscure. Here, we demonstrate robust FM properties in  $\text{MnBi}_6\text{Te}_{10}$  ( $n = 2$ ) with  $T_c \approx 12$  K and establish their origin in the Mn/Bi intermixing phenomenon by a combined experimental and theoretical study. Our measurements reveal a magnetically intact surface with a large magnetic moment, and with FM properties similar to the bulk. Our investigation thus consolidates the  $\text{MnBi}_6\text{Te}_{10}$  system as perspective for the QAHE at elevated temperatures.

## I. INTRODUCTION

Theory provides a seemingly straightforward avenue towards novel quantum effects such as the quantum anomalous Hall (QAH) effect [1–5] and axion electrodynamics [6–9], namely to induce a long-range ferromagnetic (FM) order in topological insulators (TI) [10, 11]. The vision of observing Majorana fermions and implementing topological qubits at superconductor/QAH insulator interfaces [12], ultra low-power electronics [13] and applications in spintronics [14] has ignited substantial experimental efforts in this direction. Yet, hitherto

the QAH effect (QAHE) has only been demonstrated in the (sub-) kelvin range [3, 4, 15]. The experimental realization of the QAHE is complicated by several simultaneous requirements to a candidate system: The Dirac point (DP) of the parent TI should be well within its bulk band gap; the chemical potential has to be tuned to the DP; the introduced magnetic subsystem should lead to a substantial surface ferromagnetism to open a large exchange gap at the DP; and the material's bulk should remain insulating.

The first materials to exhibit the QAHE were extrinsically doped  $(\text{V/Cr})_x(\text{Bi,Sb})_{2-x}\text{Te}_3$ , which consist of van-der-Waals coupled quintuple layers (QL, see Fig. 1). However, band engineering by tuning the Bi/Sb ratio does not move the DP sufficiently above the valence band [16], V/Cr impurity bands overlap with the alleged exchange gap [17] and residual bulk conductance destroys

\* Corresponding address: [a.isaeva@uva.nl](mailto:a.isaeva@uva.nl)

† Corresponding address: [hinkov@physik.uni-wuerzburg.de](mailto:hinkov@physik.uni-wuerzburg.de)