

The dual isogeny $\hat{\phi}$ is such that $\phi \circ \hat{\phi} = \times 3$, where $\times 3$ is the multiplication-by-3 map. The map defined by the dual isogeny $\hat{\phi}$ is given in the proof of Lemma 4.5

Consider now the exact sequence ([15], Section X.4, Remark 4.7)

$$(5) \quad 0 \rightarrow \hat{E}_D(\mathbb{Q})[\hat{\phi}]/\phi(E_{D'}(\mathbb{Q})[3]) \rightarrow \hat{E}_D(\mathbb{Q})/\phi(E_{D'}(\mathbb{Q})) \xrightarrow{\hat{\phi}} \\ \xrightarrow{\hat{\phi}} E_{D'}(\mathbb{Q})/3E_{D'}(\mathbb{Q}) \rightarrow E_{D'}(\mathbb{Q})/\hat{\phi}(\hat{E}_D(\mathbb{Q})) \rightarrow 0.$$

Since both $\ker_{\phi}(\overline{\mathbb{Q}}) = \mathcal{T}_{D'}$ and $\ker_{\hat{\phi}}(\overline{\mathbb{Q}}) = \hat{\mathcal{T}}_D$ contain no non-trivial rational point of order 3, the first quotient group of (5) vanishes and the rank $r(E_{D'})$ of $E_{D'}(\mathbb{Q})$ equals

$$(6) \quad r(E_{D'}) = \dim_{\mathbb{F}_3}(E_{D'}(\mathbb{Q})/3E_{D'}(\mathbb{Q})).$$

Consider the short exact sequence

$$(7) \quad 0 \rightarrow \mathcal{T}_{D'} \rightarrow E_{D'}(\overline{\mathbb{Q}}) \xrightarrow{\phi} \hat{E}_D(\overline{\mathbb{Q}}) \rightarrow 0.$$

From this, we obtain the long exact cohomology sequence which gives in particular the following

$$(8) \quad 0 \rightarrow \hat{E}_D(\mathbb{Q})/\phi(E_{D'}(\mathbb{Q})) \xrightarrow{\delta} H^1(G_{\mathbb{Q}}, \mathcal{T}_{D'}) \rightarrow H^1(G_{\mathbb{Q}}, E_{D'}(\overline{\mathbb{Q}}))[\phi] \rightarrow 0.$$

By localising at each prime p , we obtain the following commutative diagram, where res_p is the usual restriction map:

$$\begin{array}{ccccccc} 0 & \rightarrow & \hat{E}_D(\mathbb{Q})/\phi(E_{D'}(\mathbb{Q})) & \rightarrow & H^1(G_{\mathbb{Q}}, \mathcal{T}_{D'}) & \rightarrow & H^1(G_{\mathbb{Q}}, E_{D'}(\overline{\mathbb{Q}}))[\phi] \rightarrow 0 \\ & & \downarrow & & \downarrow res_p & & \downarrow res_p \\ 0 & \rightarrow & \hat{E}_D(\mathbb{Q}_p)/\phi(E_{D'}(\mathbb{Q}_p)) & \rightarrow & H^1(G_{\mathbb{Q}_p}, \mathcal{T}_{D'}) & \rightarrow & H^1(G_{\mathbb{Q}_p}, E_{D'}(\overline{\mathbb{Q}}_p))[\phi] \rightarrow 0 \end{array}$$

Definition 2.1. The Selmer group of $E_{D'}$ relative to the isogeny ϕ is

$$\mathcal{S}_{\phi}(E_{D'}) = \{x \in H^1(G_{\mathbb{Q}}, \mathcal{T}_{D'}) \mid res_p(x) \in \text{Im}(\hat{E}_D(\mathbb{Q}_p)/\phi(E_{D'}(\mathbb{Q}_p))) \text{ for all } p\}.$$

The Tate-Shafarevich group of $E_{D'}$ can now be defined as

$$\text{III}(E_{D'}) = \{x \in H^1(G_{\mathbb{Q}}, E_{D'}(\overline{\mathbb{Q}})) \mid res_p(x) = 0 \text{ for all } p\}.$$

These two groups are connected together as follows:

$$(9) \quad 0 \rightarrow \hat{E}_D(\mathbb{Q})/\phi(E_{D'}(\mathbb{Q})) \rightarrow \mathcal{S}_{\phi}(E_{D'}) \rightarrow \text{III}(E_{D'})[\phi] \rightarrow 0.$$

Remark 2.2. By considering the dual isogeny $\hat{\phi}$ instead, we get exact sequences analogous to (7), (8) and (9) which in turn give us the analogous definitions for $\mathcal{S}_{\hat{\phi}}(\hat{E}_D)$, $\text{III}(\hat{E}_D)$ and $\text{III}(E_D)[\hat{\phi}]$.

Remark 2.3. We also obtain exact sequences analogous to (7), (8) and (9) for $\times 3 = \phi \circ \hat{\phi}$ which in turn give us the analogous definitions for $\mathcal{S}_3(E_{D'})$, $\text{III}(E_{D'})$ and $\text{III}(E_{D'})[3]$, and similarly for \hat{E}_D .