

Supplementary materials for “scAClc: A Multi-Objective Adaptive Clustering Framework for Single-Cell Transcriptomics via Contrastive and Resolution-Aware Representation Learning”

Supplementary Tables

Table S1. The statistics of 15 real scRNA-seq datasets.

No.	dataset	organ	platform	cell types	cells	gene number
1	10X_PBMC[1]	Human PBMC	10x Genomic	8	4271	16653
2	Human_p[2]	Human pancreas	inDrop	14	3605	20125
3	Mammary_Gland[3]	Mammary Gland	10x Genomic	7	4481	23341
4	Klein[4]	Mouse embryo stem cells	inDrop	4	2717	24047
5	Adam[5]	Kidney	Drop-seq	8	3660	23797
6	Chen [6]	Mouse brain	Drop-seq	46	12089	23284
7	Turtle_b[7]	Turtle brain	Drop-seq	15	18664	23500
8	Baron_human[8]	Human pancreas	inDrop	14	8569	20125
9	Muraro[9]	Pancreas	Smart-seq2	9	2122	19046
10	Plasschaert[10]	Trachea	inDrop	8	6977	28205
11	Pollen[11]	Human tissues	SMARTer	11	301	21721
12	Celseq1[12]	Human pancreas	CEL-seq	13	1004	34363
13	Celseq2[12]	Human pancreas	CEL-seq	13	2285	23341
14	QS_Diaphragm[13]	Diaphragm	Smart-seq2	5	870	23341
15	QS_Limb_Muscle[13]	Limb Muscle	Smart-seq2	6	1090	23341

Table S2. Clustering performance comparison of the different clustering algorithms on 15 real scRNA-seq datasets measured by ARI.

Dataset	scAClc	scMGCA	scTAG	scDFN	scDeepCluster	scSCC	scVI
10X_PBMC	0.7813	0.6704	0.6698	0.7672	0.7415	0.7057	0.6956
Human_p	0.9550	0.9024	0.6954	0.6298	0.5867	0.7406	0.7019
Mammary_Gland	0.9378	0.8910	0.8054	0.8861	0.7530	0.8223	0.8371
Klein	0.9108	0.6957	0.7739	0.7560	0.4559	0.7406	0.7253
Adam	0.9349	0.8635	0.8747	0.7411	0.7882	0.7553	0.5231
Chen	0.7880	0.5971	0.5508	0.6420	0.5396	0.3430	0.6362
Turtle_b	0.8926	0.4858	0.5590	0.4128	0.4653	0.7658	0.6873
Baron_human	0.9275	0.3798	0.3649	0.8034	0.4555	0.6786	0.6902
Muraro	0.9288	0.8858	0.8896	0.9280	0.6103	0.9003	0.7615
Plasschaert	0.9125	0.8215	0.8280	0.5337	0.2805	0.7465	0.5896
Pollen	0.8952	0.9267	0.7941	0.9335	0.8642	0.7317	0.1247
Celseq1	0.8274	0.5515	0.5274	0.5686	0.6644	0.4278	0.6302
Celseq2	0.9512	0.8244	0.7956	0.6974	0.6917	0.8617	0.7117
QS_Diaphragm	0.9538	0.9652	0.9153	0.9580	0.6561	0.4645	0.8072
QS_Limb_Muscle	0.9105	0.9751	0.9667	0.9779	0.6925	0.9559	0.7769
Average	0.9005	0.7624	0.7340	0.7490	0.6164	0.7094	0.6599

Table S3. Clustering performance comparison of the different clustering algorithms on 15 real scRNA-seq datasets measured by NMI.

Dataset	scAClc	scMGCA	scTAG	scDFN	scDeepCluster	scSCC	scVI
10X_PBMC	0.7989	0.7365	0.7730	0.7665	0.7704	0.7466	0.6063
Human_p	0.9109	0.8514	0.7525	0.8160	0.7723	0.7177	0.4564
Mammary_Gland	0.9164	0.8512	0.7786	0.8858	0.8210	0.8668	0.8173
Klein	0.9154	0.6978	0.7604	0.7386	0.5604	0.7177	0.7009
Adam	0.9153	0.8595	0.8710	0.8038	0.8238	0.8363	0.4323
Chen	0.7810	0.7651	0.7643	0.7655	0.7474	0.7219	0.2373
Turtle_b	0.8628	0.7064	0.6972	0.7175	0.7411	0.7843	0.4339
Baron_human	0.8818	0.6156	0.5900	0.8541	0.7410	0.8339	0.4511
Muraro	0.8847	0.8379	0.8406	0.8924	0.7569	0.8563	0.6475
Plasschaert	0.8618	0.7602	0.7749	0.6836	0.5452	0.7528	0.3894
Pollen	0.9339	0.9197	0.8925	0.9273	0.9300	0.8067	0.0082
Celseq1	0.8206	0.7590	0.7325	0.7569	0.7778	0.5610	0.3773
Celseq2	0.9175	0.8275	0.7980	0.8350	0.8427	0.8580	0.4714
QS_Diaphragm	0.9126	0.9392	0.8987	0.9448	0.8027	0.6494	0.6768
QS_Limb_Muscle	0.8599	0.9568	0.9448	0.9614	0.8153	0.9190	0.6596
Average	0.8782	0.8056	0.7913	0.8233	0.7632	0.7752	0.4910

Table S4. The number of cells on the Chen dataset and the correspondence between real cell type and cell number.

Cell type	Cluster number	Cell number
Glu1	0	13
Glu2	1	21
Glu3	2	13
Glu4	3	131
Glu5	4	200
Glu6	5	51
Glu7	6	211
Glu8	7	50
Glu9	8	35
Glu10	9	15
Glu11	10	42
Glu12	11	24
Glu13	12	24
Glu14	13	51
Glu15	14	25
GABA1	15	23
GABA2	16	27
GABA3	17	97
GABA4	18	18
GABA5	19	73
GABA6	20	49
GABA7	21	17
GABA8	22	402
GABA9	23	71
GABA10	24	23
GABA11	25	35
GABA12	26	62
GABA13	27	165
GABA14	28	71
GABA15	29	112
GABA16	30	66
GABA17	31	50
GABA18	32	31
Hista	33	17
POPC	34	51
OPC	35	1741
IMO	36	151
MO	37	3541
Astro	38	1148
Ependy	39	413
Tany	40	609
Epith1	41	818
Epith2	42	379
Micro	43	724
Macro	44	167

Table S5. Clustering result comparison of scAClc in different module ((i) only HVGs; (ii) only RFGs; (iii) without the hierarchical gene relevance module; (iv) with the hierarchical gene relevance module (scAClc)) measured by ARI.

Dataset	only HVGs	only RFGs	without	scAClc
10X_PBMC	0.7344	0.7141	0.5766	0.7813
Human_p	0.9554	0.8934	0.8973	0.9550
Mammary_Gland	0.9539	0.6858	0.8556	0.9378
Klein	0.8075	0.9123	0.8278	0.9108
Adam	0.8141	0.7929	0.8034	0.9349
Chen	0.6557	0.7733	0.8511	0.7880
Turtle_b	0.6935	0.7796	0.7663	0.8926
Baron_human	0.6723	0.9197	0.7949	0.9275
Muraro	0.9328	0.9331	0.7290	0.9288
Plasschaert	0.7544	0.7651	0.9361	0.9125
Pollen	0.8567	0.8656	0.3177	0.8952
Celseq1	0.9214	0.8889	0.6171	0.8274
Celseq2	0.9520	0.9256	0.7307	0.9512
QS_Diaphragm	0.9822	0.9519	0.5874	0.9538
QS_Limb_Muscle	0.7026	0.8068	0.6400	0.9105
Average	0.8259	0.8405	0.7287	0.9005

Table S6. Clustering result comparison of scAClc in different module ((i) only HVGs; (ii) only RFGs; (iii) without the hierarchical gene relevance module; (iv) with the hierarchical gene relevance module (scAClc)) measured by NMI.

Dataset	only HVGs	only RFGs	without	scAClc
10X_PBMC	0.7699	0.7656	0.7018	0.7989
Human_p	0.9127	0.8702	0.8777	0.9109
Mammary_Gland	0.9278	0.8281	0.8524	0.9164
Klein	0.8271	0.9160	0.8614	0.9154
Adam	0.8698	0.8484	0.8600	0.9153
Chen	0.6904	0.7652	0.7962	0.7810
Turtle_b	0.7428	0.8305	0.7590	0.8628
Baron_human	0.8284	0.8731	0.8341	0.8818
Muraro	0.8902	0.8904	0.8282	0.8847
Plasschaert	0.7550	0.7714	0.8713	0.8618
Pollen	0.9232	0.9078	0.5801	0.9339
Celseq1	0.8931	0.8774	0.7509	0.8206
Celseq2	0.9216	0.8859	0.8602	0.9175
QS_Diaphragm	0.9686	0.9067	0.8032	0.9126
QS_Limb_Muscle	0.7666	0.7538	0.7853	0.8599
Average	0.8458	0.8460	0.8015	0.8782

Table S7. Clustering result comparison of scAClc in different module ((i) standard contrastive learning; (ii) without contrastive learning; (iii) anchor-centered contrastive learning) measured by ARI.

Dataset	standard	without	anchor-centered
10X_PBMC	0.7649	0.7787	0.7813
Human_p	0.9545	0.9577	0.9550
Mammary_Gland	0.9082	0.9373	0.9378
Klein	0.9123	0.9170	0.9108
Adam	0.8159	0.9389	0.9349
Chen	0.8269	0.5490	0.7880
Turtle_b	0.7640	0.8651	0.8926
Baron_human	0.8023	0.8025	0.9275
Muraro	0.9309	0.9288	0.9288
Plasschaert	0.7872	0.9110	0.9125
Pollen	0.7756	0.8951	0.8952
Celseql	0.6032	0.8250	0.8274
Celseq2	0.9534	0.9520	0.9512
QS_Diaphragm	0.9783	0.9574	0.9538
QS_Limb_Muscle	0.9211	0.9171	0.9105
Average	0.8466	0.8755	0.9005

Table S8. Clustering result comparison of scAClc in different module ((i) standard contrastive learning; (ii) without contrastive learning; (iii) anchor-centered contrastive learning) measured by NMI

Dataset	standard	without	anchor-centered
10X_PBMC	0.7877	0.7966	0.7989
Human_p	0.9095	0.9138	0.9109
Mammary_Gland	0.8727	0.9163	0.9164
Klein	0.9173	0.9260	0.9154
Adam	0.8712	0.9189	0.9153
Chen	0.8028	0.6551	0.7810
Turtle_b	0.8270	0.8345	0.8628
Baron_human	0.8367	0.8364	0.8818
Muraro	0.8864	0.8846	0.8847
Plasschaert	0.7786	0.8594	0.8618
Pollen	0.8960	0.9339	0.9339
Celseql	0.7618	0.9148	0.8206
Celseq2	0.9195	0.9175	0.9175
QS_Diaphragm	0.9528	0.9188	0.9126
QS_Limb_Muscle	0.8641	0.8654	0.8599
Average	0.8589	0.8728	0.8782

Table S9. Clustering result comparison of scAClc in different parameters ((i) The HVG and RFG methods screen 2000 genes respectively; (ii) The HVG and RFG methods screen 3000 genes respectively; (iii) The HVG and RFG methods screen 4000 genes respectively) measured by ARI.

Dataset	2000	3000	4000
10X_PBMC	0.7544	0.7813	0.8089
Human_p	0.9559	0.9550	0.8939
Mammary_Gland	0.8321	0.9378	0.8198
Klein	0.9111	0.9108	0.8222
Adam	0.8129	0.9349	0.8588
Chen	0.7854	0.7880	0.7875
Turtle_b	0.7682	0.8926	0.7717
Baron_human	0.9228	0.9275	0.8028
Muraro	0.9316	0.9288	0.7345
Plasschaert	0.9126	0.9125	0.5971
Pollen	0.7686	0.8952	0.5967
Celseq1	0.8195	0.8274	0.8573
Celseq2	0.7588	0.9512	0.9540
QS_Diaphragm	0.5962	0.9538	0.9682
QS_Limb_Muscle	0.9273	0.9105	0.8408
Average	0.8305	0.9005	0.8076

Table S10. Clustering result comparison of scAClc in different parameters ((i) The HVG and RFG methods screen 2000 genes respectively; (ii) The HVG and RFG methods screen 3000 genes respectively; (iii) The HVG and RFG methods screen 4000 genes respectively) measured by NMI.

Dataset	2000	3000	4000
10X_PBMC	0.7877	0.7989	0.8036
Human_p	0.9121	0.9109	0.8719
Mammary_Gland	0.8714	0.9164	0.8661
Klein	0.9134	0.9154	0.8505
Adam	0.8634	0.9153	0.8763
Chen	0.7736	0.7810	0.7825
Turtle_b	0.7495	0.8628	0.7388
Baron_human	0.8782	0.8818	0.8386
Muraro	0.8882	0.8847	0.8326
Plasschaert	0.8596	0.8618	0.6915
Pollen	0.8834	0.9339	0.8039
Celseq1	0.8132	0.8206	0.8410
Celseq2	0.8669	0.9175	0.9226
QS_Diaphragm	0.7923	0.9126	0.9339
QS_Limb_Muscle	0.8740	0.8599	0.8313
Average	0.8485	0.8782	0.8323

Table S11. Clustering result comparison of scAClc in different parameters ((i) The contrastive-loss weight ranges from 0.10 to 0.06; (ii) The contrastive-loss weight ranges from 0.12 to 0.08; (iii) The contrastive-loss weight ranges from 0.14 to 0.10) measured by ARI.

Dataset	0.10-0.06	0.12-0.08	0.14-0.10
10X_PBMC	0.7745	0.7813	0.7791
Human_p	0.9564	0.9550	0.9559
Mammary_Gland	0.9379	0.9378	0.6676
Klein	0.9150	0.9108	0.9106
Adam	0.9413	0.9349	0.9349
Chen	0.8309	0.7880	0.7753
Turtle_b	0.8310	0.8926	0.7951
Baron_human	0.9263	0.9275	0.9270
Muraro	0.9272	0.9288	0.7288
Plasschaert	0.9115	0.9125	0.6161
Pollen	0.8951	0.8952	0.8951
Celseq1	0.8274	0.8274	0.8274
Celseq2	0.9527	0.9512	0.9527
QS_Diaphragm	0.9538	0.9538	0.5857
QS_Limb_Muscle	0.9025	0.9105	0.9096
Average	0.8989	0.9005	0.8174

Table S12. Clustering result comparison of scAClc in different parameters ((i) The contrastive-loss weight ranges from 0.10 to 0.06; (ii) The contrastive-loss weight ranges from 0.12 to 0.08; (iii) The contrastive-loss weight ranges from 0.14 to 0.10) measured by NMI.

Dataset	0.10-0.06	0.12-0.08	0.14-0.10
10X_PBMC	0.7954	0.7989	0.7949
Human_p	0.9122	0.9109	0.9110
Mammary_Gland	0.9164	0.9164	0.8242
Klein	0.9222	0.9154	0.9150
Adam	0.9217	0.9153	0.9165
Chen	0.8055	0.7810	0.7634
Turtle_b	0.8017	0.8628	0.8249
Baron_human	0.8810	0.8818	0.8814
Muraro	0.8835	0.8847	0.8277
Plasschaert	0.8595	0.8618	0.6991
Pollen	0.9339	0.9339	0.9339
Celseq1	0.8206	0.8206	0.8206
Celseq2	0.9186	0.9175	0.9186
QS_Diaphragm	0.9125	0.9126	0.7686
QS_Limb_Muscle	0.8518	0.8599	0.8618
Average	0.8758	0.8782	0.8441

Reference

- [1] G. X. Zheng, J. M. Terry, P. Belgrader, P. Ryvkin, Z. W. Bent, R. Wilson, S. B. Ziraldo, T. D. Wheeler, G. P. McDermott, J. Zhu et al., “Massively parallel digital transcriptional profiling of single cells,” *Nature communications*, vol. 8, no. 1, p. 14049, 2017.
- [2] M. Baron, A. Veres, S. L. Wolock, A. L. Faust, R. Gaujoux, A. Vetere, J. H. Ryu, B. K. Wagner, S. S. Shen-Orr, A. M. Klein et al., “A single-cell transcriptomic map of the human and mouse pancreas reveals inter-and intra-cell population structure,” *Cell systems*, vol. 3, no. 4, pp. 346–360, 2016.
- [3] T. M. Consortium et al., “Single-cell transcriptomics of 20 mouse organs creates a tabula muris,” *Nature*, vol. 562, no. 7727, pp. 367– 372, 2018.
- [4] A. M. Klein, L. Mazutis, I. Akartuna, N. Tallapragada, A. Veres, V. Li, L. Peshkin, D. A. Weitz, and M. W. Kirschner, “Droplet barcoding for single-cell transcriptomics applied to embryonic stem cells,” *Cell*, vol. 161, no. 5, pp. 1187–1201, 2015.
- [5] M. Adam, A. S. Potter, and S. S. Potter, “Psychrophilic proteases dramatically reduce single-cell rna-seq artifacts: a molecular atlas of kidney development,” *Development*, vol. 144, no. 19, pp. 3625–3632, 2017.
- [6] R. Chen, X. Wu, L. Jiang, and Y. Zhang, “Single-cell rna-seq reveals hypothalamic cell diversity,” *Cell reports*, vol. 18, no. 13, pp. 3227–3241, 2017.
- [7] M. A. Tosches, T. M. Yamawaki, R. K. Naumann, A. A. Jacobi, G. Tushev, and G. Laurent, “Evolution of pallium, hippocampus, and cortical cell types revealed by single-cell transcriptomics in reptiles,” *Science*, vol. 360, no. 6391, pp. 881– 888, 2018.
- [8] Baron M, Veres A, Wolock S L, et al. A single-cell transcriptomic map of the human and mouse pancreas reveals inter-and intra-cell population structure[J]. *Cell systems*, 2016, 3(4): 346-360. e4.
- [9] M. J. Muraro, G. Dharmadhikari, D. Grün, N. Groen, T. Dielen, E. Jansen, L. Van Gurp, M. A. Engelse, F. Carlotti, E. J. De Koning et al., “A single-cell transcriptome atlas of the human pancreas,” *Cell systems*, vol. 3, no. 4, pp. 385– 394, 2016.
- [10] Plasschaert LW, Zilionis R, Choo-Wing R, Savova V, Knehr J, Roma G, et al. A single-cell atlas of the airway epithelium reveals the CFTR-rich pulmonary ionocyte. *Nature* 2018; 560(7718): 377.
- [11] A. A. Pollen, T. J. Nowakowski, J. Shuga, X. Wang, A. A. Leyrat, J. H. Lui, N. Li, L. Szpankowski, B. Fowler, P. Chen et al., “Low-coverage single-cell mrna sequencing reveals cellular heterogeneity and activated signaling pathways in developing cerebral cortex,” *Nature biotechnology*, vol. 32, no. 10, pp. 1053– 1058, 2014.
- [12] M. Baron, A. Veres, S. L. Wolock, A. L. Faust, R. Gaujoux, A. Vetere, J. H. Ryu, B. K. Wagner, S. S. Shen-Orr, A. M. Klein et al., “A single-cell transcriptomic map of the human and mouse pancreas reveals inter-and intra-cell population structure,” *Cell systems*, vol. 3, no. 4, pp. 346–360, 2016.
- [13] Schaum N, Karkanias J, Neff NF, May AP, Quake SR, Wyss-Coray T, et al. Single cell transcriptomics of 20 mouse organs creates a Tabula Muris: The Tabula Muris Consortium. *Nature* 2018; 562(7727): 367.