

Problem 1.

Derive Equation 6.3. [Hint: What fraction of the original sample decays between t and $t + dt$? What, is the (initial) probability, $p(t)dt$, of any given particle decaying between t and $t + dt$? The average lifetime is $\int_0^\infty tp(t) dt$.]

Solution

$$\tau = \frac{1}{\Gamma}, \text{ Equation (6.3)}$$

Q. What fraction of the original sample decays between t and $t + dt$?

A. $dN = -\Gamma N dt$, Equation (6.1)

Q. What is the initial probability, $p(t)dt$?

A. $N(t) = N(0)e^{-\Gamma t}$, Equation (6.2)

Using probability density function, (from wiki),

$$1 = C \int_0^\infty p(t) dt = C \int_0^\infty N(0)e^{-\Gamma t} dt, \text{ where } C \text{ is normalizing coefficient.}$$

$$= C \cdot \frac{N_0}{\Gamma}$$

$$C = \frac{\Gamma}{N_0}$$

Q. The average lifetime?

A. $\tau = \langle t \rangle = \int_0^\infty tp(t) dt = \int_0^\infty t \frac{\Gamma}{N_0} N(0)e^{-\Gamma t} dt = \Gamma \int_0^\infty te^{-\Gamma t} dt$

$$\Gamma \int_0^\infty te^{-\Gamma t} dt = (\text{integration by parts}) = \frac{1}{\Gamma} = \tau$$

reference [wiki - Exponential decay, Probability Density function, integration by parts]